



EDEXCEL STATISTICS S1 (6683) - NOVEMBER 2002
PROVISIONAL MARK SCHEME

Question Number	Scheme	Marks
1.	(a) Statistical models allow problems to be solved without the need to construct a full-scale physical model, saving time/expense. They allow parameters to be changed and refinements to be made quickly. (b) (i) Normal; (ii) Discrete uniform	B2, 1, 0 (2) B1, B1 (2) (4 marks)
2.	(a) 60A, 40S, 2M $P(\text{all only arts}) = \frac{60}{125} \times \frac{59}{124} \times \frac{58}{123} = \frac{3422}{31775} = 0.10769.....$ (b) $P(\text{exactly one only science}) = 3 \times \frac{40}{125} \times \frac{85}{124} \times \frac{84}{123}$ $= \frac{2856}{6355} = 0.44940.....$	B1 M1 A1 A1 (4) B1 M1 A1 (3) (7 marks)
3.	(a) $P(A \cap B) = P(A)P(B) = 0.25 \times 0.30 = 0.075$ (b) $P(A \cup B) = P(A) + P(B) - P(A \cap B) = 0.25 + 0.30 - 0.075 = 0.475$ (c) $P(A B') = \frac{P(A \cap B')}{P(B')} = \frac{P(A) - P(A \cap B)}{1 - P(B)}$ $= \frac{0.25 - 0.075}{1 - 0.3}$ $= 0.25$	M1 A1 (2) M1 A1 (2) M1 M1 A1ft A1 (4) (8 marks)

EDEXCEL STATISTICS S1 (6683) - Nov 2002
PROVISIONAL MARK SCHEME

Question Number	Scheme	Marks
4. (a)	 $P(L > 50.98) = 0.025$ $P\left(Z > \frac{50.98 - \mu}{0.5}\right) = 0.025$ $\therefore \frac{50.98 - \mu}{0.5} = 1.96$ $\therefore \mu = 50 \quad (*)$ $L \sim N(50, 0.5^2)$ (b) $P(49.25 < L < 50.75) = P\left(\frac{49.25 - 50}{0.5} < Z < \frac{50.75 - 50}{0.5}\right)$ $= P(-1.5 < Z < 1.5)$ -1.5 & +1.5 $= 2\Phi(1.5) - 1$ $= 0.8664$ (c) $P(\text{Both}) = (1 - 0.8664)^2$ $= 0.01784\dots$	B1 M1 A1 M1 A1 (5) M1 A1 M1 A1 (4) M1 A1 (2) (11 marks)
5. (a)	$S_{ss} = 108.07875; S_{st} = 129.1675$ $q = \frac{S_{st}}{S_{ss}} = \frac{129.1675}{108.07875} = 1.1951239\dots$ $p = \frac{65.0}{8} - (1.1951239\dots) \times \frac{48.5}{8} = 0.879561\dots$ $\therefore t = 0.879561\dots + 1.1951239\dots S$ (b) $y - 20 = 0.879561\dots + 1.1951239\dots(x - 6)$ $\therefore y = 13.709 + 1.195x$ (c) 0.943; the pmcc is an index (no units) and is not affected by linear transformations of either/both variables	B1; B1 M1, A1 M1, A1 A1 ft (7) M1, A1 ft A1 (3) B1; B1 (2) (12 marks)

EDEXCEL STATISTICS S1 (6683) - NOVEMBER 2002
PROVISIONAL MARK SCHEME

Question Number	Scheme	Marks
6.	(a) $\alpha + \beta = 0.5$ $-2\alpha + 2\beta = -0.2$ $\therefore \alpha = 0.3, \beta = 0.2$ (b) $F(0.8) = 0.6$ (c) $E(X^2) = (4 \times 0.3) + \dots + (4 \times 0.2), = 2.4$ $\therefore \text{Var}(X) = 2.4 - (-0.2)^2, = 2.36$ (d) $E(3X - 2) = 3E(X) - 2, = -2.6$ (e) $\text{Var}(2X + 6) = 4 \text{Var}(X), = 9.44$	B1 M1 M1 A1; A1 (6) B1 ft (1) M1, A1 M1, A1 (4) M1, A1 ft (2) M1, A1 ft (2) (15 marks)
7.	(a) Mode = 78 (b) $Q_1 = 56; Q_2 = 70; Q_3 = 78$ (c) $(Q_3 - Q_1) = 22$ $Q_1 - 1.0(Q_3 - Q_1) = 34 \Rightarrow 31$ & 31 are outliers $Q_3 + 1.0(Q_3 - Q_1) = 100 \Rightarrow$ no outliers (d) <i>(accurate sketch on graph paper required)</i>  boxplot scales and labels Q_1, Q_2, Q_3 31, 32, 34 (39), 92 Aptitude score 30 40 50 60 70 80 90 100 (e) $\mu = \frac{3363}{50} = 67.26$ $\sigma^2 = \frac{238305}{50} - (67.26)^2 = 242.1924$ $\therefore \sigma = \sqrt{242.1924} = 15.56253\dots$ (f) $(Q_3 - Q_2) < (Q_2 - Q_1)$, i.e. $8 < 14 \Rightarrow$ negative skew Mean < Median < Mode, i.e. $67.26 < 70 < 78 \Rightarrow$ negative skew	B1 (1) B1; B1; B1 (3) M1 A1 A1 (3) M1 B1 A1 A1 (4) B1 M1 A1 (3) M1, A1 M1, A1 (4) (18 marks)