

4727 Further Pure Mathematics 3

<p>1 (a) (i) e.g. $ap \neq pa \Rightarrow$ not commutative</p>	<p>B1 1</p>	<p>For correct reason and conclusion</p>
<p>(ii) 3</p>	<p>B1 1</p>	<p>For correct number</p>
<p>(iii) e, a, b</p>	<p>B1 1</p>	<p>For correct elements</p>
<p>(b) c^3 has order 2 c^4 has order 3 c^5 has order 6</p>	<p>B1 B1 B1 3 6</p>	<p>For correct order For correct order For correct order</p>
<p>2 $m^2 - 8m + 16 = 0$ $\Rightarrow m = 4$ \Rightarrow CF ($y =$) $(A + Bx)e^{4x}$ For PI try $y = px + q$ $\Rightarrow -8p + 16(px + q) = 4x$ $\Rightarrow p = \frac{1}{4} \quad q = \frac{1}{8}$ \Rightarrow GS $y = (A + Bx)e^{4x} + \frac{1}{4}x + \frac{1}{8}$</p>	<p>M1 A1 A1√ M1 A1 A1 B1√ 7 7</p>	<p>For stating and attempting to solve auxiliary eqn For correct solution For CF of correct form. f.t. from m For using linear expression for PI For correct coefficients For GS = CF + PI. Requires $y =$. f.t. from CF and PI with 2 arbitrary constants in CF and none in PI</p>
<p>3 (i) line segment OA</p>	<p>B1 B1 2</p>	<p>For stating line through O OR A For correct description AEF</p>
<p>(ii) $(\mathbf{r} - \mathbf{a}) \times (\mathbf{r} - \mathbf{b}) = \vec{AP} \times \vec{BP}$ $= \mathbf{AP} \mathbf{BP} \sin \pi \cdot \hat{\mathbf{n}} = \mathbf{0}$</p>	<p>B1 B1 2</p>	<p>For identifying $\mathbf{r} - \mathbf{a}$ with \vec{AP} and $\mathbf{r} - \mathbf{b}$ with \vec{BP} Allow direction errors For using \times of 2 parallel vectors = 0 OR $\sin \pi = 0$ or $\sin 0 = 0$ in an appropriate vector expression</p>
<p>(iii) line through O parallel to AB</p>	<p>B1 B1 B1 3 7</p>	<p>For stating line For stating through O For stating correct direction SR For \vec{AB} or \vec{BA} allow B1 B0 B1</p>
<p>4 $(C + iS) = \int_0^{\frac{1}{2}\pi} e^{2x} (\cos 3x + i \sin 3x) (dx)$ $\cos 3x + i \sin 3x = e^{3ix}$ $\int_0^{\frac{1}{2}\pi} e^{(2+3i)x} (dx) = \frac{1}{2+3i} \left[e^{(2+3i)x} \right]_0^{\frac{1}{2}\pi}$ $= \frac{2-3i}{4+9} \left(e^{(2+3i)\frac{1}{2}\pi} - e^0 \right) = \frac{2-3i}{13} (-ie^\pi - 1)$ $= \left\{ \frac{1}{13} (-2 - 3e^\pi + i(3 - 2e^\pi)) \right\}$ $C = -\frac{1}{13} (2 + 3e^\pi)$ $S = \frac{1}{13} (3 - 2e^\pi)$</p>	<p>B1 M1* A1 A1 M1 (dep*) M1 (dep*) A1 A1 8</p>	<p>For using de Moivre, seen or implied For writing as a single integral in exp form For correct integration (ignore limits) For substituting limits correctly (unsimplified) (may be earned at any stage) For multiplying by complex conjugate of $2+3i$ For equating real and/or imaginary parts For correct expression AG For correct expression</p>

<p>5 (i) IF $e^{\int \frac{1}{x} dx} = e^{\ln x} = x$ OR $x \frac{dy}{dx} + y = x \sin 2x$ $\Rightarrow \frac{d}{dx}(xy) = x \sin 2x$ $\Rightarrow xy = \int x \sin 2x (dx)$ $xy = -\frac{1}{2} x \cos 2x + \frac{1}{2} \int \cos 2x (dx)$ $xy = -\frac{1}{2} x \cos 2x + \frac{1}{4} \sin 2x (+c)$ $\Rightarrow y = -\frac{1}{2} \cos 2x + \frac{1}{4x} \sin 2x + \frac{c}{x}$</p>	<p>M1 A1 M1 A1 M1 A1 A1 6</p>	<p>For correct process for finding integrating factor OR for multiplying equation through by x For writing DE in this form (may be implied) For integration by parts the correct way round For 1st term correct For their 1st term and attempt at integration of $\frac{\cos}{\sin} kx$ For correct expression for y</p>
<p>(ii) $(\frac{1}{4}\pi, \frac{2}{\pi}) \Rightarrow \frac{2}{\pi} = \frac{1}{\pi} + \frac{4c}{\pi} \Rightarrow c = \frac{1}{4}$ $\Rightarrow y = -\frac{1}{2} \cos 2x + \frac{1}{4x} \sin 2x + \frac{1}{4x}$</p>	<p>M1 A1 2</p>	<p>For substituting $(\frac{1}{4}\pi, \frac{2}{\pi})$ in solution For correct solution. Requires $\boxed{y =}$.</p>
<p>(iii) $(y \approx) -\frac{1}{2} \cos 2x$</p>	<p>B1√ 1 9</p>	<p>For correct function AEF f.t. from (ii)</p>
<p>6 (i) METHOD 1 State $B = (-1, -7, 2) + t(1, 2, -2)$ On plane $\Rightarrow (-1+t) + 2(-7+2t) - 2(2-2t) = -1$ $\Rightarrow t = 2 \Rightarrow B = (1, -3, -2)$ $AB = \sqrt{2^2 + 4^2 + 4^2}$ OR $2\sqrt{1^2 + 2^2 + 2^2} = 6$</p>	<p>M1 M1 M1 A1 A1 5</p>	<p>Either coordinates or vectors may be used Methods 1 and 2 may be combined, for a maximum of 5 marks For using vector normal to plane For substituting parametric form into plane For solving a linear equation in t For correct coordinates For correct length of AB</p>
<p>METHOD 2 $AB = \frac{ -1-14-4+1 }{\sqrt{1^2+2^2+2^2}} = 6$ OR $AB = \mathbf{AC} \cdot \frac{\mathbf{AB}}{ \mathbf{AB} } = \frac{[6, 7, 1] \cdot [1, 2, -2]}{\sqrt{1^2+2^2+2^2}} = 6$ $B = (-1, -7, 2) \pm 6 \frac{(1, 2, -2)}{\sqrt{1^2+2^2+2^2}}$ $B = (-1, -7, 2) \pm (2, 4, -4)$ $B = (1, -3, -2)$</p>	<p>M1 A1 M1 B1 A1</p>	<p>For using a correct distance formula For correct length of AB For using $B = A + \text{length of } AB \times \text{unit normal}$ For checking whether + or - is needed (substitute into plane equation) For correct coordinates (allow even if B0)</p>
<p>(ii) Find vector product of any two of $\pm[6, 7, 1], \pm[6, -3, 0], \pm(0, 10, 1)$ Obtain $k[1, 2, -20]$ $\theta = \cos^{-1} \frac{ [1, 2, -2] \cdot [1, 2, -20] }{\sqrt{1^2+2^2+2^2} \sqrt{1^2+2^2+20^2}}$ $\theta = \cos^{-1} \frac{45}{\sqrt{9} \sqrt{405}} = 41.8^\circ (41.810\dots^\circ, 0.72972\dots)$</p>	<p>M1 A1 M1* M1 (dep*) A1√ A1 6 11</p>	<p>For finding vector product of two relevant vectors For correct vector \mathbf{n} For using scalar product of two normal vectors For stating both moduli in denominator For correct scalar product. f.t. from \mathbf{n} For correct angle</p>

<p>8 (i) Group A: $e = 6$ Group B: $e = 1$ Group C: $e = 2^0$ OR 1 Group D: $e = 1$</p>	$\left. \begin{array}{l} \text{B1} \\ \text{B1} \\ \mathbf{2} \end{array} \right\}$	<p>For any two correct identities For two other correct identities AEF for D, but not “$m = n$”</p>
<p>(ii) EITHER OR</p> <p>A 2 4 6 8 2 4 8 2 6 orders of elements 4 8 6 4 2 1, 2, 4, 4 6 2 4 6 8 OR cyclic group 8 6 2 8 4</p> <p>B 1 5 7 11 1 1 5 7 11 orders of elements 5 5 1 11 7 1, 2, 2, 2 7 7 11 1 5 OR non-cyclic group 11 11 7 5 1 OR Klein group</p> <p>C 2^0 2^1 2^2 2^3 2^0 2^0 2^1 2^2 2^3 orders of elements 2^1 2^1 2^2 2^3 2^0 1, 2, 4, 4 2^2 2^2 2^3 2^0 2^1 OR cyclic group 2^3 2^3 2^0 2^1 2^2</p> <p>$A \not\cong B$ $B \not\cong C$ $A \cong C$</p>	<p>B1* B1* B1 (dep*) B1 (dep*) B1 (dep*) 5</p>	<p>For showing group table OR sufficient details of orders of elements OR stating cyclic / non-cyclic / Klein group (as appropriate)</p> <p>for one of groups A, B, C for another of groups A, B, C</p> <p>For stating non-isomorphic } with sufficient detail For stating non-isomorphic } relating to the first 2 marks For stating isomorphic }</p>
<p>(iii) $\frac{1+2m}{1+2n} \times \frac{1+2p}{1+2q} = \frac{1+2m+2p+4mp}{1+2n+2q+4nq}$</p> <p>$= \frac{1+2(m+p+2mp)}{1+2(n+q+2nq)} \equiv \frac{1+2r}{1+2s}$</p>	<p>M1* M1 (dep*) A1 A1 4</p>	<p>For considering product of 2 distinct elements of this form For multiplying out For simplifying to form shown For identifying as correct form, so closed</p> <p>SR $\frac{\text{odd}}{\text{odd}} \times \frac{\text{odd}}{\text{odd}} = \frac{\text{odd}}{\text{odd}}$ earns full credit SR If clearly attempting to prove commutativity, allow at most M1</p>
<p>(iv) Closure not satisfied Identity and inverse not satisfied</p>	<p>B1 B1 2 13</p>	<p>For stating closure For stating identity and inverse SR If associativity is stated as not satisfied, then award at most B1 B0 OR B0 B1</p>