

Centre No.						Paper Reference					Surname	Initial(s)	
Candidate No.						6	6	8	1	/	0	1	Signature

Paper Reference(s)
6681/01

Edexcel GCE

Mechanics M5

Advanced/Advanced Subsidiary

Monday 28 June 2010 – Afternoon

Time: 1 hour 30 minutes

Examiner's use only

--	--	--

Team Leader's use only

--	--	--

Question Number	Leave Blank
1	
2	
3	
4	
5	
6	
Total	

<u>Materials required for examination</u>	<u>Items included with question papers</u>
Mathematical Formulae (Pink)	Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper. Answer ALL the questions. You must write your answer to each question in the space following the question. Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$. When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided. Full marks may be obtained for answers to ALL questions. The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2). There are 6 questions in this question paper. The total mark for this paper is 75. There are 24 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled. You should show sufficient working to make your methods clear to the Examiner. Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Edexcel Limited copyright policy. ©2010 Edexcel Limited.

Printer's Log. No.
N35394A
W850/R6681/57570 4/5/5/3



Turn over

2.




Figure 1

A uniform circular disc has mass $4m$, centre O and radius $4a$. The line POQ is a diameter of the disc. A circular hole of radius $2a$ is made in the disc with the centre of the hole at the point R on PQ where $QR = 5a$, as shown in Figure 1.

The resulting lamina is free to rotate about a fixed smooth horizontal axis L which passes through Q and is perpendicular to the plane of the lamina.

(a) Show that the moment of inertia of the lamina about L is $69ma^2$. (7)

The lamina is hanging at rest with P vertically below Q when it is given an angular velocity Ω . Given that the lamina turns through an angle $\frac{2\pi}{3}$ before it first comes to instantaneous rest,

(b) find Ω in terms of g and a . (6)

3. A uniform lamina ABC of mass m is in the shape of an isosceles triangle with $AB = AC = 5a$ and $BC = 8a$.

(a) Show, using integration, that the moment of inertia of the lamina about an axis through A , parallel to BC , is $\frac{9}{2}ma^2$.

(6)

The foot of the perpendicular from A to BC is D . The lamina is free to rotate in a vertical plane about a fixed smooth horizontal axis which passes through D and is perpendicular to the plane of the lamina. The lamina is released from rest when DA makes an angle α with the downward vertical. It is given that the moment of inertia of the lamina about an axis through A , perpendicular to BC and in the plane of the lamina, is $\frac{8}{3}ma^2$.

(b) Find the angular acceleration of the lamina when DA makes an angle θ with the downward vertical.

(8)

Given that α is small,

(c) find an approximate value for the period of oscillation of the lamina about the vertical.

(2)



4. Two forces $\mathbf{F}_1 = (\mathbf{i} + 2\mathbf{j} + 3\mathbf{k})$ N and $\mathbf{F}_2 = (3\mathbf{i} + \mathbf{j} + 2\mathbf{k})$ N act on a rigid body. The force \mathbf{F}_1 acts through the point with position vector $(2\mathbf{i} + \mathbf{k})$ m and the force \mathbf{F}_2 acts through the point with position vector $(\mathbf{j} + 2\mathbf{k})$ m.

(a) If the two forces are equivalent to a single force \mathbf{R} , find

(i) \mathbf{R} , (2)

(ii) a vector equation of the line of action of \mathbf{R} , in the form $\mathbf{r} = \mathbf{a} + \lambda\mathbf{b}$. (6)

(b) If the two forces are equivalent to a single force acting through the point with position vector $(\mathbf{i} + 2\mathbf{j} + \mathbf{k})$ m together with a couple of moment \mathbf{G} , find the magnitude of \mathbf{G} . (5)



Leave
blank

Question 4 continued

Lined area for writing the answer to Question 4.



Leave
blank

Question 5 continued

A series of horizontal lines for writing, consisting of 30 lines.



