



General Certificate of Education  
Advanced Level Examination  
June 2013

## Mathematics

## MFP3

### Unit Further Pure 3

Monday 10 June 2013 9.00 am to 10.30 am

**For this paper you must have:**

- the blue AQA booklet of formulae and statistical tables.  
You may use a graphics calculator.

**Time allowed**

- 1 hour 30 minutes

**Instructions**

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- Write the question part reference (eg (a), (b)(i) etc) in the left-hand margin.
- You must answer each question in the space provided for that question. If you require extra space, use an AQA supplementary answer book; do **not** use the space provided for a different question.
- Do not write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.

**Information**

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 75.

**Advice**

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.

## 2

- 1 It is given that  $y(x)$  satisfies the differential equation

$$\frac{dy}{dx} = f(x, y)$$

where  $f(x, y) = (x - y)\sqrt{x + y}$

and  $y(2) = 1$

Use the improved Euler formula

$$y_{r+1} = y_r + \frac{1}{2}(k_1 + k_2)$$

where  $k_1 = hf(x_r, y_r)$  and  $k_2 = hf(x_r + h, y_r + k_1)$  and  $h = 0.2$ , to obtain an approximation to  $y(2.2)$ , giving your answer to three decimal places. (5 marks)

---

- 2 The Cartesian equation of a circle is  $(x + 8)^2 + (y - 6)^2 = 100$ .

Using the origin  $O$  as the pole and the positive  $x$ -axis as the initial line, find the polar equation of this circle, giving your answer in the form  $r = p \sin \theta + q \cos \theta$ . (4 marks)

---

- 3 (a) Find the values of the constants  $a$ ,  $b$  and  $c$  for which  $a + bx + cxe^{-3x}$  is a particular integral of the differential equation

$$\frac{d^2y}{dx^2} + 2\frac{dy}{dx} - 3y = 3x - 8e^{-3x} \quad (5 \text{ marks})$$

- (b) Hence find the general solution of this differential equation. (3 marks)

- (c) Hence express  $y$  in terms of  $x$ , given that  $y = 1$  when  $x = 0$  and that  $\frac{dy}{dx} \rightarrow -1$  as  $x \rightarrow \infty$ . (4 marks)
- 

- 4 Evaluate the improper integral

$$\int_0^{\infty} \left( \frac{2x}{x^2 + 4} - \frac{4}{2x + 3} \right) dx$$

showing the limiting process used and giving your answer in the form  $\ln k$ , where  $k$  is a constant. (6 marks)



**5 (a)** Differentiate  $\ln(\ln x)$  with respect to  $x$ . (1 mark)

**(b) (i)** Show that  $\ln x$  is an integrating factor for the first-order differential equation

$$\frac{dy}{dx} + \frac{1}{x \ln x} y = 9x^2, \quad x > 1 \quad (2 \text{ marks})$$

**(ii)** Hence find the solution of this differential equation, given that  $y = 4e^3$  when  $x = e$ . (6 marks)

---

**6** It is given that  $y = (4 + \sin x)^{\frac{1}{2}}$ .

**(a)** Express  $y \frac{dy}{dx}$  in terms of  $\cos x$ . (2 marks)

**(b)** Find the value of  $\frac{d^3y}{dx^3}$  when  $x = 0$ . (5 marks)

**(c)** Hence, by using Maclaurin's theorem, find the first four terms in the expansion, in ascending powers of  $x$ , of  $(4 + \sin x)^{\frac{1}{2}}$ . (2 marks)

---

**7** A differential equation is given by

$$\sin^2 x \frac{d^2y}{dx^2} - 2 \sin x \cos x \frac{dy}{dx} + 2y = 2 \sin^4 x \cos x, \quad 0 < x < \pi$$

**(a)** Show that the substitution

$$y = u \sin x$$

where  $u$  is a function of  $x$ , transforms this differential equation into

$$\frac{d^2u}{dx^2} + u = \sin 2x \quad (5 \text{ marks})$$

**(b)** Hence find the general solution of the differential equation


$$\sin^2 x \frac{d^2y}{dx^2} - 2 \sin x \cos x \frac{dy}{dx} + 2y = 2 \sin^4 x \cos x$$

giving your answer in the form  $y = f(x)$ . (6 marks)

Turn over ►



- 8 The diagram shows a sketch of a curve and a circle.



The polar equation of the curve is

$$r = 3 + 2 \sin \theta, \quad 0 \leq \theta \leq 2\pi$$

The circle, whose polar equation is  $r = 2$ , intersects the curve at the points  $P$  and  $Q$ , as shown in the diagram.

- (a) Find the polar coordinates of  $P$  and the polar coordinates of  $Q$ . (3 marks)
- (b) A straight line, drawn from the point  $P$  through the pole  $O$ , intersects the curve again at the point  $A$ .
- (i) Find the polar coordinates of  $A$ . (2 marks)
- (ii) Find, in surd form, the length of  $AQ$ . (3 marks)
- (iii) Hence, or otherwise, explain why the line  $AQ$  is a tangent to the circle  $r = 2$ . (2 marks)
- (c) Find the area of the shaded region which lies inside the circle  $r = 2$  but outside the curve  $r = 3 + 2 \sin \theta$ . Give your answer in the form  $\frac{1}{6}(m\sqrt{3} + n\pi)$ , where  $m$  and  $n$  are integers. (9 marks)

