



GCE

Mathematics

Advanced GCE

Unit 4727: Further Pure Mathematics 3

Mark Scheme for January 2011

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of pupils of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, OCR Nationals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by Examiners. It does not indicate the details of the discussions which took place at an Examiners' meeting before marking commenced.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the Report on the Examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.



© OCR 2011

Any enquiries about publications should be addressed to:

OCR Publications
PO Box 5050
Annesley
NOTTINGHAM
NG15 0DL

Telephone: 0870 770 6622
Facsimile: 01223 552610
E-mail: publications@ocr.org.uk

1 (i)	Integrating factor. $e^{\int x dx} = e^{\frac{1}{2}x^2}$	B1	For correct IF
	$\Rightarrow \frac{d}{dx} \left(y e^{\frac{1}{2}x^2} \right) = x e^{x^2}$	M1	For $\frac{d}{dx} (y \cdot \text{their IF}) = x e^{\frac{1}{2}x^2} \cdot \text{their IF}$
	$\Rightarrow y e^{\frac{1}{2}x^2} = \frac{1}{2} e^{x^2} (+c)$	A1	For correct integration both sides
	$\Rightarrow y = e^{-\frac{1}{2}x^2} \left(\frac{1}{2} e^{x^2} + c \right) = \frac{1}{2} e^{\frac{1}{2}x^2} + c e^{-\frac{1}{2}x^2}$	A1 4	For correct solution AEF as $y = f(x)$
(ii)	$(0, 1) \Rightarrow c = \frac{1}{2}$	M1	For substituting (0, 1) into their GS, solving for c and obtaining a solution of the DE
	$\Rightarrow y = \frac{1}{2} \left(e^{\frac{1}{2}x^2} + e^{-\frac{1}{2}x^2} \right)$	A1 2	For correct solution AEF Allow $y = \cosh\left(\frac{1}{2}x^2\right)$
6			
2 (i)	$\mathbf{n} = [2, 1, -3] \times [-1, 2, 4]$	M1	For using \times of direction vectors
	$= [10, -5, 5] = k[2, -1, 1]$	A1	For correct \mathbf{n}
	$(1, 3, 4) \Rightarrow 2x - y + z = 3$	A1 3	For substituting (1, 3, 4) and obtaining AG (Verification only M0)
(ii)	METHOD 1	M1	For $21 - 3$ OR $[1, 3, 4] \cdot [2, -1, 1] - 21$
	distance = $\frac{21-3}{ \mathbf{n} }$ OR $\frac{[1, 3, 4] \cdot [2, -1, 1] - 21}{ \mathbf{n} }$		OR $ (1, 3, 4) - [a, b, c] \cdot [2, -1, 1] $ soi
	OR $\frac{ (1, 3, 4) - [a, b, c] \cdot [2, -1, 1] }{ \mathbf{n} }$ where (a, b, c) is on q	B1	For $ \mathbf{n} = \sqrt{6}$ soi
	$= \frac{18}{\sqrt{6}} = 3\sqrt{6}$	A1 3	For correct distance AEF
METHOD 2	$[1+2t, 3-t, 4+t]$ on q	M1	For forming and solving an equation in t
	$\Rightarrow 2(1+2t) - (3-t) + (4+t) = 21 \Rightarrow t = 3$	B1	For $ \mathbf{n} = \sqrt{6}$ soi
	$\Rightarrow \text{distance} = 3 \mathbf{n} = 3\sqrt{6}$	A1	For correct distance AEF
METHOD 3	As Method 2 to $t = 3 \Rightarrow (7, 0, 7)$ on q	M1*	For finding point where normal meets q
	distance from (1, 3, 4)	M1	For finding distance from (1, 3, 4)
	$= \sqrt{(7-1)^2 + (0-3)^2 + (7-4)^2} = \sqrt{54} = 3\sqrt{6}$	(*dep) A1	For correct distance AEF
6			
3 (i)	$\sin \theta = \frac{1}{2i} (e^{i\theta} - e^{-i\theta})$	B1	z or $e^{i\theta}$ may be used throughout For correct expression for $\sin \theta$ soi
	$\sin^4 \theta = \frac{1}{16} (z^4 - 4z^2 + 6 - 4z^{-2} + z^{-4})$	M1	For expanding $(e^{i\theta} - e^{-i\theta})^4$ (with at least 3 terms and 1 binomial coefficient)
	$\Rightarrow \sin^4 \theta = \frac{1}{16} (2 \cos 4\theta - 8 \cos 2\theta + 6)$	M1	For grouping terms and using multiple angles
	$\Rightarrow \sin^4 \theta = \frac{1}{8} (\cos 4\theta - 4 \cos 2\theta + 3)$	A1 4	For answer obtained correctly AG
(ii)	$\int_0^{\frac{1}{6}\pi} \sin^4 \theta d\theta = \frac{1}{8} \left[\frac{1}{4} \sin 4\theta - 2 \sin 2\theta + 3\theta \right]_0^{\frac{1}{6}\pi}$	M1	For integrating (i) to $A \sin 4\theta + B \sin 2\theta + C\theta$
		A1	For correct integration
	$= \frac{1}{8} \left(\frac{1}{8} \sqrt{3} - \sqrt{3} + \frac{1}{2} \pi \right) = \frac{1}{64} (4\pi - 7\sqrt{3})$	M1	For completing integration and substituting limits
		A1 4	For correct answer AEF (exact)
8			

4 (i)	<p><i>EITHER</i> $1 + \omega + \omega^2$ $=$ sum of roots of $(z^3 - 1) = 0$</p> <hr/> <p><i>OR</i> $\omega^3 = 1 \Rightarrow (\omega - 1)(\omega^2 + \omega + 1) = 0$ $\Rightarrow 1 + \omega + \omega^2 = 0$ (for $\omega \neq 1$)</p> <hr/> <p><i>OR</i> sum of G.P. $1 + \omega + \omega^2 = \frac{1 - \omega^3}{1 - \omega} \left(= \frac{0}{1 - \omega} \right) = 0$</p> <hr/> <p><i>OR</i>  shown on Argand diagram or explained in terms of vectors</p> <hr/> <p><i>OR</i> $1 + \text{cis } \frac{2}{3}\pi + \text{cis } \frac{4}{3}\pi = 1 + \left(-\frac{1}{2} + \frac{\sqrt{3}}{2}i\right) + \left(-\frac{1}{2} - \frac{\sqrt{3}}{2}i\right) = 0$</p>	M1 A1 2	For result shown by any correct method AG
(ii)	<p>Multiplication by $\omega \Rightarrow$ rotation through $\frac{2}{3}\pi$ </p> <p>$z_1 - z_3 = \vec{CA}$, $z_3 - z_2 = \vec{BC}$</p> <p>\vec{BC} rotates through $\frac{2}{3}\pi$ to direction of \vec{CA}</p> <p>ΔABC has $BC = CA$, hence result</p>	B1 B1 M1 A1 4	<p>For correct interpretation of \times by ω (allow 120° and omission of, or error in, \odot)</p> <p>For identification of vectors soi (ignore direction errors)</p> <p>For linking BC and CA by rotation of $\frac{2}{3}\pi$ <i>OR</i> ω</p> <p>For stating equal magnitudes \Rightarrow AG</p>
(iii)	<p>(ii) $\Rightarrow z_1 + \omega z_2 - (1 + \omega)z_3 = 0$</p> <p>$1 + \omega + \omega^2 = 0 \Rightarrow z_1 + \omega z_2 + \omega^2 z_3 = 0$</p>	M1 A1 2	<p>For using $1 + \omega + \omega^2 = 0$ in (ii)</p> <p>For obtaining AG</p>
8			
5 (i)	<p>Aux. equation $3m^2 + 5m - 2 (= 0)$</p> <p>$\Rightarrow m = \frac{1}{3}, -2$</p> <p>CF ($y =$) $Ae^{\frac{1}{3}x} + Be^{-2x}$</p> <p>PI ($y =$) $px + q \Rightarrow 5p - 2(px + q) = -2x + 13$ $\Rightarrow p = 1, q = -4$</p> <p>GS ($y =$) $Ae^{\frac{1}{3}x} + Be^{-2x} + x - 4$</p>	M1 A1 A1√ M1 A1 A1 B1√ 7	<p>For correct auxiliary equation seen and solution attempted</p> <p>For correct roots</p> <p>For correct CF f.t. from m with 2 arbitrary constants</p> <p>For stating and substituting PI of correct form</p> <p>For correct value of p, and of q</p> <p>For GS f.t. from their CF+PI with 2 arbitrary constants in CF and none in PI</p>
(ii)	<p>$\left(0, -\frac{7}{2}\right) \Rightarrow A + B = \frac{1}{2}$</p> <p>$y' = \frac{1}{3}Ae^{\frac{1}{3}x} - 2Be^{-2x} + 1, (0, 0) \Rightarrow A - 6B = -3$</p> <p>$\Rightarrow A = 0, B = \frac{1}{2}$</p> <p>$\Rightarrow (y =) \frac{1}{2}e^{-2x} + x - 4$</p>	M1 M1 M1 A1 B1√ 5	<p>For substituting $\left(0, -\frac{7}{2}\right)$ in their GS and obtaining an equation in A and B</p> <p>For finding y', substituting $(0, 0)$ and obtaining an equation in A and B</p> <p>For solving their 2 equations in A and B</p> <p>For correct A and B CAO</p> <p>For correct solution f.t. with their A and B in their GS</p>
(iii)	<p>x large $\Rightarrow (y =) x - 4$</p>	B1√ 1	<p>For correct equation or function (allow \approx and \rightarrow) WWW f.t. from (ii) if valid</p>
13			

4727

Mark Scheme

January 2011

6 (i)	$a^4 = r^6 = e \Rightarrow a$ has order 4, a^2 has order 2 $(a^3)^4 = a^{12} = e \Rightarrow a^3$ has order 4 $(r^2)^3 = e \Rightarrow r^2$ has order 3	M1 A1 A1 B1	For considering powers of a For order of any one of a, a^2, a^3 correct For all correct For order of r^2 correct																		
(ii)	<p>G order 4</p> <table border="1" data-bbox="263 414 742 481"> <tr> <td>Order of element</td> <td>1</td> <td>2</td> <td>(4)</td> </tr> <tr> <td>Number of elements</td> <td>1</td> <td>3</td> <td>(0)</td> </tr> </table> <p>H order 6</p> <table border="1" data-bbox="263 504 813 571"> <tr> <td>Order of element</td> <td>1</td> <td>2</td> <td>3</td> <td>(6)</td> </tr> <tr> <td>Number of elements</td> <td>1</td> <td>3</td> <td>2</td> <td>(0)</td> </tr> </table> <p>G and H are the only non-cyclic groups of order which divides 12 Q has 1 element of order 2, G and H have 3, so no non-cyclic subgroups in Q</p>	Order of element	1	2	(4)	Number of elements	1	3	(0)	Order of element	1	2	3	(6)	Number of elements	1	3	2	(0)	M1 A1 A1 B1 B1	For top line in either table Allow inclusion of 4 and 6 respectively (and other orders if 0 appears below) For order 4 table For order 6 table For stating that only G and H need be considered AEF For argument completed by elements of order 2 AG SR Allow equivalent arguments for B1 B1
Order of element	1	2	(4)																		
Number of elements	1	3	(0)																		
Order of element	1	2	3	(6)																	
Number of elements	1	3	2	(0)																	
9																					
7 (i)	$[1, 1, -2] \times [1, -1, 3] = (\pm)[1, -5, -2]$ $[1, -1, 3] \times [1, 5, -12] = (\pm)[-3, 15, 6]$ $[-3, 15, 6] = k[1, -5, -2] \Rightarrow$ parallel	M1 A1 M1 A1 A1	For using \times of direction vectors For correct direction For using \times of direction vectors For correct direction For correct direction For argument completed AG ($k = -3$ not essential)																		
(ii)	Line of intersection is parallel to l and m	B1	1 For correct statement																		
(iii)	<p>METHOD 1</p> $\left. \begin{array}{l} x + y - 2z = 5 \\ x - y + 3z = 6 \end{array} \right\} \text{e.g. } z = 0 \Rightarrow \left(\frac{11}{2}, -\frac{1}{2}, 0\right) \text{ on } l$ $\left. \begin{array}{l} x - y + 3z = 6 \\ x + 5y - 12z = 12 \end{array} \right\} \text{e.g. } z = 0 \Rightarrow (7, 1, 0) \text{ on } m$ $\left. \begin{array}{l} x + y - 2z = 5 \\ x + 5y - 12z = 12 \end{array} \right\} \text{e.g. } z = 0 \Rightarrow \left(\frac{13}{4}, \frac{7}{4}, 0\right) \text{ on } l_3$ <p>Different points \Rightarrow no common line of intersection</p>	M1 A1 A1 A1	For attempt to find points on 2 lines For a correct point on one line For a correct point on another line For correct answer																		
METHOD 2	$\left. \begin{array}{l} x + y - 2z = 5 \\ x - y + 3z = 6 \end{array} \right\} \text{e.g. } \Rightarrow z = 11 - 2x, y = 27 - 5x$ <p>LHS of eqn 3 = $x + (135 - 25x) - (132 - 24x) = 3 \neq 12$ \Rightarrow no common line of intersection</p>	M1 A1 A1	For finding (e.g.) y and z in terms of x OR eliminating one variable For correct expressions OR equations For obtaining a contradiction from 3rd equation For correct answer																		
METHOD 3	<p>LHS $II_3 = 3II_1 - 2II_2$ RHS $3 \times 5 - 2 \times 6 = 3 \neq 12$ \Rightarrow no common line of intersection</p>	M2 A1 A1	For attempt to link 3 equations For obtaining a contradiction For correct answer																		
SR Variations on all methods may gain full credit	SR f.t. may be allowed from relevant working																				
10																					

4727

Mark Scheme

January 2011

8 (i)	$((a,b)*(c,d))*(e,f) = (ac, ad+b)*(e,f)$	M1	For 3 distinct elements bracketed and attempt to expand
	$= (ace, acf + ad + b)$	A1	For correct expression
	$(a,b)*((c,d)*(e,f)) = (a,b)*(ce, cf + d)$ $= (ace, acf + ad + b)$	A1	3 For correct expression again
(ii)	$(a,b)*(1,1) = (a, a+b), (1,1)*(a,b) = (a, b+1)$	M1	For combining both ways round
	$a+b = b+1 \Rightarrow a = 1$	M1	For equating components (allow from incorrect pairs)
	$\Rightarrow (1, b) \forall b$	A1	3 For correct elements AEF
(iii)	$(mp, mq+n) \text{ OR } (pm, pn+q) = (1, 0)$	M1	For either element on LHS
	$\Rightarrow (p, q) = \left(\frac{1}{m}, -\frac{n}{m}\right)$	A1	2 For correct inverse
(iv)	$(a,b)*(a,b) = (a^2, ab+b) = (1, 0)$	M1	For attempt to find self-inverses
	$\text{OR } (a,b) = \left(\frac{1}{a}, -\frac{b}{a}\right) \Rightarrow a^2 = 1, ab = -b$	B1	For (1, 0). For (-1, b) AEF
	\Rightarrow self-inverse elements (1, 0) and $(-1, b) \forall b$	A1	
(v)	$(0, y)$ has no inverse for any $y \Rightarrow$ not a group	B1	1 For stating any one element with no inverse. Allow $x \neq 0$ required, provided reference to inverse is made "Some elements have no inverse" B0

12

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

14 – 19 Qualifications (General)

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553

