

| | |
|---|------------------|
| Write your name here | |
| Surname | Other names |
| Pearson Edexcel | Centre Number |
| International | Candidate Number |
| Advanced Level | |
| <h1 style="margin: 0;">Chemistry</h1> <h2 style="margin: 0;">Advanced Subsidiary</h2> <h3 style="margin: 0;">Unit 3: Chemistry Laboratory Skills I</h3> | |
| Wednesday 7 May 2014 – Morning | Paper Reference |
| Time: 1 hour 15 minutes | WCH03/01 |
| Candidates may use a calculator. | Total Marks |

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
– *there may be more space than you need.*

Information

- The total mark for this paper is 50.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*
- You will be assessed on your ability to organise and present information, ideas, descriptions and arguments clearly and logically, including your use of grammar, punctuation and spelling.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Keep an eye on the time.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P42980A

©2014 Pearson Education Ltd.

6/6/6/6/



PEARSON

Answer ALL the questions. Write your answers in the spaces provided.

1 A series of tests was carried out on **A**, a white powder. **A** is known to contain one cation and one anion. Complete the table below. You may use names or formulae in your answers.

| | Test | Observation | Inference | |
|-----|---|--|---|-----|
| (a) | Carry out a flame test on A . | | Cation is calcium. | (1) |
| (b) | Add a few drops of dilute nitric acid to an aqueous solution of A , followed by aqueous silver nitrate. Then add concentrated aqueous ammonia solution. | | Anion is probably iodide. This confirms the anion is iodide. | (2) |
| (c) | Add an aqueous solution of chlorine to an aqueous solution of A . | The colour of the resulting solution is | The colour is due to the formation of | (2) |
| (d) | Add an aqueous solution of starch to the mixture formed in (c). | The colour of the resulting mixture is | This confirms the inference made in (c). | (1) |
| (e) | Add a solution of sodium carbonate to an aqueous solution of A . When there is no further change, add dilute hydrochloric acid to the mixture. | A white precipitate forms. The precipitate dissolves in the acid and bubbles of gas are seen. | The precipitate is The gas is | (2) |



(f) When **concentrated** sulfuric acid is added to a **solid** sample of **A**, there is a vigorous redox reaction.

(i) Identify, by name or formula, the product formed by the oxidation of the iodide ion in this reaction. Describe the appearance of this product.

(2)

Product

Appearance

(ii) Identify, by name or formula, one product formed when the concentrated sulfuric acid is reduced. Describe an observation you could make that shows this product has formed.

(2)

Product

Observation

.....

(Total for Question 1 = 12 marks)



2 This question concerns the analysis of an organic compound.

- (a) (i) How can the relative molecular mass of a compound be found from its mass spectrum?

(1)

- (ii) The general formula of an alcohol can be written ROH, where R is an alkyl group.

The relative molecular mass of an alcohol **Q** is 88. The formula of the alkyl group may be represented as C_xH_y.

State the values of x and y.

(1)

x y

- (b) When **Q** was warmed with a mixture of sulfuric acid and aqueous potassium dichromate(VI), there was no colour change.

Deduce the displayed formula of alcohol **Q**.

(1)



(c) When a sample of **Q** was reacted with phosphorus(V) chloride, PCl_5 , steamy fumes were seen.

(i) Identify these steamy fumes by name or formula.

(1)

(ii) The steamy fumes were tested by reacting them with ammonia gas. A white smoke was seen.

Write an equation, including state symbols, for the reaction in which the white smoke was formed.

(2)

(d) One of the isomers of the alcohol **Q** is an ether. Ethers contain two alkyl groups linked by an oxygen atom and can be represented as R-O-R.

Explain how the information in an **infrared** spectrum would be used to decide whether the spectrum is produced by an alcohol or an ether. Wavenumber data are not required.

(1)

(Total for Question 2 = 7 marks)



- 3 (a) The concentrations of acids and alkalis can be found by titration using a suitable indicator.

Give the colours which are seen if the indicator phenolphthalein is used.

(2)

Colour in acid

Colour in alkali

- (b) Another type of titration is a **thermometric** titration.


In a thermometric titration, 20.0 cm^3 of 1.50 mol dm^{-3} hydrochloric acid was placed in a well-insulated cup, and its temperature was measured. Portions of sodium hydroxide solution were added from a burette. The mixture was stirred continuously and the temperature measured after each addition.

| | | | | | | | |
|--|------|------|-------|-------|-------|-------|-------|
| Total volume of sodium hydroxide added / cm^3 | 0.00 | 5.00 | 10.00 | 15.00 | 20.00 | 25.00 | 30.00 |
| Temperature / $^{\circ}\text{C}$ | 20.4 | 22.8 | 25.5 | 28.0 | 27.2 | 24.1 | 20.8 |

On the axes opposite, plot a graph of temperature against the total volume of sodium hydroxide added. Draw two straight lines on your graph and extrapolate the lines until they intersect. Hence find the maximum temperature of the reaction mixture and the total volume of sodium hydroxide which just neutralized the hydrochloric acid.

(4)





Maximum temperature.....

Total volume of sodium hydroxide that just neutralized the hydrochloric acid.

.....



(c) In an experiment using a **different** sample of sodium hydroxide solution, 20.0 cm³ of 1.50 mol dm⁻³ hydrochloric acid was neutralized by 15.50 cm³ of sodium hydroxide solution. The starting temperature was 20.4°C and the temperature at neutralization was 30.6°C.

(i) Calculate the energy, in joules, transferred when the acid is just neutralized.

$$\begin{array}{ccccccc} \text{Energy transferred} & = & \text{total mass of solution} & \times & 4.18 & \times & \text{temperature rise} \\ \text{(J)} & & \text{(g)} & & (\text{J g}^{-1} \text{ } ^\circ\text{C}^{-1}) & & \text{(} ^\circ\text{C)} \end{array}$$

Assume that the density of the solution is 1 g cm⁻³.

(1)

(ii) The number of moles of hydrochloric acid used was 3.00×10^{-2} .

Calculate the enthalpy change of the reaction, in kJ mol⁻¹, for the neutralization of one mole of hydrochloric acid.

Give your answer to **three** significant figures and include a sign.

(2)

$\Delta H = \dots\dots\dots$ kJ mol⁻¹



(iii) Why is it important that the temperature readings are taken as quickly as possible?

(1)

.....

.....

(iv) Thermometric titrations can also be carried out using an electronic probe connected to a computer, instead of a thermometer.

The sodium hydroxide is run into the acid from the burette at a steady rate. The acid is in an insulated beaker with a magnetic stirrer. The computer then produces a plot of the results.

Explain why this modified method can give improved results, other than because of any increase in accuracy of the temperature readings by the electronic probe.

(2)

.....

.....

.....

.....

(d) (i) Calculate the concentration, in mol dm⁻³, of the sodium hydroxide used when 20.0 cm³ of 1.50 mol dm⁻³ hydrochloric acid is neutralized by 15.50 cm³ of sodium hydroxide.

(2)



(ii) Each time a burette is read, the error is $\pm 0.05 \text{ cm}^3$.

Calculate the percentage error in using a burette to measure a volume of 5.00 cm^3 of sodium hydroxide.

(1)

(e) (i) When a titration is carried out using an indicator, the concentrations of acid and alkali are usually between 0.05 and 0.20 mol dm^{-3} .

Explain why more concentrated solutions are used in thermometric titrations.

(1)

.....

.....

.....

.....

(ii) Sodium hydroxide is described as an irritant at concentrations less than 0.50 mol dm^{-3} .

In what way is more concentrated sodium hydroxide hazardous?

(1)

.....

.....

(Total for Question 3 = 17 marks)



BLANK PAGE



- 4 Butanone, $\text{CH}_3\text{COCH}_2\text{CH}_3$, can be prepared from butan-2-ol, $\text{CH}_3\text{CH}(\text{OH})\text{CH}_2\text{CH}_3$, using the procedure below.

An organic solvent suitable for this procedure has a low boiling temperature and is extremely flammable, so adequate safety precautions must be taken.

Procedure

1. Place about 10 g of sodium dichromate(VI) and 20 cm^3 of distilled water in a conical flask. Shake the flask to dissolve the solid. Then slowly add about 8 cm^3 of concentrated sulfuric acid.
 2. Dissolve 5.00 g of butan-2-ol in the organic solvent in a round-bottom flask. Stand the flask in a large beaker containing ice and water. Slowly add the acidified sodium dichromate(VI) solution through a funnel to the butan-2-ol solution in the flask.
 3. When the addition is finished, leave the mixture to cool and separate the organic layer, which contains the butanone, from the aqueous layer.
 4. Wash the organic layer with sodium hydrogencarbonate solution, and then with water. Discard the aqueous layer.
 5. Add some sodium sulfate, Na_2SO_4 , to the organic layer and wait until this solution is clear.
 6. Decant the solution into a flask, and add a few anti-bumping granules. Use distillation to remove the solvent, which has a **lower** boiling temperature than butanone. The solvent boils between 32°C and 36°C .
- (a) What colour change will be seen when the acidified sodium dichromate(VI) reacts with the butan-2-ol?

(1)

From to



(b) The reaction is exothermic. Other than the risk of explosion, why is it important to cool the flask in a beaker of ice and water in **step 2**?

(1)

.....

.....

.....

(c) State the purpose of washing the crude butanone in **step 4** with sodium hydrogencarbonate solution. Describe the method used to carry out this process, naming the piece of apparatus used.

(3)

Purpose

Method

.....

.....

.....

.....

(d) What is the purpose of adding sodium sulfate in **step 5**?

(1)

.....

.....



- (e) Draw a labelled diagram of the apparatus used in **step 6** to distil off the solvent from the organic layer. The diagram should show at least one precaution which must be taken when distilling an extremely flammable liquid.

(4)

- (f) (i) Calculate the volume, in cm^3 , of 5.00 g of butan-2-ol.

The density of butan-2-ol is 0.805 g cm^{-3} .

(1)



- (ii) Each mole of butan-2-ol can produce a maximum yield of one mole of butanone.

Calculate the mass of butan-2-ol that would be required to make 3.00 g of butanone if the yield is 64%.

Relative molecular masses:

| | |
|------------|------|
| butan-2-ol | 74.1 |
| butanone | 72.1 |

(3)

(Total for Question 4 = 14 marks)

TOTAL FOR PAPER = 50 MARKS



The Periodic Table of Elements

| | 1 | 2 | | | | | | | | | | | 3 | 4 | 5 | 6 | 7 | 0 (8) | | | | | | | | | | | | | | | | | | | | | | | | | |
|-------|-----------------------------|---------------------|------------------------------|------|----------------------------|-------------------------------|---------------------------------|-----------------------------------|------------------------------|-------|------------------------------|-------|--------------------------------|-------|-------------------------------|-------------------------------|------------------------------|------------------------------|--------------------------------|-------|--------------------------------|-------|----------------------------------|-------|---------------------------------|---------------------------------|------------------------------|----------------------------|--------------------------------|-------|-----------------------------|-------|------------------------------|------|-------|-----------------------------|-------|------------------------------|-------|--------------------------|----------------------------|-------|----------------------------|
| | | | (13) | (14) | (15) | (16) | (17) | | | | | | | | | | | (18) | | | | | | | | | | | | | | | | | | | | | | | | | |
| 6.9 | Li lithium 3 | 9.0 | Be beryllium 4 | | | | | | | | | | | 10.8 | B boron 5 | 12.0 | C carbon 6 | 14.0 | N nitrogen 7 | 16.0 | O oxygen 8 | 19.0 | F fluorine 9 | 20.2 | Ne neon 10 | | | | | | | | | | | | | | | | | | |
| 23.0 | Na sodium 11 | 24.3 | Mg magnesium 12 | | | | | | | | | | | 27.0 | Al aluminium 13 | 28.1 | Si silicon 14 | 31.0 | P phosphorus 15 | 32.1 | S sulfur 16 | 35.5 | Cl chlorine 17 | 39.9 | Ar argon 18 | | | | | | | | | | | | | | | | | | |
| 39.1 | K potassium 19 | 40.1 | Ca calcium 20 | (3) | 45.0 | Sc scandium 21 | 47.9 | Ti titanium 22 | (4) | 50.9 | V vanadium 23 | 52.0 | Cr chromium 24 | (5) | 54.9 | Mn manganese 25 | 55.8 | Fe iron 26 | (6) | 58.9 | Co cobalt 27 | 58.7 | Ni nickel 28 | (7) | 59.9 | Cu copper 29 | 63.5 | Zn zinc 30 | (8) | 65.4 | Ga gallium 31 | 69.7 | Ge germanium 32 | (9) | 72.6 | As arsenic 33 | 74.9 | Se selenium 34 | (10) | 79.0 | Br bromine 35 | 83.8 | Kr krypton 36 |
| 85.5 | Rb rubidium 37 | 87.6 | Sr strontium 38 | (4) | 88.9 | Y yttrium 39 | 91.2 | Zr zirconium 40 | (5) | 92.9 | Nb niobium 41 | 95.9 | Mo molybdenum 42 | (6) | [98] | Tc technetium 43 | 101.1 | Ru ruthenium 44 | (7) | 102.9 | Rh rhodium 45 | 106.4 | Pd palladium 46 | (8) | 107.9 | Ag silver 47 | 112.4 | Cd cadmium 48 | (9) | 114.8 | In indium 49 | 118.7 | Sn tin 50 | (10) | 121.8 | Sb antimony 51 | 126.9 | Te tellurium 52 | (11) | 127.6 | I iodine 53 | 131.3 | Xe xenon 54 |
| 132.9 | Cs caesium 55 | 137.3 | Ba barium 56 | (3) | 138.9 | La* lanthanum 57 | 178.5 | Hf hafnium 72 | (4) | 180.9 | Ta tantalum 73 | 183.8 | W tungsten 74 | (5) | 186.2 | Re rhenium 75 | 190.2 | Os osmium 76 | (6) | 192.2 | Ir iridium 77 | 195.1 | Pt platinum 78 | (7) | 197.0 | Au gold 79 | 200.6 | Hg mercury 80 | (8) | 204.4 | Tl thallium 81 | 207.2 | Pb lead 82 | (9) | 209.0 | Bi bismuth 83 | [210] | At astatine 85 | [222] | Rn radon 86 | | | |
| [223] | Fr francium 87 | [226] | Ra radium 88 | (3) | [227] | Ac* actinium 89 | [261] | Rf rutherfordium 104 | (4) | [262] | Db dubnium 105 | [266] | Sg seaborgium 106 | (5) | [264] | Bh bohrium 107 | [277] | Hs hassium 108 | (6) | [268] | Mt meitnerium 109 | [271] | Ds darmstadtium 110 | (7) | [272] | Rg roentgenium 111 | | | | | | | | | | | | | | | | | |
| | | * Lanthanide series | | 140 | Ce cerium 58 | 141 | Pr praseodymium 59 | 144 | Nd neodymium 60 | 150 | Sm samarium 62 | 152 | Eu europium 63 | 157 | Gd gadolinium 64 | 159 | Tb terbium 65 | 163 | Dy dysprosium 66 | 165 | Ho holmium 67 | 167 | Er erbium 68 | 169 | Tm thulium 69 | 173 | Yb ytterbium 70 | 175 | Lu lutetium 71 | | | | | | | | | | | | | | |
| | | * Actinide series | | 232 | Th thorium 90 | [231] | Pa protactinium 91 | 238 | U uranium 92 | [242] | Pu plutonium 94 | [243] | Am americium 95 | [247] | Cm curium 96 | [245] | Bk berkelium 97 | [251] | Cf californium 98 | [254] | Es einsteinium 99 | [253] | Fm fermium 100 | [256] | Md mendelevium 101 | [254] | No nobelium 102 | [257] | Lr lawrencium 103 | | | | | | | | | | | | | | |

Elements with atomic numbers 112-116 have been reported but not fully authenticated

Key

relative atomic mass
atomic symbol
name
atomic (proton) number

1.0
H
hydrogen
1

