Paper Reference(s)

6686/01 Edexcel GCE

Statistics S4

Advanced Subsidiary

Thursday 16 June 2005 – Afternoon

Time: 1 hour 30 minutes

Materials required for examination

Items included with question papers

Answer Book (AB16) Graph Paper (ASG2) Mathematical Formulae (Lilac)

Candidates may use any calculator EXCEPT those with the facility for symbolic algebra, differentiation and/or integration. Thus candidates may NOT use calculators such as the Texas Instruments TI 89, TI 92, Casio CFX 9970G, Hewlett Packard HP 48G.

Instructions to Candidates

In the boxes on the answer book, write the name of the examining body (Edexcel), your centre number, candidate number, the unit title (Statistics S4), the paper reference (6686), your surname, other name and signature.

Values from the statistical tables should be quoted in full. When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

This paper has seven questions.

The total mark for this paper is 75.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled.

You must show sufficient working to make your methods clear to the Examiner. Answers without working may gain no credit.

T	he	rand	lom va	ariab	$ext{le } X ext{ ha}$	as a	χ²-di	stribu	ition wi	th 9	degree	es of	f freedo	om.				
(0	a)	Finc	d P(2.0)88 <	< <i>X</i> < 1	9.02	23).											(:
T	`he	ranc	lom va	ariab	ole <i>Y</i> fo	ollow	/s an	<i>F</i> -dis	stributio	n w	ith 12	and	5 degre	ees	of freed	lom	1.	(-
(ł	b)) Find the upper and lower 5% critical values for <i>Y</i> .																
																(".	Total	6 mark
y	ard	wa	s 8 m	m. A		nd t	imbe				-			-	-		-	a timbe
																		0.0
(0					_										e that the		_	arly.
		post	s prod	luce	d by th	ese t	imbe	r yar	ds diffe	r in	variab	ility	. State	you		hes	_	early.
		post	s prod	luce	d by th	ese t	imbe	r yar	ds diffe	r in	variab	ility	. State	you	r hypot	hes	ses cle	arly.
(1	<i>b</i>)	State	es prod	ssum	d by th	you	imbe	r yar	ds diffe	r in er to	variab	out	. State	you t in	r hypot	hes	ses cle	early.
(<i>l</i>	h)	State	e an as	ssum	o fill ba	you ags wine i	have vith f	made	ds diffe e in ord	r in er to	carry e mean	out we	. State the test	you t in 101	part (a) 0 grams	(T	Fotal	early.
(<i>l</i>	h)	State	e an as	set to the r, in	o fill ba	ese t you ags v ine i	have vith f	made	ds diffee in ord	er to	carry e mean	out we	. State the test	you t in 101	part (a) 0 grams	(T	Fotal	arly. (£ 6 mark
(ll A T w	A m To coveig	State achiece by our ring	ne is sk that of flour ta su the bag	set to the r, in	o fill bamachigrams,	ags viine ii, in e	with fas wo each to 100 me 5% 1010	lour sorking bag is 6 sign gram	such that g prope as foll	er to	e mean a rand	out we lom	ight is sampl	you t in 101 e or	part (a) 0 grams f 8 bag 1012 or not th	(T. s. is	Fotal s sele 1007	arly. (£ 6 mark

- **4.** A farmer set up a trial to assess the effect of two different diets on the increase in the weight of his lambs. He randomly selected 20 lambs. Ten of the lambs were given diet *A* and the other 10 lambs were given diet *B*. The gain in weight, in kg, of each lamb over the period of the trial was recorded.
 - (a) State why a paired t-test is not suitable for use with these data.

(1)

(b) Suggest an alternative method for selecting the sample which would make the use of a paired t-test valid.

(1)

(c) Suggest two other factors that the farmer might consider when selecting the sample.

(2)

The following paired data were collected.

Diet A	5	6	7	4.6	6.1	5.7	6.2	7.4	5	3
Diet B	7	7.2	8	6.4	5.1	7.9	8.2	6.2	6.1	5.8

(d) Using a paired t-test, at the 5% significance level, test whether or not there is evidence of a difference in the weight gained by the lambs using diet A compared with those using diet B.

(8)

(e) State, giving a reason, which diet you would recommend the farmer to use for his lambs.

(1)

(Total 13 marks)

5. Define

(a) a Type I error, (1)

(b) the size of a test. (1)

Jane claims that she can read Alan's mind. To test this claim Alan randomly chooses a card with one of 4 symbols on it. He then concentrates on the symbol. Jane then attempts to read Alan's mind by stating what symbol she thinks is on the card. The experiment is carried out 8 times and the number of times *X* that Jane is correct is recorded.

The probability of Jane stating the correct symbol is denoted by p.

To test the hypothesis H_0 : p = 0.25 against H_1 : p > 0.25, a critical region of X > 6 is used.

(c) Find the size of this test. (3)

(d) Show that the power function of this test is $8p^7 - 7p^8$. (3)

Given that p = 0.3, calculate

(e) the power of this test, (1)

(f) the probability of a Type II error. (2)

(g) Suggest two ways in which you might reduce the probability of a Type II error. (2)

(Total 12 marks)

- 6. Brickland and Goodbrick are two manufacturers of bricks. The lengths of the bricks produced by each manufacturer can be assumed to be normally distributed. A random sample of 20 bricks is taken from Brickland and the length, x mm, of each brick is recorded. The mean of this sample is 207.1 mm and the variance is 3.2 mm^2 .
 - (a) Calculate the 98% confidence interval for the mean length of brick from Brickland.

(4)

A random sample of 10 bricks is selected from those manufactured by Goodbrick. The length of each brick, *y* mm, is recorded. The results are summarised as follows.

$$\sum y = 2046.2 \qquad \sum y^2 = 418785.4$$

The variances of the length of brick for each manufacturer are assumed to be the same.

(b) Find a 90% confidence interval for the value by which the mean length of brick made by Brickland exceeds the mean length of brick made by Goodbrick.

(8)

(Total 12 marks)

7. A bag contains marbles of which an unknown proportion p is red. A random sample of n marbles is drawn, with replacement, from the bag. The number X of red marbles drawn is noted.

A second random sample of m marbles is drawn, with replacement. The number Y of red marbles drawn is noted.

Given that $p_1 = \frac{aX}{n} + \frac{bY}{m}$ is an unbiased estimator of p,

(a) show that a + b = 1. (4)

Given that $p_2 = \frac{(X+Y)}{n+m}$,

(b) show that p_2 is an unbiased estimator for p. (3)

(c) Show that the variance of p_1 is $p(1-p)\left(\frac{a^2}{n} + \frac{b^2}{m}\right)$.

(d) Find the variance of p_2 . (3)

(e) Given that a = 0.4, m = 10 and n = 20, explain which estimator p_1 or p_2 you should use. (4)

(Total 17 marks)

(3)

TOTAL FOR PAPER: 75 MARKS

END