


Mark Scheme (Results) Summer 2009


GCE

GCE Mathematics (6678/01)

June 2009
6678 Mechanics M2
Mark Scheme

Question Number	Scheme	Marks
Q1	$\mathbf{I} = m\mathbf{v} - m\mathbf{u}$ $5\mathbf{i} - 3\mathbf{j} = \frac{1}{4} \mathbf{v} - \frac{1}{4} (3\mathbf{i} + 7\mathbf{j})$ $\mathbf{v} = 23\mathbf{i} - 5\mathbf{j}$ $ \mathbf{v} = \sqrt{23^2 + 5^2} = 23.5$	<p>M1A1</p> <p>A1</p> <p>M1A1</p> <p style="text-align: right;">[5]</p>
Q2	<p>(a)</p> $\frac{dv}{dt} = 8 - 2t$ $8 - 2t = 0$ $\text{Max } v = 8 \times 4 - 4^2 = 16 \text{ (ms}^{-1}\text{)}$ <p>(b)</p> $\int 8t - t^2 dt = 4t^2 - \frac{1}{3}t^3 (+c)$ <p>($t=0$, displacement = 0 $\Rightarrow c=0$)</p> $4T^2 - \frac{1}{3}T^3 = 0$ $T^2(4 - \frac{T}{3}) = 0 \Rightarrow T = 0, 12$ $T = 12 \text{ (seconds)}$	<p>M1</p> <p>M1</p> <p>M1A1</p> <p style="text-align: right;">(4)</p> <p>M1A1</p> <p>DM1</p> <p>DM1</p> <p>A1</p> <p style="text-align: right;">(5)</p> <p style="text-align: right;">[9]</p>
Q3	<p>(a) Constant $v \Rightarrow$ driving force = resistance $\Rightarrow F=120 \text{ (N)}$ $\Rightarrow P=120 \times 10 = 1200\text{W}$</p> <p>(b) Resolving parallel to the slope, zero acceleration: $\frac{P}{v} = 120 + 300g \sin \theta (= 330)$ $\Rightarrow v = \frac{1200}{330} = 3.6 \text{ (ms}^{-1}\text{)}$</p>	<p>M1</p> <p>M1</p> <p style="text-align: right;">(2)</p> <p>M1A1A1</p> <p>A1</p> <p style="text-align: right;">(4)</p> <p style="text-align: right;">[6]</p>

Question Number	Scheme	Marks
<p>Q4 (a)</p>  <p>(b)</p>	<p>Taking moments about A:</p> $3g \times 0.75 = \frac{T}{\sqrt{2}} \times 0.5$ $T = 3\sqrt{2}g \times \frac{7.5}{5} = \frac{9\sqrt{2}g}{2} (= 62.4N)$ <p> $\leftarrow \pm H = \frac{T}{\sqrt{2}} (= \frac{9g}{2} \approx 44.1N)$ </p> <p> $\uparrow \pm V + \frac{T}{\sqrt{2}} = 3g \quad (\Rightarrow V = 3g - \frac{9g}{2} = \frac{-3g}{2} \approx -14.7N)$ </p> <p> $\Rightarrow R = \sqrt{81+9} \times \frac{g}{2} \approx 46.5(N)$ </p> <p> at angle $\tan^{-1} \frac{1}{3} = 18.4^\circ$ (0.322 radians) below the line of BA 161.6° (2.82 radians) below the line of AB $(108.4^\circ$ or 1.89 radians to upward vertical) </p>	<p>M1A1A1</p> <p>A1</p> <p>(4)</p> <p>B1</p> <p>M1A1</p> <p>M1A1</p> <p>M1A1</p> <p>(7) [11]</p>
<p>Q5 (a)</p> <p>(b)</p>	<p>Ratio of areas triangle:sign:rectangle = 1 : 5 : 6 (1800:9000:10800) Centre of mass of the triangle is 20cm down from AD (seen or implied)</p> <p> $\Rightarrow 6 \times 45 - 1 \times 20 = 5 \times \bar{y}$ $\bar{y} = 50cm$ </p> <p>Distance of centre of mass from AB is 60cm</p> <p>Required angle is $\tan^{-1} \frac{60}{50}$ $= 50.2^\circ$ (0.876 rads)</p> <p>(their values)</p>	<p>B1</p> <p>B1</p> <p>M1A1</p> <p>A1</p> <p>(5)</p> <p>B1</p> <p>M1A1ft</p> <p>A1</p> <p>(4) [9]</p>

Question Number	Scheme	Marks
Q7 (a)	 <p> $KE \text{ at } X = \frac{1}{2}mv^2 = \frac{1}{2} \times 2 \times 14^2$ $GPE \text{ at } Y = mgd \sin \alpha \left(= 2 \times g \times d \times \frac{7}{25} \right)$ $\text{Normal reaction } R = mg \cos \alpha$ $\text{Friction} = \mu \times R = \frac{1}{8} \times 2g \times \frac{24}{25}$ </p> <p> $\text{Work Energy: } \frac{1}{2}mv^2 - mgd \sin \alpha = \mu \times R \times d \text{ or equivalent}$ $196 = \frac{14gd}{25} + \frac{6gd}{25} = \frac{20gd}{25}$ $d = 25 \text{ m}$ </p> <p>(b) Work Energy</p> <p> $\text{First time at } X: \frac{1}{2}mv^2 = \frac{1}{2}m14^2$ $\text{Work done} = \mu \times R \times 2d = \frac{1}{8} \times 2g \times \frac{24}{25} \times 2d$ $\text{Return to } X: \frac{1}{2}mv^2 = \frac{1}{2}m14^2 - \frac{1}{8} \times 2g \times \frac{24}{25} \times 50$ $v = 8.9 \text{ ms}^{-1} \quad (\text{accept } 8.85 \text{ ms}^{-1})$ </p> <p> $\text{OR: Resolve parallel to } XY \text{ to find the acceleration and use of } v^2 = u^2 + 2as$ $2a = 2g \sin \alpha - F_{\max} = 2g \times \frac{7}{25} - \frac{6g}{25} = \frac{8g}{25}$ $v^2 = (0+)^2 + 2 \times a \times s = 8g ; v = 8.9 \quad (\text{accept } 8.85 \text{ ms}^{-1})$ </p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>M1</p> <p>M1A1</p> <p>A1</p> <p>(7)</p> <p>M1A1</p> <p>DM1A1</p> <p>(4)</p> <p>M1A1</p> <p>DM1;A1</p> <p>[11]</p>

Question Number	Scheme	Marks
<p>Q8 (a)</p> <div style="text-align: center; margin: 20px 0;"> </div> <p>Conservation of momentum: $4mu - 3mv = 3mkv$</p> <p>Impact law: $kv = \frac{3}{4}(u + v)$</p> <p>Eliminate k: $4mu - 3mv = 3m \times \frac{3}{4}(u + v)$</p> <p style="text-align: center;">$u = 3v$ (Answer given)</p> <p>(b) $kv = \frac{3}{4}(3v + v), k = 3$</p> <p>(c) Impact law: $(kv + 2v)e = v_C - v_B$ ($5ve = v_C - v_B$)</p> <p>Conservation of momentum: $3 \times kv - 1 \times 2v = 3v_B + v_C$ ($7v = 3v_B + v_C$)</p> <p>Eliminate v_C: $v_B = \frac{v}{4}(7 - 5e) > 0$ hence no further collision with A.</p>		<p>M1A1</p> <p>M1A1</p> <p>DM1</p> <p>A1</p> <p style="text-align: right;">(6)</p> <p>M1,A1</p> <p style="text-align: right;">(2)</p> <p>B1</p> <p>B1</p> <p>M1 A1</p> <p style="text-align: right;">(4)</p> <p style="text-align: right;">[12]</p>