更多咨询请登录 Version 1





General Certificate of Education (A-level) January 2011

Chemistry

CHEM5

(Specification 2420)

Unit 5: Energetics, Redox and Inorganic Chemistry

Final



Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all examiners participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the candidates" responses to questions and that every examiner understands and applies it in the same correct way. As preparation for standardisation each examiner analyses a number of candidates" scripts: alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, examiners encounter unusual answers which have not been raised they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates" reactions to a particular paper. Assumptions about future mark schemes on the basis of one year"s document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from: aqa.org.uk

Copyright © 2010 AQA and its licensors. All rights reserved.

Copyright

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

The Assessment and Qualifications Alliance (AQA) is a company limited by guarantee registered in England and Wales (company number 3644723) and a registered charity (registered charity number 1073334). Registered address: AQA, Devas Street, Manchester M15 6EX.

Question	Marking Guidance	Mark	Comments
1(a)	Enthalpy change for the formation of <u>1 mol</u> of <u>gaseous atoms</u> From the <u>element</u> (in its standard state) Enthalpy change to separate <u>1 mol</u> of an <u>ionic</u> lattice/solid/compound Into (its component) <u>gaseous ions</u>	1 1 1 1	allow <u>heat energy change</u> for <u>enthalpy change</u> ignore reference to conditions enthalpy change not required but penalise energy mark all points independently
1(b)	$\Delta H_{L} = \Delta H_{f} + \Delta H_{a} + 1.E. + 1/2E(Cl-Cl) + EA$ = +411 + 109 + 494 + 121 - 364 = +771 (kJ mol ⁻¹)	1 1 1	Or correct Born-Haber cycle drawn out -771 scores 2/3 +892 scores 1/3 -51 scores 1/3 -892 scores zero +51 scores zero ignore units
1(c)(i)	lons are perfect spheres (or point charges) <u>Only</u> electrostatic attraction/no covalent interaction	1	mention of molecules/intermolecular forces/covalent bonds CE = 0 allow ionic bonding <u>only</u> If mention of atoms CE = 0 for M2
1(c)(ii)	Ionic	1	Allow no covalent character/bonding

1(c)(iii)	Ionic with additional covalent bonding	1	Or has covalent character/partially covalent
			Allow mention of polarisation of ions or description of polarisation

Question	Marking Guidance	Mark	Comments
2(a)	Because it is a <u>gas</u> compared with <u>solid</u> carbon	1	Mark independently
	Nitrogen is more disordered/random/chaotic/free to move	1	
2(b)	0 K / –273 C / absolute zero	1	
2(c)	$\Delta G = \Delta H - T \Delta S$	1	Allow $\Delta H = \Delta G - T \Delta S$
			$T\Delta S = \Delta H - \Delta G$
			$\Delta S = (\Delta H - \Delta G)/T$
			Ignore θ in ΔG^{θ}
2(d)	ΔG is less than or equal to zero ($\Delta G \leq 0$)	1	Allow ΔG is less than zero ($\Delta G < 0$)
			Allow ΔG is equal to zero ($\Delta G = 0$)
			Allow ΔG is negative
2(e)	When $\Delta G = 0$ $T = \Delta H / \Delta S$	1	
	$\Delta H = +90.4$	1	Allow $\Delta H = +90$
	$\Delta S = \sum S(\text{products}) - \sum S(\text{reactants})$	1	
	ΔS = 211.1 – 205.3/2 – 192.2/2 = <u>12.35</u>	1	
	<i>T</i> = (90.4 x 1000)/12.35 = 7320 К /7319.8 <u>К</u>	1	Allow 7230 to 7350 <u>K</u> (Note 7.32 K scores 4 marks)
			Units of temperature essential to score the mark

2(g)	$\Delta H = 1.9 \text{ (kJ mol}^{-1}\text{)}$ $\Delta S = 2.4 - 5.7 = -3.3 \text{ (J K}^{-1} \text{ mol}^{-1}\text{)}$ ΔG is always positive	1 1 1	for M1 and M2 allow no units, penalise wrong units This mark can only be scored if ΔH is +ve and ΔS is –ve
2(f)	Activation energy is high	1	Allow chemical explanation of activation energy Allow needs route with lower activation energy Allow catalyst lowers activation energy

Question	Marking Guidance	Mark	Comments
3(a)	Na ₂ O ionic	1	mention of molecules/intermolecular forces/delocalised electrons, CE = 0
	Strong forces between ions/strong ionic bonding	1	Allow lots of energy to break bonds provided M1 scored
	SiO ₂ macromolecular	1	Allow giant molecular/giant covalent.
			If ions mentioned, $CE = 0$
	Strong <u>covalent bonds</u> (between atoms)	1	Allow lots of energy to break <u>covalent</u> bonds
			If breaking intermolecular forces are mentioned, CE = 0 for M4
3(b)	Higher	1	
	Li^{+} (or Li ion) smaller than Na^{+}	1	Must imply Li⁺ ion
			Allow Li [⁺] has higher charge/size ratio not charge/mass
	Attracts O ^{2−} ion more strongly	1	Allow stronger ionic bonding
			Allow additional attraction due to polarisation in Li ₂ O
			M3 can only be scored if M2 gained
3(c)(i)	Molecular Covalent bonds (between P and O)	1	Do not allow simple covalent BUT simple covalent molecule scores M1 and M2 Ignore reference to van der Waals [®] or dipole-dipole

Mark Scheme – General Certificate of Educatio	n (A-level) Ch	nemistry – Unit 5	: Energetics, Redox an	d Inorganic Chemi	stry – January 2011
	· · · · ·	2	U	0	

3(c)(ii)	Weak van der Waals" forces and/or dipole-dipole forces <u>between</u> molecules	1	Allow weak <u>inter-molecular</u> forces – can score "between" molecules in (c)(i) CE = 0 if ionic or macromolecular mentioned in (c)(i) Must state van der Waals" forces are weak OR low energy needed to break van der Waals" forces
3(d)	Allow -1 to +2	1	Allow balanced equations to form HPO_4^{2-} or $H_2PO_4^{-}$
	$P_4O_{10} + 6H_2O \rightarrow 12H^+ + 4PO_4^{3-}$ (or $4H_3PO_4$)	1	ignore state symbols
	Allow 12 to 14	1	Allow $2Na^+ + O^{2-}$ on LHS, 2NaOH on RHS, ignore s.s.
	$Na_2O + H_2O \rightarrow 2Na^+ + 2OH^-$	1	Mark independently
3(e)	$6Na_2O + P_4O_{10} \rightarrow 4Na_3PO_4$	1	Allow neutralisation, mark independently of M1
	Acid-base	1	Do not allow Acid + Base \rightarrow Salt + Water

Question	Marking Guidance	Mark	Comments
4(a)	Incomplete (or partially filled) d orbitals/sub-shells	1	Do not allow d shell
4(b)	Variable oxidation states	1	
4(c)(i)	[H ₃ N–Ag–NH ₃] ⁺	1	Allow [Cl–Ag–Cl] ⁻ or similar Cu(I) ion Allow compounds in (i), (ii) and (iii) (eg Cl-Be-Cl) Allow no charge shown, penalise wrong charge(s)
4(c)(ii)	Cis platin drawn out as square planar	1	Allow NiX ₄ ^{2–} etc
4(c)(iii)	[CuCl ₄] ²⁻ drawn out as tetrahedral ion	1	Or [CoCl ₄] ^{2–} drawn out
4(d)(i)	$SO_2 + 1/2O_2 \rightarrow SO_3$	1	Allow multiples Allow SO ₂ + $1/2O_2$ + $H_2O \rightarrow H_2SO_4$ ignore state symbols
4(d)(ii)	In a different phase/state (from the reactants)	1	
4(d)(iii)	$V_2O_5 + SO_2 \rightarrow V_2O_4 + SO_3$ $V_2O_4 + 1/2O_2 \rightarrow V_2O_5$	1 1	can be in either order allow multiples
4(d)(iv)	Surface area is increased By use of powder or granules or finely divided	1 1	Allow suspending/spreading out onto a mesh or support

4(e)(i)	Forms two or more co-ordinate bonds	1	Allow more than one co-ordinate bond or <u>donates</u> more than 1 electron pair. Do not allow "has more than one electron pair" Allow uses more than one atom to bond (to TM)
4(e)(ii)	Number of product particles > Number of reactant particles Disorder increases or entropy increases (or entropy change is positive)	1	Allow molecules/entities instead of particles Penalise incorrect numbers (should be $2\rightarrow 5$) Allow ΔG must be negative because $\Delta H = 0$ and ΔS is +ve
4(e)(iii)	6 Cyanide strongly bound to Co (by co-ordinate/covalent bond)	1 1	

Question	Marking Guidance	Mark	Comments
5(a)(i)	Co/Cobalt	1	If Co or Cobalt not given CE = 0
	(+) 4	1	Allow 4 and 2 in either order
	(+) 3	I	
5(a)(ii)	Li → Li ⁺ + e ⁻	1	Ignore state symbols Allow e without -ve sign Do not allow equilibrium sign
5(a)(iii)	Platinum is a conductor (Platinum is) unreactive/inert	1	Ignore mention of surface area or catalyst Allow 2 marks if two properties given on one answer line Apply list principle to contradictions/wrong answers Do not allow platinum resists corrosion
5(a)(iv)	Li reacts with water/forms lithium hydroxide	1	Allow water breaks down (or is electrolysed) on re- charge

Mark Scheme – General Certificate of Education	(A-level) (Chemistry – U	nit 5: Energetics,	Redox and Inorganic	Chemistry -	January 2011
	· /	2		0		

5(b)(i)	Pt SO ₃ ²⁻ (aq), SO ₄ ²⁻ (aq) ClO ₃ ⁻ (aq), Cl ⁻ (aq) Pt	2	State symbols and "" not necessary Allow in place of ',' NOT "," inplace of Ignore H ⁺ and H ₂ O Deduct one mark for each mistake (e.g. Pt missed twice counts as two mistakes) Allow reverse order for whole cell Pt Cl^- , $ClO_3^- \parallel SO_4^{2-}$, $SO_3^{2-} \mid Pt$
5(b)(ii)	$ClO_3^- + 3SO_3^{2-} \rightarrow Cl^- + 3SO_4^{2-}$ Oxidising agent ClO_3^- Reducing agent SO_3^{2-}	1 1 1	

Question	Marking Guidance	Mark	Comments
6(a)	Brown ppt/solid	1	Must be stated Allow CO, such ad De not allow CO
	Gas evolved/enervescence		alone CO_2 evolved. Do not allow CO_2
	$2[Fe(H_2O)_6]^{3^+} + 3CO_3^{2^-} \rightarrow 2Fe(H_2O)_3(OH)_3 + 3CO_2 + 3H_2O$	2	Correct iron product (1) allow Fe(OH) ₃ and in equation Balanced equation (1)
6(b)	White ppt/solid	1	
	Colourless <u>Solution</u>	1	Only award M2 if M1 given or initial ppt mentioned
	$[Al(H_2O)_6]^{3+} + 3OH^- \rightarrow Al(H_2O)_3(OH)_3 + 3H_2O$	1	Allow $[Al(H_2O)_6]^{3+} + 3OH^- \rightarrow Al(OH)_3 + 6H_2O$
	$Al(H_2O)_3(OH)_3 + 3OH^- \rightarrow [Al(OH)_6]^{3-} + 3H_2O$	1	Allow formation of $[Al(H_2O)_{6-x}(OH)_x]^{(x-3)-}$ where x=4,5,6
			Allow product without water ligands
			Allow formation of correct product from $[Al(H_2O)_6]^{3+}$
6(c)	Blue ppt/solid	1	
	(Dissolves to give a) deep blue <u>solution</u>	1	Only award M2 if M1 given or initial ppt mentioned
	$[\operatorname{Cu}(\operatorname{H}_2\operatorname{O})_6]^{2+} + 2\operatorname{NH}_3 \to \operatorname{Cu}(\operatorname{H}_2\operatorname{O})_4(\operatorname{OH})_2 + 2\operatorname{NH}_4^+$	1	Allow $[Cu(H_2O)_6]^{2+} + 2NH_3 \rightarrow Cu(OH)_2 + 2NH_4^+ + 4H_2O$
			Allow two equations: $NH_3 + H_2O \rightarrow NH_4^+ + OH^-$
			then $[Cu(H_2O)_6]^{2+} + 2OH^- \rightarrow Cu(OH)_2 + 4H_2O$ etc
	$Cu(H_2O)_4(OH)_2 + 4NH_3 \rightarrow [Cu(H_2O)_2(NH_3)_4]^{2+} + 2OH^- + 2H_2O$	1	Allow $[Cu(H_2O)_6]^{2+} + 4NH_3 \rightarrow [Cu(H_2O)_2(NH_3)_4]^{2+} + 4H_2O$
6(d)	Green/yellow <u>solution</u>	1	
	$[\operatorname{Cu}(\operatorname{H}_2\operatorname{O})_6]^{2^+} + 4\operatorname{Cl}^- \to [\operatorname{Cu}\operatorname{Cl}_4]^{2^-} + 6\operatorname{H}_2\operatorname{O}$	1	

Question	Marking Guidance	Mark	Comments
7(a)(i)	Ammonia	1	If reagent is missing or incorrect cannot score M3
	Starts as a pink (solution)	1	
	Changes to a yellow/straw (solution)	1	Allow pale brown
			Do not allow reference to a precipitate
7(a)(ii)	(dark) brown	1	Do not allow pale/straw/yellow-brown (i.e. these and other shades except for dark brown)
7(b)(i)	Ruby / red-blue / purple / violet / green	1	Do not allow red or blue
			If ppt mentioned contradiction/CE =0
	Green	1	If ppt mentioned contradiction/CE =0
	$[Cr(H_2O)_6]^{3+} + 6OH^- \rightarrow [Cr(OH)_6]^{3-} + 6H_2O$	1	
	Formula of product	1	Can score this mark in (b) (ii)
7(b)(ii)	$H_2O_2 + 2e^- \rightarrow 2OH^-$	1	
	$2[Cr(OH)_6]^{3-} + 3H_2O_2 \rightarrow 2CrO_4^{2-} + 8H_2O + 2OH^-$	2	Allow 1 mark out of 2 for a balanced half-equation such as Cr(III) \rightarrow Cr(VI) + 3e ⁻
			or $Cr^{3+} + 4H_2O \rightarrow CrO_4^{2-} + 8H^+ + 3e^- etc$
			also for 2Cr(III) + $3H_2O_2 \rightarrow 2CrO_4^{2-}$ (unbalanced)
	Yellow	1	Do not allow orange

7(c)	$2MnO_4^- + 6H^+ + 5H_2O_2 \rightarrow 2Mn^{2+} + 8H_2O + 5O_2$	1	if no equation and uses given ratio can score M2, M3, M4 & M5
	Moles $MnO_4^- = (24.35/1000) \times 0.0187 = 4.55 \times 10^{-4}$	1	Note value must be quoted to at least 3 sig. figs.
		·	M2 is for 4.55 x 10 ⁻⁴
	Moles $H_2O_2 = (4.55 \times 10^{-4}) \times \frac{5/2}{2} = 1.138 \times 10^{-3}$	1	M3 is for x 5/2 (or7/3)
			Mark consequential on molar ratio from candidate's equation
	Moles $H_2 \Omega_2$ in 5 cm ³ original	1	M4 is for x 10
	$= (1 \ 138 \ \times \ 10^{-3}) \times 10 = 0.01138$		
	Original $[H_2O_2] = 0.01138 \times (1000/5) = 2.28 \text{ mol dm}^{-3}$ (allow 2.25-2.30)	1	M5 is for consequentially correct answer from (answer to mark 4) x (1000/5)
			Note an answer of between 2.25 and 2.30 is worth 4 marks)
			If candidate uses given ratio 3/7 max 4 marks:
			M1 : Moles of $MnO_4^- = 4.55 \times 10^{-4}$
			M2 : Moles $H_2O_2 = (4.55 \times 10^{-4}) \times 7/3 = 1.0617 \times 10^{-3}$
			M3 : Moles H ₂ O ₂ in 5 cm ³ original
			= $(1.0617 \times 10^{-3}) \times 10 = 0.01062$
			M4 : Original $[H_2O_2] = 0.01062 \text{ x} (1000/5) = 2.12 \text{ mol} dm^{-3}$
			(allow 2.10 to 2.15)