

Centre No.					Paper Reference	Surname	Initial(s)
Candidate No.					6 6 6 7 / 0 1	Signature	

Paper Reference(s)

6667/01

Edexcel GCE

Further Pure Mathematics FP1 Advanced/Advanced Subsidiary

Monday 1 February 2010 – Afternoon

Time: 1 hour 30 minutes

Examiner’s use only

--	--	--

Team Leader’s use only

--	--	--

Materials required for examination

Mathematical Formulae (Pink)

Items included with question papers

Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Question Number	Leave Blank
1	
2	
3	
4	
5	
6	
7	
8	
9	
Total	

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper. Answer ALL the questions. You must write your answer to each question in the space following the question. When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet ‘Mathematical Formulae and Statistical Tables’ is provided. Full marks may be obtained for answers to ALL questions. The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2). There are 9 questions in this question paper. The total mark for this paper is 75. There are 24 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled. You should show sufficient working to make your methods clear to the Examiner. Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Edexcel Limited copyright policy.
©2010 Edexcel Limited.

Printer’s Log. No.
N35143A

W850/R6667/57570 3/5/5/3



Turn over

1. The complex numbers z_1 and z_2 are given by

$$z_1 = 2 + 8i \quad \text{and} \quad z_2 = 1 - i$$

Find, showing your working,

(a) $\frac{z_1}{z_2}$ in the form $a + bi$, where a and b are real, **(3)**

(b) the value of $\left| \frac{z_1}{z_2} \right|$, **(2)**

(c) the value of $\arg \frac{z_1}{z_2}$, giving your answer in radians to 2 decimal places. **(2)**



2.

$$f(x) = 3x^2 - \frac{11}{x^2}$$

- (a) Write down, to 3 decimal places, the value of $f(1.3)$ and the value of $f(1.4)$. (1)

The equation $f(x) = 0$ has a root α between 1.3 and 1.4

- (b) Starting with the interval $[1.3, 1.4]$, use interval bisection to find an interval of width 0.025 which contains α . (3)

- (c) Taking 1.4 as a first approximation to α , apply the Newton-Raphson procedure once to $f(x)$ to obtain a second approximation to α , giving your answer to 3 decimal places. (5)



4.




Figure 1

Figure 1 shows a sketch of part of the parabola with equation $y^2 = 12x$.

The point P on the parabola has x -coordinate $\frac{1}{3}$.

The point S is the focus of the parabola.

(a) Write down the coordinates of S .

(1)

The points A and B lie on the directrix of the parabola.

The point A is on the x -axis and the y -coordinate of B is positive.

Given that $ABPS$ is a trapezium,

(b) calculate the perimeter of $ABPS$.

(5)



5. $\mathbf{A} = \begin{pmatrix} a & -5 \\ 2 & a+4 \end{pmatrix}$, where a is real.

(a) Find $\det \mathbf{A}$ in terms of a .

(2)

(b) Show that the matrix \mathbf{A} is non-singular for all values of a .

(3)

Given that $a = 0$,

(c) find \mathbf{A}^{-1} .

(3)



7. The rectangular hyperbola H has equation $xy = c^2$, where c is a constant.

The point $P\left(ct, \frac{c}{t}\right)$ is a general point on H .

(a) Show that the tangent to H at P has equation

$$t^2 y + x = 2ct$$

(4)

The tangents to H at the points A and B meet at the point $(15c, -c)$.

(b) Find, in terms of c , the coordinates of A and B .

(5)



9.

$$\mathbf{M} = \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix}$$

- (a) Describe fully the geometrical transformation represented by the matrix \mathbf{M} . (2)

The transformation represented by \mathbf{M} maps the point A with coordinates (p, q) onto the point B with coordinates $(3\sqrt{2}, 4\sqrt{2})$.

- (b) Find the value of p and the value of q . (4)

- (c) Find, in its simplest surd form, the length OA , where O is the origin. (2)

- (d) Find \mathbf{M}^2 . (2)

The point B is mapped onto the point C by the transformation represented by \mathbf{M}^2 .

- (e) Find the coordinates of C . (2)

