



GCE

Mathematics

Advanced Subsidiary GCE

Unit **4721**: Core Mathematics 1

Mark Scheme for June 2011

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of pupils of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, OCR Nationals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by Examiners. It does not indicate the details of the discussions which took place at an Examiners' meeting before marking commenced.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the Report on the Examination.


OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2011

Any enquiries about publications should be addressed to:

OCR Publications
PO Box 5050
Annesley
NOTTINGHAM
NG15 0DL

Telephone: 0870 770 6622
Facsimile: 01223 552610
E-mail: publications@ocr.org.uk

<p>1</p> $3(x^2 - 6x) + 4$ $= 3[(x - 3)^2 - 9] + 4$ $= 3(x - 3)^2 - 23$	<p>B1 $p = 3$</p> <p>B1 $(x - 3)^2$ seen or $q = -3$</p> <p>M1 $4 - 3q^2$ or $\frac{4}{3} - q^2$ (their q)</p> <p>A1 $r = -23$</p> <p>4 4</p>	<p>If p, q, r found correctly, then ISW slips in format.</p> <p>$3(x - 3)^2 + 23$ B1 B1 M0 A0</p> <p>$3(x - 3) - 23$ B1 B1 M1 A1 (BOD)</p> <p>$3(x - 3x)^2 - 23$ B1 B0 M1 A0</p> <p>$3(x^2 - 3)^2 - 23$ B1 B0 M1 A0</p> <p>$3(x + 3)^2 - 23$ B1 B0 M1 A1 (BOD)</p> <p>$3x(x - 3)^2 - 23$ B0 B1M1A1</p>
<p>2 (i)</p> 	<p>B1 Reasonably correct curve for $y = \frac{1}{x}$ in 1st and 3rd quadrants only</p> <p>B1 2 Very good curves for $y = \frac{1}{x}$ in 1st and 3rd quadrants</p> <p>SC If 0, very good single curve in either 1st or 3rd quadrant and nothing in other three quadrants. B1</p>	<p>N.B. Ignore 'feathering' now that answers are scanned. Reasonably correct shape, not touching axes more than twice.</p> <p>Correct shape, not touching axes, asymptotes clearly the axes. Allow slight movement away from asymptote at one end but not more. Not finite.</p>
<p>(ii) Translation 4 units parallel to y axis</p>	<p>B1 Must be translation/translated – not shift, move etc.</p> <p>B1 2 4 Or $\begin{pmatrix} 0 \\ 4 \end{pmatrix}$</p>	<p>For "parallel to the y axis" allow "vertically", "up", "in the (positive) y direction". Do not accept "in/on/across/up/along the y axis"</p>
<p>3 (i)</p> $\frac{16x^2 \times 2x^3}{x}$ $= 32x^4$	<p>B1 32</p> <p>B1 2 x^4</p>	
<p>(ii) $\frac{1}{6}x$</p>	<p>M1 6 or $\frac{1}{36^{\frac{1}{2}}}$ or $\frac{1}{\sqrt{36}}$ seen</p> <p>A1 $\frac{1}{6}$ in final answer</p> <p>B1 $\frac{3}{5}x$ (Allow x^1) in final answer</p>	<p>$\frac{1}{\sqrt{36}}$ is M0</p> <p>$\pm \frac{1}{6}$ is A0</p>

4	$2x^2 - 8x + 8 = 26 - 3x$	M1	Attempt to eliminate x or y	Must be a clear attempt to reduce to one variable. Condone poor algebra for first mark. <u>If x eliminated:</u> $y = 2\left(\frac{26 - y}{3} - 2\right)^2$ Leading to $2y^2 - 89y + 800 = 0$ $(2y - 25)(y - 32) = 0$ etc.
	$2x^2 - 5x - 18 (= 0)$	A1	Correct 3 term quadratic (not necessarily all in one side)	
	$(2x - 9)(x + 2) (= 0)$	M1	Correct method to solve quadratic	
	$x = \frac{9}{2}, x = -2$	A1	x values correct	
	$y = \frac{25}{2}, y = 32$	A1	5 y values correct	
		5	SR If A0 A0, one correct pair of values, spotted or from correct factorisation www B1	
5 (i)	$10\sqrt{3} - 4\sqrt{3}$	M1	Attempt to express both surds in terms of $\sqrt{3}$	e.g. $\sqrt{3} \times 100 - \sqrt{3} \times 16$
		B1	One term correct	
	$= 6\sqrt{3}$	A1	3 Fully correct (not $\pm 6\sqrt{3}$)	
(ii)	$\frac{\sqrt{5}(15 + \sqrt{40})}{5}$	M1	Multiply numerator and denominator by $\sqrt{5}$ or $-\sqrt{5}$ or attempt to express both terms of numerator in terms of $\sqrt{5}$ (e.g. dividing both terms by $\sqrt{5}$)	Check both numerator and denominator have been multiplied
	$= \frac{15\sqrt{5} + 10\sqrt{2}}{5}$	B1	One of a, b correctly obtained	
	$= 3\sqrt{5} + 2\sqrt{2}$	A1	3 Both a = 3 and b=2 correctly obtained	
		6		

6	$k = x^{\frac{1}{4}}$ $3k^2 - 8k + 4 = 0$ $(3k - 2)(k - 2) = 0$ $k = \frac{2}{3}$ or $k = 2$ $x = \left(\frac{2}{3}\right)^4$ or $x = 2^4$ $x = \frac{16}{81}$ or $x = 16$	M1* DM1 A1 M1 A1 5 5	Use a substitution to obtain a quadratic or factorise into 2 brackets each containing $x^{\frac{1}{4}}$ Correct method to solve a quadratic Attempt to calculate k^4	No marks unless evidence of substitution (quadratic seen or square rooting or squaring of roots found). = 0 may be implied. Allow $x = x^{\frac{1}{4}}$ as a substitution. No marks if straight to quadratic formula to get $x = \frac{2}{3}$ or $x = 2$ and no further working No marks if $k = x^{\frac{1}{4}}$ then $3k - 8k^2 + 4 = 0$ SC If M0 Spotted solutions www B1 each Justifies 2 solutions exactly B3
	<u>If candidates use $k = x^{\frac{1}{2}}$ and rearrange:</u> $3k - 8\sqrt{k} + 4 = 0$ $8\sqrt{k} = 3k + 4$ $64k = 9k^2 + 24k + 16$ $9k^2 - 40k + 16 = 0$ $(9k - 4)(k - 4) = 0$ $k = \frac{4}{9}$ or $k = 4$ $x = \left(\frac{4}{9}\right)^2$ or $x = 4^2$ $x = \frac{16}{81}$ or $x = 16$	M1* DM1 A1 M1 A1	Substitute, rearrange and square both sides Correct method to solve quadratic Attempt to calculate k^2	
7 (i)	$-14 \leq 6x \leq -5$ $-\frac{7}{3} \leq x \leq -\frac{5}{6}$	M1 A1 A1 3	2 equations or inequalities both dealing with all 3 terms resulting in $a \leq 6x \leq b$, $a \neq -9$, $b \neq 0$ -14 and -5 seen www Accept as two separate inequalities provided not linked by "or" (must be \leq)	Do not ISW after correct answer if contradictory inequality seen. Allow $-\frac{14}{6} \leq x \leq -\frac{5}{6}$
(ii)	$0 < x^2 - 4x - 12$ $(x - 6)(x + 2)$ $x > 6, x < -2$	M1 M1 A1 M1 A1 5 8	Rearrange to collect all terms on one side Correct method to find roots 6, -2 seen Correct method to solve quadratic inequality i.e. $x >$ their higher root, $x <$ their lower root (not wrapped, strict inequalities, no 'and')	Do not ISW after correct answer if contradictory inequality seen. e.g. for last two marks, $-2 > x > 6$ scores M1 A0

8 (i)	$\frac{dy}{dx} = 6x + 6x^{-2}$	M1	Attempt to differentiate (one non-zero term correct)	NB $x = -1$ (and therefore possibly $y = 7$) can be found from equating the incorrect differential	
		A1	Completely correct		
	$6x + \frac{6}{x^2} = 0$	M1	Sets their $\frac{dy}{dx} = 0$	$\frac{dy}{dx} = 6x + 6$ to 0. This could score M1A0 M1A0A1 ft	
	$x = -1$	A1	Correct value for x - www		
$y = 7$	A1 ft	5	Correct value of y for <i>their</i> value of x	If more than one value of x found, allow A1 ft for one correct value of y	
(ii)	$\frac{d^2y}{dx^2} = 6 - 12x^{-3}$	M1	Correct method e.g. substitutes their x from (i) into their $\frac{d^2y}{dx^2}$ (must involve x) and considers sign.	Allow comparing signs of their $\frac{dy}{dx}$ either side of their “- 1”, comparing values of y to their “7”	
	When $x = -1$, $\frac{d^2y}{dx^2} > 0$ so minimum pt	A1 ft	2	ft from their $\frac{dy}{dx}$ differentiated correctly and correct	SC $\frac{d^2y}{dx^2} = a$ constant correctly obtained from their $\frac{dy}{dx}$ and correct conclusion (ft) B1
		7	7	substitution of <i>their</i> value of x and consistent final conclusion NB If second derivate evaluated, it must be correct (18 for $x = -1$). If more than one value of x used, max M1 A0	

<p>9 (i)</p> <p>Gradient of $AB = \frac{1-3}{7-1} = -\frac{1}{3}$</p> <p>Gradient of $AC = \frac{-9-3}{-3-1} = 3$</p> <p>Vertex A OR: Length of $AB = \sqrt{(7-1)^2 + (1-3)^2} = \sqrt{40}$ $AC = \sqrt{(-3-1)^2 + (-9-3)^2} = \sqrt{160}$ $BC = \sqrt{(-3-7)^2 + (-9-1)^2} = \sqrt{200}$ Shows that $AB^2 + AC^2 = BC^2$ Vertex A</p>	M1*	Uses $\frac{y_2 - y_1}{x_2 - x_1}$ for any 2 points	
	A1	One correct gradient (may be for gradient of BC)	
	A1	=1)	
	M1	Gradients for both AB and AC found correctly	Do not allow final mark if vertex A found from wrong working. (Dependent on 1 st M 1 A1 A1)
	DB1	Attempts to show that $m_1 \times m_2 = -1$ oe, accept “negative reciprocal”	Accept $B\hat{A}C$ etc for vertex A or “between AB and AC” Allow if marked on diagram.
	M1*	Correct use of Pythagoras, square rooting not needed	
	A1	Any length or length squared correct	
	A1	All three correct	
	M1	5 Correct use of Pythagoras to show $AB^2 + AC^2 = BC^2$	i.e must add squares of shorter two lengths
	DB1		
<p>9 (ii)</p> <p>Midpoint of BC is $\left(\frac{7+(-3)}{2}, \frac{1+(-9)}{2}\right)$ $= (2, -4)$</p> <p>Length of $BC = \sqrt{(-3-7)^2 + (-9-1)^2} = \sqrt{200} = 10\sqrt{2}$</p> <p>Radius = $5\sqrt{2}$ $(x-2)^2 + (y+4)^2 = (5\sqrt{2})^2$ $(x-2)^2 + (y+4)^2 = 50$ $x^2 + y^2 - 4x + 8y - 30 = 0$</p>	M1*	Uses $\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right)$ o.e. for BC, AB or	<u>Substitution method 1</u> (into $x^2 + y^2 + ax + by + c = 0$) Substitutes all 3 points to get 3 equations in a, b, c M1 At least 2 equations correct A1 Correct method to find one variable M1 One of a, b, c correct A1 Correct method to find other values M1 All values correct A1 Correct equation in required form A1
	A1	AC (3 out of 4 subs correct) Correct centre (cao)	<u>Alternative markscheme for last 4 marks with f, g, c method:</u> $x^2 - 4x + y^2 + 8y$ for their centre DM1* $c = (\pm 2)^2 + 4^2 - 50$ DM1** $c = -30$ A1 Correct equation in required form A1
	M1**	Correct method to find d or r or d^2 or r^2 o.e. for BC, AB or AC (must be consistent with their midpoint if found)	<u>Ends of diameter method (p, q) to (c, d):</u> Attempts to use $(x-p)(x-c) + (y-q)(y-d) = 0$ for BC, AC or AB M2 $(x-7)(x+3) + (y-1)(y+9) = 0$ A2 for both x brackets correct, A2 for both y brackets correct $x^2 + y^2 - 4x + 8y - 30 = 0$ A1 SC If M2 A0 A0 then B1 if both x brackets correct and B1 if both y brackets correct for AC or AB
	DM1*	7 $(x-a)^2 + (y-b)^2$ seen for their centre	
	DM1**	12 $(x-a)^2 + (y-b)^2 = \text{their } r^2$	
	A1	Correct equation	
	A1	Correct equation in required form	


4721

Mark Scheme

June 2011

Substitution method 2 into $(x-p)^2 + (y-q)^2 = \text{their } r^2$
 Correct method to find d or r or d^2 or r^2 *M1
 Substitutes all 3 points to get 3 equations in p, q DM1
 At least 2 equations correct A1
 Correct method to find one variable M1
 One of p, q correct A1
 Correct equation $[(x-2)^2 + (y+4)^2 = 50]$ A1
 Correct equation in required form
 $[x^2 + y^2 - 4x + 8y - 30 = 0]$ A1

10(i)



B1 +ve cubic with 3 distinct roots
B1 (0, 3) labelled or indicated on y-axis
B1 3 (-3, 0), $(\frac{1}{2}, 0)$ and (1, 0) labelled or indicated on x-axis and no other x- intercepts

For first **B1**, left end of curve must finish below x axis and right end must end above x axis. Allow slight wrong curvature at one end but not both ends. No cusp at either turning point. No straight lines drawn with a ruler. Condone (0, 3) as maximum point.
 To gain second and third **B** marks, there must be an attempt at a curve, not just points on axes.
 Final **B1** can be awarded for a negative cubic.

(ii) $2x^2 + 5x - 3, x^2 + 2x - 3, 2x^2 - 3x + 1$ **B1** Obtain one quadratic factor (can be unsimplified)
 $(2x^2 + 5x - 3)(x - 1)$ **M1** Attempt to multiply a quadratic by a linear factor
 $2x^3 + 3x^2 - 8x + 3$ **A1**
 $\frac{dy}{dx} = 6x^2 + 6x - 8$ **M1** Attempt to differentiate (one non-zero term correct)
 When $x = 1$, gradient = 4 **A1** Fully correct expression **www**
A1 6 Confirms gradient = 4 at $x = 1$ **AG

Alternative for first 3 marks:
 Attempt to expand all 3 brackets with an appropriate number of terms (including an x^3 term) **M1**
 Expansion with at most 1 incorrect term **A1**
 Correct, answer (can be unsimplified) **A1**
 Allow if done in part(i) please check.

(iii) Gradient of $l = 4$ **B1** May be embedded in equation of line
 On curve, when $x = -2, y = 15$ **B1** Correct y coordinate
 $y - 15 = 4(x + 2)$ **M1** Correct equation of line using their values
 $y = 4x + 23$ **A1** 4 Correct answer **in correct form**

M mark is for any equation of line with any non-zero numerical gradient through (-2, their evaluated y)

(iv) Attempt to find gradient of curve when $x = -2$ **M1** Substitute $x = -2$ into their $\frac{dy}{dx}$
 $6(-2)^2 + 6(-2) - 8 = 4$ **A1** Obtain gradient of 4 **CWO**
 So line is a tangent **A1** 3 Correct conclusion
16

Alternatives
1) Equates equation of l to equation of curve and attempts to divide resulting cubic by $(x + 2)$ **M1**
 Obtains $(x + 2)^2 (2x - 5) (=0)$ **A1**
 Concludes repeated root implies tangent at $x = -2$ **A1**
2) Equates their gradient function to 4 and uses correct method to solve the resulting quadratic **M1**
 Obtains $(x + 2)(x - 1) = 0$ oe **A1**
 Correctly concludes gradient = 4 when $x = -2$ **A1**

Allocation of method mark for solving a quadratic

e.g. $2x^2 - 5x - 18 = 0$

1) If the candidate attempts to solve by factorisation, their attempt when expanded must produce the **correct quadratic term** and **one other correct term** (with correct sign):

$(2x + 2)(x - 9) = 0$

M1 $2x^2$ and -18 obtained from expansion

$(2x + 3)(x - 4) = 0$

M1 $2x^2$ and $-5x$ obtained from expansion

$(2x - 9)(x - 2) = 0$

M0 only $2x^2$ term correct

2) If the candidate attempts to solve by using the formula

a) If the formula is quoted incorrectly then **M0**.

b) If the formula is quoted correctly then one **sign slip** is permitted. Substituting the wrong numerical value for a or b or c scores **M0**

$$\frac{-5 \pm \sqrt{(-5)^2 - 4 \times 2 \times -18}}{2 \times 2}$$

earns **M1** (minus sign incorrect at start of formula)

$$\frac{5 \pm \sqrt{(-5)^2 - 4 \times 2 \times 18}}{2 \times 2}$$

earns **M1** (18 for c instead of -18)

$$\frac{-5 \pm \sqrt{(-5)^2 - 4 \times 2 \times 18}}{2 \times 2}$$

M0 (2 sign errors: initial sign and c incorrect)

$$\frac{5 \pm \sqrt{(-5)^2 - 4 \times 2 \times -18}}{2 \times -5}$$

M0 (2b on the denominator)

Notes – for equations such as $2x^2 - 5x - 18 = 0$, then $b^2 = 5^2$ would be condoned in the discriminant and would not be counted as a sign error. Repeating the sign error for a in both occurrences in the formula would be two sign errors and score **M0**.

c) If the formula is not quoted at all, substitution must be completely correct to earn the **M1**

3) If the candidate attempts to complete the square, they must get to the “square root stage” involving \pm ; we are looking for evidence that the candidate knows a quadratic has two solutions!

$2x^2 - 5x - 18 = 0$

$2\left(x^2 - \frac{5}{2}x\right) - 18 = 0$

$2\left[\left(x - \frac{5}{4}\right)^2 - \frac{25}{16}\right] - 18 = 0$

$\left(x - \frac{5}{4}\right)^2 = \frac{169}{16}$

$x - \frac{5}{4} = \pm \sqrt{\frac{169}{16}}$

← This is where the **M1** is awarded – arithmetical errors may be condoned provided $x - \frac{5}{4}$ seen or implied

If a candidate makes repeated attempts (e.g. fails to factorise and then tries the formula), mark only what you consider to be their last full attempt.

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

14 – 19 Qualifications (General)

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553

