



GCE MARKING SCHEME


**CHEMISTRY
AS/Advanced**

SUMMER 2011


CHEMISTRY - CH2**SECTION A**

Q.1	(a)	Calcium carbonate	[1]
	(b)	Sodium carbonate	[1]
Q.2		Metallic (1) Covalent and van der Waals (1)	[2]
Q.3		$\text{Ca}_3(\text{PO}_4)_2$	[1]
Q.4		D	[1]
Q.5		Materials that change their properties in response to a change in conditions / environment / surroundings	[1]
Q.6	(a)	Alkene / double bond (1) Alcohol / hydroxyl / hydroxy (1)	[2]
	(b)	$\text{C}_5\text{H}_{10}\text{O}$	[1]
			Total [10]

SECTION B


- Q.7** (a) Compound that contains no double bonds / single bonds only
(Accept contains maximum number of hydrogens) [1]
- (b) (i) $C_3H_8 + 5O_2 \longrightarrow 3CO_2 + 4H_2O$ [2]
products (1)
balancing (1)
- (ii)  [1]
- (c) Cracking (1)
Heat fraction strongly / heat over a catalyst (1)
Accept equation or description of cracking [2]
- (d) Planar molecule with trigonal arrangement about each atom / bond angles roughly 120° (1)
Four (single) **covalent** C – H bonds and one C = C double bond (1)
 π bond in C = C formed by sideways overlap of p orbital (1) [3]
QWC: Information is organised clearly and coherently, using specialist vocabulary where appropriate. [1]
- (e) Electrophilic addition (1)
$$\begin{array}{c} \text{H} & \text{H} \\ | & | \\ \text{Br}-\text{C} & -\text{C}-\text{Br} \\ | & | \\ \text{H} & \text{H} \end{array} \quad \text{accept} \quad \begin{array}{c} \text{H} & \text{H} \\ | & | \\ \text{Br}-\text{C} & -\text{C}-\text{OH} \\ | & | \\ \text{H} & \text{H} \end{array}$$
 [2]
- (f) Phosphoric acid [1]
- (g) Moles ethanol = $\frac{230}{46} = 5$ (1)
Moles glucose = 2.5 (1)
Mass glucose = $2.5 \times 180 = 450$ g (1) [3]

Total [16]





UV light (1)

any of following for 4 max




Free radical substitution / photochlorination (1)



[6]

QWC: Selection of form and style of writing appropriate to purpose and to complexity of subject matter. [1]



Nucleophilic substitution / hydrolysis

[2]

(c) Heat with NaOH (1)

Add HNO_3 then AgNO_3 (1)

White precipitate seen (1)


[3]

(d) Ozone layer depleted / (leads to) increased incidence of skin cancer


Contributes to greenhouse effect / increases global warming

[1]

Total [13]

- Q.9** (a) C=O absorption at 1650–1750 cm^{-1}
 C–O absorption at 1000–1300 cm^{-1}
 O–H absorption at 2500–3500 cm^{-1}
 3 correct peaks labelled [2]
 (2 correct peaks labelled 1 mark)
- (b) Molecular ion at m/z 60 shows that M_r is 60 (1)
 Peak at m/z 15 shows CH_3 group / peak at m/z 45 shows COOH group (1) [2]
- (c) (i)  [1]
 (Accept 1 hydrogen bond)
- (ii) (Intermolecular bond formed) when hydrogen attached to a highly electronegative atom (oxygen) (1)
 is bonded to an electronegative atom in another molecule (1)
 forming very strong dipole – dipole attraction (1) [3]
QWC: Legibility of text; accuracy of spelling, punctuation and grammar, clarity of meaning [1]
- (d) (i) Acidified and heat / reflux [1]
 (ii) Colour change from orange to green [1]
- (e) Propane would be lower as it cannot form hydrogen bonds / only forms van der Waals forces between molecules (1)
 Butan-1-ol would be higher as it (also has hydrogen bonds but) has more van der Waals forces between molecules (1) [2]

Total [13]



(ii)

<i>Element</i>	<i>Initial Oxidation State</i>	<i>Final Oxidation State</i>
Nitrogen	-3	2
Hydrogen	1	1
Oxygen	0	-2

All three rows correct (2)
(1 mark if two rows correct)


Nitrogen oxidised as its oxidation state has increased (1) [3]

(iii) NH_3 has 3 bonding and 1 non bonding pair of electrons (1)

BF_3 has 3 bonding pairs only (1)

Electron pairs position themselves as far apart as possible
(to minimise repulsion) (1) [3]

(b) (i) A covalent bond where one of the atoms has donated both electrons
in the shared pair [1]



charge spread over ion (1)

correct bonding (1)

(iii) Tetrahedral (1) [2]

$109\frac{1}{2}^\circ$ (1) (accept 109°) [2]

(iv) Water is polar / a polar solvent (1)

Anion is attracted to H^{\oplus} / cation is attracted to O^{\ominus} (1) [2]

Total [14]

- Q.11** (a) (i) Lilac flame (1)
White solid / white fumes / potassium melts (1) [2]
- (ii) $4\text{K} + \text{O}_2 \longrightarrow 2\text{K}_2\text{O}$ [1]
- (iii) More reactive (1)
Electrons in rubidium lost more easily / ionisation energy is less /
explanation e.g. increased sheilding (1) [2]
(Need reason to get first mark but accept more reactive as reactivity
increases down group for 1 mark)
- (b) (i) No. moles = $\frac{0.098}{23} = 0.00426$ [1]
- (ii) Moles $\text{H}_2 = 0.00213$ (1)
Volume $\text{H}_2 = 0.00213 \times 24 = 0.0511 \text{ dm}^3$ (1) [2]
- (iii) Moles $\text{NaOH} = 0.00426$ (1)
Concentration $\text{NaOH} = \frac{0.00426}{0.200} = 0.0213 \text{ mol dm}^{-3}$ (1) [2]
- (c) (i) Do the experiment in a fume cupboard [1]
- (ii) I 6:6 [1]
II Electrostatic forces between the oppositely charged ions (1)
ionic bonds are / ionic lattice is very strong so large amount of
energy needed (1) [2]

Total [14]

Section B Total [70]