

# 4724 Core Mathematics 4

1 Attempt to factorise numerator and denominator M1  $\frac{A}{f(x)} + \frac{B}{g(x)}$ ; fg =  $6x^2 - 24x$   
 Any (part) factorisation of both num and denom A1 Corres identity/cover-up  
 Final answer =  $-\frac{5}{6x}, \frac{-5}{6x}, \frac{5}{-6x}, -\frac{5}{6}x^{-1}$  Not  $-\frac{5}{6x}$  A1  
3

---

2 Use parts with  $u = x, dv = \sec^2 x$  M1 result  $f(x) + / - \int g(x) dx$   
 Obtain correct result  $x \tan x - \int \tan x dx$  A1  
 $\int \tan x dx = k \ln |\sec x|$  or  $k \ln |\cos x|$ , where  $k = 1$  or  $-1$  B1 or  $k \ln |\sec x|$  or  $k \ln |\cos x|$   
 Final answer =  $x \tan x - \ln |\sec x| + c$  or  $x \tan x + \ln |\cos x| + c$  A1  
4

---

3 (i)  $1 + \frac{1}{2} \cdot 2x + \frac{\frac{1}{2} \cdot -\frac{1}{2}}{2} (4x^2 \text{ or } 2x^2) + \frac{\frac{1}{2} \cdot -\frac{1}{2} \cdot -\frac{3}{2}}{6} (8x^3 \text{ or } 2x^3)$  M1  
 $= 1 + x$  B1  
 $\dots -\frac{1}{2}x^2 + \frac{1}{2}x^3$  (AE fract coeffs) A1 (3) For both terms

(ii)  $(1+x)^{-3} = 1 - 3x + 6x^2 - 10x^3$  B1 or  $(1+x)^3 = 1 + 3x + 3x^2 + x^3$   
 Either attempt at their (i) multiplied by  $(1+x)^{-3}$  M1 or (i) long div by  $(1+x)^3$   
 $1 - 2x \dots \quad \sqrt{1 + (a-3)x}$  A1 f.t. (i) =  $1 + ax + bx^2 + cx^3$   
 $\dots + \frac{5}{2}x^2 \dots \quad \sqrt{(-3a+b+6)x^2}$  A1  
 $\dots - 2x^3 \quad \sqrt{(6a-3b+c-10)x^3}$  A1 (5) (AE fract.coeffs)

(iii)  $-\frac{1}{2} < x < \frac{1}{2}$ , or  $|x| < \frac{1}{2}$  B1 (1)  
9

---

4	Attempt to expand $(1 + \sin x)^2$ and integrate it	*M1	Minimum of $1 + \sin^2 x$
	Attempt to change $\sin^2 x$ into $f(\cos 2x)$	M1	
	Use $\sin^2 x = \frac{1}{2}(1 - \cos 2x)$	A1	dep M1 + M1
	Use $\int \cos 2x \, dx = \frac{1}{2} \sin 2x$	A1	dep M1 + M1
	Use limits correctly on an attempt at integration	dep* M1	Tolerate $g\left(\frac{1}{4}\pi\right) - 0$
	$\frac{3}{8}\pi - \sqrt{2} + \frac{7}{4}$ AE(3-term)F	A1	WW 1.51... → M1 A0

**6**

5 (i)	Attempt to connect $du$ and $dx$ , find $\frac{du}{dx}$ or $\frac{dx}{du}$	M1	But not e.g. $du = dx$
	Any correct relationship, however used, such as $dx = 2u \, du$	A1	or $\frac{du}{dx} = \frac{1}{2}x^{-1/2}$
	Subst with clear reduction ( $\geq 1$ inter step) to <b>AG</b>	A1 (3)	WWW

(ii)	Attempt partial fractions	M1	
	$\frac{2}{u} - \frac{2}{1+u}$	A1	
	$\sqrt{A \ln u + B \ln(1+u)}$	√A1	Based on $\frac{A}{u} + \frac{B}{1+u}$
	Attempt integ, change limits & use on $f(u)$	M1	or re-subst & use 1 & 9
	$\ln \frac{9}{4}$ AEexactF (e.g. $2 \ln 3 - 2 \ln 4 + 2 \ln 2$ )	A1 (5)	Not involving $\ln 1$

**8**

<p><b>6 (i)</b> Solve <math>0 = t - 3</math> &amp; subst into <math>x = t^2 - 6t + 4</math></p> <p>Obtain <math>x = -5</math></p> <p>N.B. If <b>(ii)</b> completed first, subst <math>y = 0</math> into their cartesian eqn (M1) &amp; find <math>x</math> (no f.t.) (A1)</p>	<p>M1</p> <p>A1 <b>(2)</b> <math>(-5,0)</math> need not be quoted</p> <p>(M1) &amp; find <math>x</math> (no f.t.) (A1)</p>
-----	
<p><b>(ii)</b> Attempt to eliminate <math>t</math></p> <p>Simplify to <math>x = y^2 - 5</math> ISW</p>	<p>M1</p> <p>A1 <b>(2)</b></p>
-----	
<p><b>(iii)</b> Attempt to find <math>\frac{dy}{dx}</math> or <math>\frac{dx}{dy}</math> from cartes or para form</p> <p>Obtain <math>\frac{dy}{dx} = \frac{1}{2t-6}</math> or <math>\frac{1}{2y}</math> or <math>(-)\frac{1}{2}(x+5)^{-\frac{1}{2}}</math></p> <p>If <math>t = 2</math>, <math>x = -4</math> and <math>y = -1</math></p> <p>Using their num <math>(x, y)</math> &amp; their num <math>\frac{dy}{dx}</math>, find tgt eqn</p> <p><math>x + 2y + 6 = 0</math> AEF(without fractions) ISW</p>	<p>M1 Award anywhere in Que</p> <p>A1</p> <p>B1 Awarded anywhere in <b>(iii)</b></p> <p>M1</p> <p>A1 <b>(5)</b></p>
<div style="border: 1px solid black; display: inline-block; padding: 2px 5px;">9</div>	

<p><b>7 (i)</b> Attempt direction vector between the 2 given points</p> <p>State eqn of line using format <math>(\mathbf{r}) = (\text{either end}) + s(\text{dir vec})</math></p> <p>Produce 2/3 eqns containing <math>t</math> and <math>s</math></p> <p>Solve giving <math>t = 3</math>, <math>s = -2</math> or 2 or <math>-1</math> or 1</p> <p>Show consistency</p> <p>Point of intersection = <math>(5,9,-1)</math></p>	<p>M1</p> <p>M1 's' can be 't'</p> <p>M1 2 different parameters</p> <p>A1</p> <p>B1</p> <p>A1 <b>(6)</b></p>
-----	
<p><b>(ii)</b> Correct method for scalar product of 'any' 2 vectors</p> <p>Correct method for magnitude of 'any' vector</p> <p>Use <math>\cos \theta = \frac{\mathbf{a} \cdot \mathbf{b}}{ \mathbf{a}   \mathbf{b} }</math> for the correct 2 vectors <math>\begin{pmatrix} 1 \\ 4 \\ -2 \end{pmatrix}</math> &amp; <math>\begin{pmatrix} 2 \\ -1 \\ 3 \end{pmatrix}</math></p> <p>62.2 (62.188157...) 1.09 (1.0853881)</p>	<p>M1 Vectors from this question</p> <p>M1 Vector from this question</p> <p>M1 Vects may be mults of dvs</p> <p>A1 <b>(4)</b></p>

- 8 (i)  $\frac{d}{dx}(y^3) = 3y^2 \frac{dy}{dx}$  B1
- Consider  $\frac{d}{dx}(xy)$  as a product M1
- $= x \frac{dy}{dx} + y$  A1 Tolerate omission of '6'
- $\frac{dy}{dx} = \frac{6y - 3x^2}{3y^2 - 6x}$  ISW AEF A1 (4)

- (ii)  $x^3 = 2^4$  or 16 and  $y^3 = 2^5$  or 32 \*B1
- Satisfactory conclusion dep\* B1 AG
- Substitute  $\left(2^{\frac{4}{3}}, 2^{\frac{5}{3}}\right)$  into their  $\frac{dy}{dx}$  M1 or the numerator of  $\frac{dy}{dx}$
- Show or use calc to demo that num = 0, ignore denom AG A1 (4)

- (iii) Substitute  $(a, a)$  into eqn of curve M1 & attempt to state 'a = ...'
- $a = 3$  only with clear ref to  $a \neq 0$  A1
- Substitute  $(3,3)$  or (their  $a$ , their  $a$ ) into their  $\frac{dy}{dx}$  M1
- 1 only WWW A1 (4) from (their  $a$ , their  $a$ )
- 12**

- 9 (i)  $\frac{d\theta}{dt} = \dots$  B1
- $k(160 - \theta)$  B1 (2) The 2 @ 'B1' are indep
- (ii) Separate variables with  $(160 - \theta)$  in denom; or invert \*M1  $\int \frac{1}{160 - \theta} d\theta = \int k, \frac{1}{k}, 1 dt$
- Indication that LHS =  $\ln f(\theta)$  A1 If wrong ln, final 3@A = 0
- RHS =  $kt$  or  $\frac{1}{k}t$  or  $t$  (+ c) A1
- Subst.  $t = 0, \theta = 20$  into equation containing 'c' dep\* M1
- Subst  $t = 5, \theta = 65$  into equation containing 'c' & 'k' dep\* M1
- $c = -\ln 140$  (-4.94) ISW A1
- $k = \frac{1}{5} \ln \frac{140}{95}$  ( $\approx 0.077$  or  $0.078$ ) ISW A1
- Using their 'c' & 'k', subst  $t = 10$  & evaluate  $\theta$  dep\* M1
- $\theta = 96(95.535714)$   $\left(95 \frac{15}{28}\right)$  A1 (9)

**11**