

Centre Number						Candidate Number				
Surname										
Other Names										
Candidate Signature										

For Examiner's Use	
Examiner's Initials	
Question	Mark
1	
2	
3	
4	
5	
TOTAL	



General Certificate of Education  
Advanced Level Examination  
June 2015

# Physics A

# PHYA5/1R

## Unit 5 Nuclear and Thermal Physics Section A

Thursday 18 June 2015 9.00 am to 10.45 am

**For this paper you must have:**

- a calculator
- a pencil and a ruler
- a question paper/answer book for Section B (enclosed).

**Time allowed**

- The total time for both sections of this paper is 1 hour 45 minutes.  
You are advised to spend approximately 55 minutes on this section.

**Instructions**

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided. Answers written in margins or on blank pages will not be marked.
- Do all rough work in this book. Cross through any work you do not want to be marked.
- Show all your working.

**Information**

- The marks for questions are shown in brackets.
- The maximum mark for this section is 40.
- You are expected to use a calculator, where appropriate.
- A *Data and Formulae Booklet* is provided as a loose insert in Section B.
- You will be marked on your ability to:
  - use good English
  - organise information clearly
  - use specialist vocabulary where appropriate.



J U N 1 5 P H Y A 5 1 R 0 1

**Section A**

The maximum mark for this section is 40.  
You are advised to spend approximately 55 minutes on this section.

- 1 (a)** Which ionizing radiation produces the greatest number of ion pairs per mm in air?  
Tick (✓) the correct answer.

**[1 mark]**

$\alpha$ particles	
$\beta$ particles	
$\gamma$ rays	
X-rays	

- 1 (b) (i)** Complete **Table 1** below showing the typical maximum range in air for  $\alpha$  and  $\beta$  particles.

**[2 marks]**

**Table 1**

Type of radiation	Typical range in air / m
$\alpha$	
$\beta$	

- 1 (b) (ii)**  $\gamma$  rays have a range of at least 1 km in air.  
However, a  $\gamma$  ray detector placed 0.5 m from a  $\gamma$  ray source detects a noticeably smaller count-rate as it is moved a few centimetres further away from the source.

Explain this observation.

**[1 mark]**

.....

.....

.....



**1 (c)** Following an accident, a room is contaminated with dust containing americium which is an  $\alpha$ -emitter.

Explain the most hazardous aspect of the presence of this dust to an unprotected human entering the room.

**[2 marks]**

.....

.....

.....

.....

6

**Turn over for the next question**

**Turn over ▶**



**2 (a)** Scattering experiments are used to investigate the nuclei of gold atoms. In one experiment, alpha particles, all of the same energy (monoenergetic), are incident on a foil made from a single isotope of gold.

**2 (a) (i)** State the main interaction when an alpha particle is scattered by a gold nucleus. **[1 mark]**

.....

.....

**2 (a) (ii)** The gold foil is replaced with another foil of the same size made from a mixture of isotopes of gold. Nothing else in the experiment is changed.

Explain whether or not the scattering distribution of the monoenergetic alpha particles remains the same.

**[1 mark]**


.....

.....

.....

**2 (b)** Data from alpha-particle scattering experiments using elements other than gold allow scientists to relate the radius  $R$ , of a nucleus, to its nucleon number,  $A$ . **Figure 1** shows the relationship obtained from the data in a graphical form, which obeys the relationship  $R = r_0 A^{\frac{1}{3}}$ .

**Figure 1**



**2 (b) (i)** Use information from **Figure 1** to show that  $r_0$  is about  $1.4 \times 10^{-15}$  m.

**[1 mark]**

**2 (b) (ii)** Show that the radius of a  ${}_{23}^{51}\text{V}$  nucleus is about  $5 \times 10^{-15}$  m.

**[2 marks]**

**2 (c)** Calculate the density of a  ${}_{23}^{51}\text{V}$  nucleus.

State an appropriate unit for your answer.

**[3 marks]**

density ..... unit .....



**3** A rod made from uranium-238 ( ${}^{238}_{92}\text{U}$ ) is placed in the core of a nuclear reactor where it absorbs free neutrons.

When a nucleus of uranium-238 absorbs a neutron it becomes unstable and decays to neptunium-239 ( ${}^{239}_{93}\text{Np}$ ), which in turn decays to plutonium-239 ( ${}^{239}_{94}\text{Pu}$ ).


**3 (a)** Write down the nuclear equation that represents the decay of neptunium-239 into plutonium-239.

[2 marks]

**3 (b)** A sample of the rod is removed from the core and its radiation is monitored from time  $t = 0$  s.

The variation of the activity with time is shown in **Figure 2**.

**Figure 2**



**3 (b) (i)** Show that the decay constant of the sample is about  $3.4 \times 10^{-6} \text{ s}^{-1}$ .

**[2 marks]**

**3 (b) (ii)** Assume that the activity shown in **Figure 2** comes only from the decay of neptunium.

Estimate the number of neptunium nuclei present in the sample at time  $t = 5.0 \times 10^5 \text{ s}$ .

**[1 mark]**

number of nuclei .....

**Question 3 continues on the next page**

**Turn over ►**



**3 (c) (i)** A chain reaction is maintained in the core of a thermal nuclear reactor that is operating normally.

Explain what is meant by a chain reaction, naming the materials and particles involved.

**[2 marks]**

.....  
.....  
.....  
.....

**3 (c) (ii)** Explain the purpose of a moderator in a thermal nuclear reactor.

**[2 marks]**

.....  
.....  
.....  
.....  
.....  
.....

**3 (c) (iii)** Substantial shielding around the core protects nearby workers from the most hazardous radiations. Radiation from the core includes  $\alpha$  and  $\beta$  particles,  $\gamma$  rays, X-rays, neutrons and neutrinos.

Explain why the shielding becomes radioactive.

**[2 marks]**

.....  
.....  
.....  
.....





**4 (a)** Lead has a specific heat capacity of  $130 \text{ J kg}^{-1} \text{ K}^{-1}$ .

Explain what is meant by this statement.

**[1 mark]**

.....

.....

.....

.....

**4 (b)** Lead of mass  $0.75 \text{ kg}$  is heated from  $21 \text{ }^\circ\text{C}$  to its melting point and continues to be heated until it has all melted.

Calculate how much energy is supplied to the lead.  
Give your answer to an appropriate number of significant figures.

melting point of lead =  $327.5 \text{ }^\circ\text{C}$   
specific latent heat of fusion of lead =  $23\,000 \text{ J kg}^{-1}$

**[3 marks]**

energy supplied ..... J

4

**Turn over ►**





.....

.....

.....

.....

.....

.....

.....

.....

**Question 5 continues on the next page**

**Turn over ▶**



**5 (b) (i)** State **two** assumptions about the **movement** of molecules that are used when deriving the equation of state,  $pV = \frac{1}{3} Nm (c_{\text{rms}})^2$  for an ideal gas.

**[2 marks]**

1 .....

.....

2 .....

.....

**5 (b) (ii)** Three molecules move at the speeds shown in **Table 2**.

**Table 2**

molecule	speed / m s <sup>-1</sup>
1	2000
2	3000
3	7000

Calculate their mean square speed.

**[1 mark]**

mean square speed ..... m<sup>2</sup> s<sup>-2</sup>



**5 (c)** The average molecular kinetic energy of an ideal gas is  $6.6 \times 10^{-21}$  J.  
Calculate the temperature of the gas.

**[2 marks]**

temperature ..... K

11

**END OF SECTION A**

**Turn over ►**



**There are no questions printed on this page**

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**



**There are no questions printed on this page**

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**



**There are no questions printed on this page**

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**

