

## PH5

## SECTION A

Question			Marking details	Marks Available
1	(a)	(i)	$84.6 \times 10^{-9}$ [C] [for 4.7 nF] (1) $73.8 \times 10^{-9}$ [C][and 73.8 nC or clearly stated same for other 8.2 nF] (1)	2
		(ii)	$E = \frac{1}{2}CV^2$ or other equation used correctly or $C$ total = 8.8 nF (1) Answer = $1.43 \times 10^{-6}$ [J] <b>ecf</b> on $Q$ but not $V$ (1)	2
	(b)	(i)	Points taken from the curve e.g. $Q_0 = 85$ nC and (50 ms, 6 nC) (or $85$ nC/ $e = 31$ nC) (1)  Values substituted correctly e.g. $6 = 85e^{-0.05/CR}$ or $CR = 18$ ms (1)  Answer $R = 3.8 \times 10^6$ [ $\Omega$ ] (1) <b>Award 1 mark</b> for use of $\frac{\Delta Q}{t}$ <b>t</b> or $11$ M $\Omega$	3
		(ii)	$I = \frac{V}{R}$ used or tangent drawn at $t = 0$ (1)  Answer = $4.7 \times 10^{-6}$ [A] <b>ecf</b> (1)	2
		(iii)	After $41 \pm 1$ ms 10% charge left [or 90% discharged] Or other valid method e.g. taking logs and getting time (1)  $83 \times 10^{-3}$ [s] (first step can be implied) <b>ecf</b> (1)	2
	<b>Question 1 Total</b>			<b>[11]</b>


Question		Marking details	Marks Available
2	(a)	(i) 0	1
		(ii) $\phi = B \times l^2(1)$ Answer = $4.32 \times 10^{-5}$ [Wb] (1)	2
	(b)	Change in flux or Faraday's law gives emf (1) Complete circuit or accept emf gives current (1) <b>Award 1 mark only for:</b> Current due to Faraday's law	2
	(c)	Force / current / emf <b>opposes the change</b> (1) Force on PQ opposite to SR <b>or</b> the force is clockwise (1)	2
	(d)	$I = \frac{V}{R}$ used (1) $A = \pi \frac{d^2}{4}$ <i>or</i> $\pi \times 3^2 (\times 10^{-6})$ i.e. $\pi^2$ used (1) $R = \frac{\rho \times l}{A}$ used (1) $V = \frac{\Delta N \phi}{\Delta t}$ used (1) Answer = 0.19 [A] <b>ecf</b> on $\phi$ and $\pi d^2$ (1)	5
		<b>Question 2 Total</b>	<b>[12]</b>

Question		Marking details	Marks Available
3	(a)	<p>Low <math>A</math> numbers do fusion (or arrow / label used) (1)</p> <p>High <math>A</math> numbers do fission (or arrow / label used) (1)</p> <p>Moving toward high BE/nucleon (around Fe-56) or Fe-56 is the most stable (or low PE/nucleon or accept work done by strong nuclear force) (1)</p> <p>Higher BE/nucleon is more stable (or low PE/nucleon more stable or more work done more stable) (1)</p>	4
	(b)	<p><math>1.1 \pm 0.1</math> MeV identified from graph for <math>{}^2_1\text{H}</math> (1)</p> <p><math>\times 2 = 2.2</math> [MeV] <b>ecf</b> (1)</p>	2
	(c)	<p><math>7.6 \pm 0.2, 8.4 \pm 0.2, 8.7 \pm 0.2</math> (1)</p> <p>Correct multipliers for each i.e. <math>235 \times 7.6, 137 \times 8.4, 96 \times 8.7</math> (1)</p> <p>RHS – LHS or reverse (1)</p> <p>Correct answer e.g. 201 MeV <b>UNIT mark</b> (1) [dependent on BE/<math>A</math> approximations]</p>	4
		<b>Question 3 Total</b>	<b>[10]</b>

Question		Marking details	Marks Available
4	(a)	360 ± 10 [minutes]	1
	(b)	No [significant] drop after paper [no α] (1)  [Small drop after aluminium] so small amount of $\gamma$ being absorbed / most $\gamma$ passes through i.e. could be $\beta$ but some $\gamma$ would be absorbed ok <b>Or</b> accept drop could be attributable to randomness of decay (1)  $\gamma$ present because something gets through 3 mm Al <b>or</b> $\gamma$ present because <b>bigger</b> drop after 10 cm Pb [than 3 mm Al] <b>or</b> $\gamma$ present because <u>only</u> absorbed by the Pb (1)	3
	(c)	Activity = $\frac{450}{0.006} = 75\,000$ (1)  Activity = $\lambda N$ or $t_{\frac{1}{2}} = \frac{\ln 2}{\lambda}$ used (1)  $N = 2.34 \times 10^9$ (1)  Mass = $99 \times 1.66 \times 10^{-27} \times 2.34 \times 10^9 = 3.84 \times 10^{-16}$ kg <b>UNIT mark</b> (1) <b>ecf</b> on $A$ and $t_{\frac{1}{2}}$ and $N$	4
	<b>Question 4 Total</b>		<b>[8]</b>

Question		Marking details	Marks Available
5	(a)	$n = \frac{12\,000}{1.8} (1)$ $B = \mu_0 nI = 0.019 \text{ [T]} (1)$	2
	(b)	<p>(i) Either <math>Bev = \frac{mv^2}{r}</math> or <math>Bev = m\omega^2 r</math> (1)</p> <p><math>v = \omega r</math> and <math>\omega = 2\pi f</math> quoted (1)</p> <p>Clear algebra (if not immediately understandable then not clear) (1)</p> <p>(ii) <math>f = \frac{3.3 \times 6 \times 1.6 \times 10^{-19}}{2\pi \times 12 \times 1.66 \times 10^{-27}} (1)</math></p> <p>Answer = <math>25.3 \times 10^6</math> [Hz] (1)</p> <p>(iii) <math>6e \times 14.5 \text{ kV} \times 24 [= 2.09 \text{ MeV}] (1)</math></p> <p>Conversion to J i.e. look out for <math>\times 1.6 \times 10^{-19}</math> (1)</p> <p>Equating some related energy to <math>\frac{1}{2}mv^2</math> e.g. <math>\frac{1}{2}mv^2 = 14\,500</math> (1)</p> <p>Answer = <math>5.8 \times 10^6</math> [<math>\text{m s}^{-1}</math>] (1)</p> <p>(<b>ecf</b> on these values only <math>2.4 \times 10^6</math> and <math>4.1 \times 10^6</math> which correspond to <math>q = 1e</math> and 12 kicks respectively)</p> <p><b>Question 5 Total</b></p>	3            <b>[11]</b>

Question		Marking details	Marks Available
6	(a)	(i) +ve correct	1
		(ii) voltmeter correct	1
	(b)	$V = Ed$ or $V_H = Bvd$ (1) $Bev = eE$ quoted or $d = 5 \times 10^{-3}$ (1) Answer = $6.3 \times 10^{-6}$ [V] (1)	3
	(c)	Electrons do not move in the direction of the Hall field (or accept in the direction of the Hall voltage)	1
	(d)	Correct use of $I = nAve$ or $n = \frac{BI}{V_H te}$ (or equiv equation) (1) Answer $I = 0.30 \times 10^{-3}$ [A] (1)	2
		<b>Question 6 Total</b>	<b>[8]</b>



**SECTION B**



Question		Marking details	Marks Available
7	(a)	<p>KE given to water (1)</p> <p>Air resistance (1)</p> <p><b>Award 2 marks for:</b>                      Water runs out before pressure drops to 1 atm</p> <p>Accept PE given to water [even though water runs out in 1.6 m]</p> <p>Accept viscosity [or friction] of water</p> <p>Accept KE given to bottle [only 1/10th of mass of ball]</p> <p>Don't accept heat or sound</p>	2
	(b)	<p><math>v = -20 \ln \left( 1 - \frac{5.9}{1.5} \times 0.175 \right)</math> (1)</p> <p><math>v = 23.3 \text{ [m s}^{-1}\text{]}</math> (1)</p>	2
	(c)	<p>kg s<sup>-1</sup>, m<sup>2</sup>, kg m<sup>-3</sup> and m s<sup>-1</sup>                      (0 or 1 unit correct 0 marks)                      (2 or 3 units correct 1 mark)                      (All 4 correct 2 marks)                      If all 4 units correct but algebra is incorrect then deduct 1 mark</p>	2
	(d)	<p>Rearranging i.e. <math>u = \frac{\Delta m / \Delta t}{\pi r^2 \rho}</math> (1)</p> <p>Answer = 25 [m s<sup>-1</sup>] (1)</p>	2
	(e)	<p><b>Any 3 × (1) from:</b></p> <p>Rocket equation assumes constant <math>u</math></p> <p>Volume increasing so pressure decreasing</p> <p>Pressure is decreasing so <math>u</math> is decreasing</p> <p>Thrust is decreasing</p>	3

Question		Marking details	Marks Available
	(f)	$\Delta p = 6.8 \times 10^5 \text{ [Pa]} \text{ (1)}$ $u = \sqrt{\frac{2 \times 6.8 \times 10^5}{1000}} = 36.9 \text{ [m s}^{-1}\text{]} \text{ (1)}$	2
	(g)	<p><math>mg</math> - weight or gravitational force <b>and</b> <math>0.0107v^2</math> - air resistance / drag accept <u>skin</u> / <u>air</u> friction (1)</p> <p>Weight decreases (1)</p> <p>Air resistance increases (1)</p>	3
	(h)	<p>Squaring equation 6 or rearranging equation 5 (1)</p> <p>Convincing algebra (1)</p>	2
	(i)	I Fast process or no time for heat to flow or equivalent	1
		II The gas does <u>work</u> so <u>internal energy</u> decreases	1
		<b>Or <math>\Delta U = -W</math></b>	
		<b>Question 7 Total</b>	<b>[20]</b>



SECTION C

Question		Marking details	Marks Available
8	(a)	(i) $\omega L = \frac{1}{\omega C}$ or $f = \frac{1}{2\pi\sqrt{LC}}$ (1) $= \frac{1}{2\pi\sqrt{0.022 \times 15 \times 10^{-9}}}$ or $f = \frac{1}{2\pi\sqrt{0.022 \times 90 \times 10^{-9}}}$ (1) 8 761 [Hz] and 3 577 [Hz] (1)	3
		(ii) $Q = \frac{2\pi fL}{R}$ or $Q = \frac{1}{R}\sqrt{\frac{L}{C}}$ (1) Correct matching of $f$ and $R$ or $R$ and $C$ (1) 61 (1) 10 (1)	4
	(b)	$Z = \sqrt{\left(\omega L - \frac{1}{\omega C}\right)^2 + R^2}$ used (1) $I = \frac{3.2}{Z}$ i.e. $I = \frac{V}{Z}$ used (1) 1.7 [mA] (1)	3

Question			Marking details	Marks Available
8	(c)	(i)	 <p>Shapes similar and asymptotic to original (1)</p> <p>20 <math>\Omega</math> with higher peak and 50 <math>\Omega</math> with lower (implied if not labelled) (1)</p> <p>20 <math>\Omega</math> with peak current of 160 mA (1)</p> <p>50 <math>\Omega</math> with peak current of 64 mA (1)</p>	4
		(ii)	<p><math>\omega L</math> increases (wrt <math>f</math>) or graph (1)</p> <p><math>\frac{1}{\omega C}</math> decreases (wrt <math>f</math>) (1)</p> <p>low frequency behaviour explained e.g. <math>X_C</math> very large, <math>Z</math> large at low freq (1)</p>  <p>high frequency behaviour explained e.g. <math>X_L</math> very large, <math>Z</math> large at high freq (1)</p> <p><math>\omega L - \frac{1}{\omega C} = 0</math> at resonance making <math>Z</math> a minimum (1)</p> <p><math>I \propto \frac{1}{Z}</math> or equivalent equation etc. (1)</p>	6
<b>Question 8 Total</b>				<b>[20]</b>

Question		Marking details	Marks Available	
9	(a)	(i) Diagram showing either angle (accept $\theta$ ) or baseline (1)  Attempt to use $b = r\theta$ and indication that $\theta$ must be in radians, Or attempt to use $b/2 = r \tan (\theta/2)$ or equiv. Or by implication. (1)  $r = 1.96 \times 10^6$ [km] or $1.92 \times 10^6$ [km] or convincing answer. (1)	3	
		(ii) Showed comet (far) beyond Moon. (1) But according to Aristotle nothing changes beyond Moon [yet comet was new – and went away]. (1)	2	
	(b)	(i) Diagram showing relevant areas (1) $r_P, r_A, v_P\Delta t, v_A\Delta t$ marked on diagram or meanings otherwise shown (1) $(\frac{1}{2})r_Pv_P\Delta t = (\frac{1}{2})r_Av_A\Delta t$ or equivalent (1)	3	
		(ii) Use of $\frac{v_P}{v_A} = \frac{r_A}{r_P}$ [= 1.10] or by implication (1) 10 % [increase] (1)	2	
		(iii) Explicit use of $\frac{mv^2}{r}$ (1)  $\frac{v_P}{v_A} = \frac{r_A}{r_P}$ used convincingly to give $\frac{F_P}{F_A} = \frac{r_A^2}{r_P^2}$ or equiv. (1)	2	
	(c)	(i) Towards S or equivalent	1	
		(ii) Any 3 of ... • Sun at S, planet's path ABCDEF... • If time interval is shrunk, path becomes smooth • Equal areas swept out in equal times • Showed that for an elliptical path ... • ... force had to vary as inverse square of Sun-planet distance	3	
		(iii) Planets swirled in whirlpool (vortex) around the Sun (1)  Any 2 of .... (2) • easy to understand • gave a <i>mechanism</i> • Newton didn't say what <i>caused</i> gravitation  Descartes' vortex theory can't be made to account for actual orbits [that is for Kepler's laws] or Newton's theory accounted for so many phenomena so economically [or similar point] (1)	4	
	<b>Question 9 Total</b>			<b>[20]</b>

Question			Marking details	Marks Available
10	(a)	(i)	Strong (covalent) bonds between ions in structure. Accept molecules arranged irregularly or amorphous structure present (1) [No dislocations present] so no slip (accept no movement of dislocations). Accept different sized atoms seize up the structure (1) Do not accept 'untangle'.	2
		(ii)	Scratches (on surface) weaken material or break surface bonds. Scratches have stress concentrations at their tips- can be awarded from diagram. Cracks propagate through material. Correct direction of bending is to open the crack [ANY 2] (credit well annotated diagrams) [Local] stress cannot be relieved by slip / plastic flow/ dislocation movement.	2
		(iii)	Compression (1) [Do not accept 'stress'] More difficult for cracks to develop/ widen/ propagate (1)	2
	(b)	(i)	Gradient shown = 80 GPa e.g. $\frac{800 \times 10^6}{0.01}$ seen.	1
		(ii)	Area under graph = $[\frac{1}{2} \times 0.01 \times 800 \times 10^6 + \frac{1}{2} \times 0.08 \times 100 \times 10^6 + 0.08 \times 800 \times 10^6] = 72 \text{ [MJ m}^{-3}]$ (1) Volume = $\pi \times (1.25 \times 10^{-3})^2 \times 2.5 = 12.3 \times 10^{-6} \text{ [m}^3]$ (1) Work done = $72 \times 10^6 \times 12.3 \times 10^{-6} = 884 \text{ [J]}$ (1) ( <b>ecf</b> on both area and volume).	3
		(iii)	Initial straight line of same gradient. (1) Yield point at 1 000 MPa. (1) Linear plastic region of small slope (accept zero slope) stopping at 5% strain. (1)	3
		(iv)	(I) Creep: [ <b>Gradual/slow/Over time</b> ] AND [ <b>extension/stretching/deformity or increase in strain</b> ] (1) (under a constant load). Necking: <b>Localised</b> (or reference to 'section' or 'region') <b>thinning</b> (of structure/material before breaking- accept diagram) (1)	2
		(II)	Same shaped curve but steeper gradient (1) Stopped at 15% and $t < 400$ hrs (approx.) (1)	2
	(v)	Repeated bending, stretching or hammering of metal alloy (1) Dislocations become tangled / traffic jam effect <b>or</b> new dislocations created (1) Stopping each other from moving (or inhibiting plastic deformation or collect at grain boundaries) (1)	3	
	<b>Question 10 Total</b>			<b>[20]</b>

Question		Marking details	Marks Available
11	(a)	(i) A/B/D	1
		(ii) C	1
		(iii) A	1
	(b)	$V = \frac{hc}{e\lambda}$ (must rearrange) (1)	2
		$6.2 \times 10^4 \text{ V}$ (must have valid unit) (1)	
	(c)	Reduces scattering/ spreading accept 'ensures (X-rays) are all parallel / perpendicular [to the patient] (1) [leading to] sharper image / better resolution (1)	2
	(d)	(i) Radio (waves)	1
		(ii) Cause Hydrogen atoms to resonate (1) Flip alignment producing a magnetic field (1)	2
		(iii) Not good for dense objects/bone/ Uncomfortable/ Claustrophobic/cannot be used with pacemakers/ expensive	1
	(e)	(i) Depolarization of ventricles/ repolarisation of atria (1) Contraction of ventricles (1)	2
		(ii) Repolarization of ventricles (1) Relaxation of ventricles/ ventricles return to normal (1) Do NOT accept ventricles expand	2
	(f)	(i) Doppler	1
		(ii) $0.4 \times \frac{1500}{500} = 2v$ (1) $v = 0.6 \text{ [m s}^{-1}\text{]}$ allow 1 mark only for $1.2 \text{ m s}^{-1}$ (1)	2
(g)	(i) Gamma / $\gamma$	1	
	(ii) Very expensive/need a cyclotron / particle accelerator Ignore any reference to radiation dose	1	
<b>Question 11 Total</b>			<b>[20]</b>

Question		Marking details	Marks Available
12	(a)	<p>Any 3 of:</p> <p>More than one isotope of uranium</p> <p>Only U-235 fissile / U-238 unsuitable for fission</p> <p>Need higher concentration of U-235</p> <p>Get rid of U-238 since neutron absorber/'sink'</p>	3
	(b)	<p>More fissile nuclei obtained (1)</p> <p>U-238 captures neutrons <b>or</b> decays to plutonium 239 via beta (1)</p>	2
	(c)	<p>Any 2 of:</p> <p>More easily controllable or no chain reaction</p> <p>Hydrogen &amp; deuterium more plentiful / more easily sourced / can be extracted from sea water</p> <p>No [long term] radioactive products.</p> <p>More energy released per fusion</p>	2
	(d)	<p>(i) Overcome electrostatic repulsion /forces (1)</p> <p>Needs high / enough (K)E (or velocity, speed) of deuterium/tritium (1)</p> <p><u>KE proportional to temperature</u> or only tail end of distribution (1) with high enough energy</p>	3

Question		Marking details	Marks Available
	(ii)	$n = or \geq \frac{3.5 \times 10^{28}}{0.9 \times 120000000} (1)$ <p>Multiplying by <math>2.5 \times 1.67 \times 10^{-27} (1)</math></p> <p>Answer = <math>1.35 \times 10^{-6} [kg m^{-3}] (1)</math></p>	3
	(e)	<p><math>9 \times 10^{16} (1)</math></p> <p><math>\times 2 = 18 \times 10^{16} [J] (1)</math></p>	2
	(f)	<p>Coal - <b>any 2 of:</b> acid rain, CO<sub>2</sub>, non-renewable, lasts hundreds of years, increase asthma, damage to buildings/trees/etc, global warming (1)</p> <p>Natural gas - <b>any 2 of:</b> less acid rain, CO<sub>2</sub>, non-renewable lasts tens or hundreds of years, global warming (1)</p> <p>Biomass - <b>any 2 of:</b> acid rain, CO<sub>2</sub> neutral (roughly), renewable/lasts millions of years, increase asthma, damage to buildings/trees/etc, (roughly) no global warming, large land area needed (1)</p> <p>Uranium-235 <b>any 2 of:</b> no acid rain, no CO<sub>2</sub>, lasts thousands of years, no increase asthma, no damage to buildings/trees/etc, (little or ) no global warming, leak or explosion risk, disposal of waste, hazardous waste, large energy output per kg of fuel (1)</p> <p>Wind <b>any 2 of:</b> no acid rain, no CO<sub>2</sub>, renewable/lasts millions of years, no increase asthma, no damage to buildings/trees/etc, no global warming, low power, weather dependent/unreliable, large area (1)</p> <p>(ugly/kills birds/noisy/disrupts TV signals OK but only 1 point max for these)</p> <p>NOTE: No marks for cheap/expensive - eliminated in stem!!</p>	5
<b>Question 12 Total</b>			<b>[20]</b>