

Mark Scheme (Results)

Summer 2009

GCE

GCE Mathematics (6686/01)

June 2009
6686 Statistics S4
Mark Scheme

Question Number	Scheme	Marks
Q1	<p>$H_0: \mu = 5; H_1: \mu < 5$</p> <p>CR: $t_9(0.01) > 2.821$</p> <p>$\bar{x} = 4.91$</p> $s^2 = \frac{1}{9} \left(241.2 - \frac{49.1^2}{10} \right) = 0.0132222$ $t = \frac{ 4.91 - 5 }{\frac{\sqrt{0.013222}}{\sqrt{10}}} = \pm 2.475$ <p>Since 2.475 is not in the critical region there is insufficient evidence to reject H_0 and conclude that the mean diameter of the bolts is not less than (not equal to) 5mm.</p>	<p style="text-align: center;">both</p> <p>B1 B1 B1</p> <p>s= awrt 0.115 M1 A1</p> <p>2.47 – 2.48 M1 A1</p> <p>A1ft</p> <p style="text-align: right;">[8]</p>

Question Number	Scheme	Marks
Q2	<p>(a) The differences are normally distributed</p> <p>(b) The data is collected in pairs or small sample size and variance unknown or samples not independent</p> <p>(c) $d: 2.5, 1.6, 1.6, -1.9, -0.6, 4.5$ at least 2 correct $(\Sigma d = 7.7, \Sigma d^2 = 35.59) \bar{d} = \pm 1.2833, sd = 2.2675. (Var = 5.141)$ $H_0: \mu_d = 0, H_1: \mu_d > 0$ ($H_1: \mu_d < 0$ if $d = -2.5, -1.6, -1.6$ etc) both depend on their d's $t = \frac{\pm 1.2833\sqrt{6}}{2.2675} = \pm 1.386\dots$ formula and substitution, 1.38 – 1.39 Critical value $t_5(5\%) = 2.015$ (1 tail) Not significant. Insufficient evidence to support that the device reduces CO₂ emissions.</p> <p>(d) The idea that the device reduces CO₂ emissions has been rejected when in fact it does reduce emissions. OR Concluding that the device does not reduce emissions when in fact it does (if not in context can get B1 only)</p> <p>(b) Allow because the same car has been used (c) awrt $\pm 1.28, 2.27$</p>	<p>B1 (1)</p> <p>B1 (1)</p> <p>M1 A1, A1 B1 M1, A1 B1 A1 ft (8)</p> <p>B1 B1 (2)</p> <p>[12]</p>

Question Number	Scheme	Marks
3	(a) Size is the probability of H_0 being rejected when it is in fact true. or $P(\text{reject } H_0 / H_0 \text{ is true})$ oe	B1 (1)
	(b) The power of the test is the probability of rejecting H_0 when H_1 is true. or $P(\text{rejecting } H_0 / H_1 \text{ is true}) / P(\text{rejecting } H_0 / H_0 \text{ is false})$ oe	B1 (1)
	(c) $X \sim B(12, 0.5)$ $P(X \leq 2) = 0.0193$ $P(X \geq 10) = 0.0193$	B1 M1
	\therefore critical region is $\{X \leq 2 \cup X \geq 10\}$	A1A1 (4)
	(d)(i) $P(\text{Type II error}) = P(3 \leq X \leq 9 \mid p = 0.4)$ $= P(X \leq 9) - P(X \leq 2)$ $= 0.9972 - 0.0834$ $= 0.9138$	M1 M1dep A1
	(ii) Power = $1 - 0.9138$	B1 ft (4)
	(e) = 0.0862 Increase the sample size Increase the significance level/larger critical region	B1 B1 (2)
	Notes (d) (i) first M1 for either correct area or follow through from their critical region 2nd M1 dependent on them having the first M1. for finding their area correctly A1 cao (ii) B1 follow through from their (i)	[12]

Question Number	Scheme	Marks
<p>Q4 (a)</p> <p>$H_0 : \sigma_A^2 = \sigma_B^2, H_1 : \sigma_A^2 \neq \sigma_B^2$</p> <p>critical values $F_{12,8}=3.28$ and $\frac{1}{F_{8,12}} = 0.35$</p> <p>$\frac{s_B^2}{s_A^2} = 2.40 \left(\frac{s_A^2}{s_B^2} = 0.416 \right)$</p> <p>Since 2.40 (0.416) is not in the critical region we accept H_0 and conclude there is no evidence that the two variances are different.</p> <p>(b)</p> <p>$S_p^2 = \frac{8 \times 1.02 + 12 \times 2.45}{20}$</p> <p>$= 1.878$</p> <p>$(27.94 - 25.54) \pm 2.086 \times \sqrt{1.878} \times \sqrt{\frac{1}{9} + \frac{1}{13}}$</p> <p>(1.16, 3.64)</p> <p>(c) To calculate the confidence interval the variances need to be equal. In part (a) the test showed they are equal.</p>		<p>B1</p> <p>B1</p> <p>M1A1</p> <p>A1ft (5)</p> <p>M1</p> <p>A1</p> <p>B1M1 A1ft</p> <p>A1 A1 (7)</p> <p>B1</p> <p>B1 (2)</p> <p>[14]</p>

Question Number	Scheme	Marks
Q5 (a)	95% confidence interval for μ is 2.145 $560 \pm t_{14}(2.5\%) \sqrt{\frac{25.2}{15}} = 560 \pm 2.145 \sqrt{\frac{25.2}{15}} = (557.2, 562.8)$	B1 M1 A1 A1 (4)
(b)	95% confidence interval for σ^2 is $5.629 < \frac{14 \times 25.2}{\sigma^2} < 26.119$ $\sigma^2 < 62.675 \quad \sigma^2 > 13.507$ $13.507 < \sigma^2 < 62.675$ awrt 13.5, 62.7	B1, M1, B1 A1, A1 (5)
(c)	Require $P(X > 565) = P\left(Z > \frac{565 - \mu}{\sigma}\right)$ to be as large as possible OR $\frac{565 - \mu}{\sigma}$ to be as small as possible; both imply highest σ and μ . $\frac{565 - 562.8}{\sqrt{62.675}} = 0.28$ $P(Z > 0.28) = 1 - 0.6103 = 0.3897$ awrt 0.39 – 0.40	M1 M1A1 M1 A1 (5) [14]
	(c) M1 for using their largest σ and μ M1 for using $\frac{x - \mu}{\sigma}$ M1 1 – their prob	

Question Number	Scheme	Marks
Q6	<p>(a) $E(\frac{2}{3}X_1 + \frac{1}{2}X_2 + \frac{5}{6}X_3) = \frac{2}{3} \times \frac{k}{2} + \frac{1}{2} \times \frac{k}{2} + \frac{5}{6} \times \frac{k}{2} = k$ $E(X_1 + X_2 + X_3) = k \Rightarrow$ unbiased</p> <p>(b) $E(aX_1 + bX_2) = a\frac{k}{2} + b\frac{k}{2} = k$ $a + b = 2$ $\text{Var}(aX_1 + bX_2) = a^2\frac{k^2}{12} + b^2\frac{k^2}{12}$ $= a^2\frac{k^2}{12} + (2-a)^2\frac{k^2}{12}$ $= (2a^2 - 4a + 4)\frac{k^2}{12}$ $= (a^2 - 2a + 2)\frac{k^2}{6}$ (*) since answer given</p> <p>(c) Min value when $(2a-2)\frac{k^2}{6} = 0$ $\frac{d}{da}(\text{Var}) = 0$, all correct, condone missing $\frac{k^2}{6}$ $\Rightarrow 2a - 2 = 0$ $a = 1, b = 1.$ $\frac{d^2(\text{Var})}{da^2} = \frac{2k^2}{6} > 0$ since $k^2 > 0$ therefore it is a minimum min variance $= (1 - 2 + 2)\frac{k^2}{6}$ $= \frac{k^2}{6}$</p> <p>Alternative $\frac{k^2}{6}(a-1)^2 - \frac{k^2}{6} + \frac{2k^2}{6}$ $\frac{k^2}{6}(a-1)^2 + \frac{k^2}{6}$ Min when $\frac{k^2}{6}(a-1)^2 = 0$ $a = 1, b = 1$ min var $= k^2/6$</p>	<p>M1 A1 B1 (3)</p> <p>M1 A1 M1A1 M1</p> <p>A1 cso (6)</p> <p>M1A1 A1A1 M1</p> <p>B1 (6)</p> <p>M1 A1 M1 A1A1 B1</p>