

# 4723 Core Mathematics 3

1 Obtain integral of form  $k(2x-7)^{-1}$  M1 any constant  $k$   
 Obtain correct  $-5(2x-7)^{-1}$  A1 or equiv  
 Include ... +  $c$  B1 **3** at least once; following any integral  
3

2 (i) Use  $\sin 2\theta = 2\sin\theta\cos\theta$  B1  
 Attempt value of  $\sin\theta$  from  $k\sin\theta\cos\theta = 5\cos\theta$  M1 any constant  $k$ ; or equiv  
 Obtain  $\frac{5}{12}$  A1 **3** or exact equiv; ignore subsequent work

(ii) Use  $\operatorname{cosec}\theta = \frac{1}{\sin\theta}$  or  $\operatorname{cosec}^2\theta = 1 + \cot^2\theta$  B1 or equiv  
 Attempt to produce equation involving  $\cos\theta$  only M1 using  $\sin^2\theta = \pm 1 \pm \cos^2\theta$  or equiv  
 Obtain  $3\cos^2\theta + 8\cos\theta - 3 = 0$  A1 or equiv  
 Attempt solution of 3-term quadratic equation M1 using formula or factorisation or equiv  
 Obtain  $\frac{1}{3}$  as only final value of  $\cos\theta$  A1 **5** or exact equiv; ignore subsequent work  
8

3 (i) Obtain or clearly imply  $60\ln x$  B1  
 Obtain  $(60\ln 20 - 60\ln 10)$  and hence  $60\ln 2$  B1 **2** with no error seen

(ii) Attempt calculation of form  $k(y_0 + 4y_1 + y_2)$  M1 any constant  $k$ ; using  $y$ -value attempts  
 Identify  $k$  as  $\frac{5}{3}$  A1  
 Obtain  $\frac{5}{3}(6 + 4 \times 4 + 3)$  and hence  $\frac{125}{3}$  or 41.7 A1 **3** or equiv

(iii) Equate answers to parts (i) and (ii) M1 provided  $\ln 2$  involved  
 Obtain  $60\ln 2 = \frac{125}{3}$  and hence  $\frac{25}{36}$  A1 **2** AG; necessary detail required including clear use of an exact value from (ii)  
7

4 (i) Attempt correct process for composition M1 numerical or algebraic  
 Obtain  $(7)$  and hence  $0$  A1 **2**

(ii) Attempt to find  $x$ -intercept M1  
 Obtain  $x \leq 7$  A1 **2** or equiv; condone use of  $<$

(iii) Attempt correct process for finding inverse M1  
 Obtain  $\pm(2-y)^3 - 1$  or  $\pm(2-x)^3 - 1$  A1  
 Obtain correct  $(2-x)^3 - 1$  A1 **3** or equiv in terms of  $x$

(iv) Refer to reflection in  $y = x$  B1 **1** or clear equiv  
8

4723

Mark Scheme

January 2010

<p>5 (i) Obtain derivative of form <math>kx(x^2 + 1)^7</math>                  Obtain <math>16x(x^2 + 1)^7</math>                  Equate first derivative to 0 and confirm <math>x = 0</math> or substitute <math>x = 0</math> and verify first derivative zero                  Refer, in some way, to <math>x^2 + 1 = 0</math> having no root</p>	<p>M1 any constant <math>k</math>                  A1 or equiv                  M1 AG; allow for deriv of form <math>kx(x^2 + 1)^7</math>                  A1 <b>4</b> or equiv</p>
-----	
<p>(ii) Attempt use of product rule                  Obtain <math>16(x^2 + 1)^7 + \dots</math>                  Obtain <math>\dots + 224x^2(x^2 + 1)^6</math>                    Substitute 0 in attempt at second derivative                  Obtain 16</p>	<p>*M1 obtaining <math>\dots + \dots</math> form                  A1<math>\surd</math> follow their <math>kx(x^2 + 1)^7</math>                  A1<math>\surd</math> follow their <math>kx(x^2 + 1)^7</math>; or unsimplified equiv                  M1 dep *M                  A1 <b>5</b> from second derivative which is correct at some point</p>
<b>9</b>	

<p>6 Integrate <math>e^{3x}</math> to obtain <math>\frac{1}{3}e^{3x}</math> or <math>e^{-\frac{1}{2}x}</math> to obtain <math>-2e^{-\frac{1}{2}x}</math>                  Obtain indefinite integral of form <math>m_1e^{3x} + m_2e^{-\frac{1}{2}x}</math>                  Obtain correct <math>\frac{1}{3}ke^{3x} - 2(k - 2)e^{-\frac{1}{2}x}</math>                    Obtain <math>e^{3\ln 4} = 64</math> or <math>e^{-\frac{1}{2}\ln 4} = \frac{1}{2}</math>                  Apply limits and equate to 185                  Obtain <math>\frac{64}{3}k - (k - 2) - \frac{1}{3}k + 2(k - 2) = 185</math>                  Obtain <math>\frac{17}{2}</math></p>	<p>B1 or both                  M1 any constants <math>m_1</math> and <math>m_2</math>                  A1 or equiv                    B1 or both                  M1 including substitution of lower limit                  A1 or equiv                  A1 <b>7</b> or equiv</p>
<b>7</b>	

<p>7 (a) <u>Either</u>: State or imply either <math>\frac{dA}{dr} = 2\pi r</math> or <math>\frac{dA}{dt} = 250</math>                  Attempt manipulation of derivatives to find <math>\frac{dr}{dt}</math>                  Obtain correct <math>\frac{250}{2\pi r}</math>                  Obtain 1.6</p> <p><u>Or</u>: Attempt to express <math>r</math> in terms of <math>t</math>                  Obtain <math>r = \sqrt{\frac{250t}{\pi}}</math>                  Differentiate <math>kt^{\frac{1}{2}}</math> to produce <math>\frac{1}{2}kt^{-\frac{1}{2}}</math>                  Substitute <math>t = 7.6</math> to obtain 1.6</p>	<p>B1 or both                    M1 using multiplication / division                  A1 or equiv                  A1 <b>4</b> or equiv; allow greater accuracy                    M1 using <math>A = 250t</math>                  A1 or equiv                  M1 any constant <math>k</math>                  A1 (<b>4</b>) allow greater accuracy</p>
--	---

4723

Mark Scheme

January 2010

- (b) State  $\frac{dm}{dt} = -150ke^{-kt}$  B1  
 Equate to  $(\pm)3$  and attempt value for  $t$  M1 using valid process; condone sign confusion  
 Obtain  $-\frac{1}{k}\ln\left(\frac{1}{50k}\right)$  or  $\frac{1}{k}\ln(50k)$  or  $\frac{\ln 50 + \ln k}{k}$  A1 3 or equiv but with correct treatment of signs  
7

- 8 (i) State scale factor is  $\sqrt{2}$  B1 allow 1.4  
 State translation is in negative  $x$ -direction ... B1 or clear equiv  
 ... by  $\frac{3}{2}$  units B1 3
- 
- (ii) Draw (more or less) correct sketch of  $y = \sqrt{2x+3}$  B1 'starting' at point on negative  $x$ -axis  
 Draw (more or less) correct sketch of  $y = \frac{N}{x^3}$  B1 showing both branches  
 Indicate one point of intersection B1 3 with both sketches correct  
 [SC: if neither sketch complete or correct but diagram correct for both in first quadrant B1]
- 
- (iii) (a) Substitute 1.9037 into  $x = N^{\frac{1}{3}}(2x+3)^{-\frac{1}{6}}$  M1 or into equation  $\sqrt{2x+3} = \frac{N}{x^3}$ ; or equiv  
 Obtain 18 or value rounding to 18 A1 2 with no error seen
- 
- (b) State or imply  $2.6282 = N^{\frac{1}{3}}(2 \times 2.6022 + 3)^{-\frac{1}{6}}$  B1  
 Attempt solution for  $N$  M1 using correct process  
 Obtain 52 A1 3 concluding with integer value  
11

- 9 (i) Identify  $\tan 55^\circ$  as  $\tan(45^\circ + 10^\circ)$  B1 or equiv  
 Use correct angle sum formula for  $\tan(A+B)$  M1 or equiv  
 Obtain  $\frac{1+p}{1-p}$  A1 3 with  $\tan 45^\circ$  replaced by 1
- 
- (ii) Either: Attempt use of identity for  $\tan 2A$  \*M1 linking  $10^\circ$  and  $5^\circ$   
 Obtain  $p = \frac{2t}{1-t^2}$  A1  
 Attempt solution for  $t$  of quadratic equation M1 dep \*M  
 Obtain  $\frac{-1 + \sqrt{1+p^2}}{p}$  A1 4 or equiv; and no second expression
- Or (1): Attempt expansion of  $\tan(60^\circ - 55^\circ)$  \*M1  
 Obtain  $\frac{\sqrt{3} - \frac{1+p}{1-p}}{1 + \sqrt{3} \frac{1+p}{1-p}}$  A1√ follow their answer from (i)  
 Attempt simplification to remove denominators M1 dep \*M  
 Obtain  $\frac{\sqrt{3}(1-p) - (1+p)}{1-p + \sqrt{3}(1+p)}$  A1 (4) or equiv

4723

Mark Scheme

January 2010

Or (2): State or imply  $\tan 15^\circ = 2 - \sqrt{3}$  B1  
 Attempt expansion of  $\tan(15^\circ - 10^\circ)$  M1 with exact attempt for  $\tan 15^\circ$   
 Obtain  $\frac{2 - \sqrt{3} - p}{1 + p(2 - \sqrt{3})}$  A2 (4)

Or (3): State or imply  $\tan 15^\circ = \frac{\sqrt{3}-1}{\sqrt{3}+1}$  B1 or exact equiv  
 Attempt expansion of  $\tan(15^\circ - 10^\circ)$  M1 with exact attempt for  $\tan 15^\circ$   
 Obtain  $\frac{\sqrt{3}-1-p\sqrt{3}-p}{\sqrt{3}+1+p\sqrt{3}-p}$  A2 (4) or equiv

Or (4): Attempt expansion of  $\tan(10^\circ - 5^\circ)$  \*M1  
 Obtain  $t = \frac{p-t}{1+pt}$  A1  
 Attempt solution for  $t$  of quadratic equation M1 dep \*M  
 Obtain  $\frac{-2 + \sqrt{4+4p^2}}{2p}$  A1 (4) or equiv; and no second expression

-----  
 (iii) Attempt expansion of both sides M1  
 Obtain  $3\sin\theta\cos 10^\circ + 3\cos\theta\sin 10^\circ =$   
 $7\cos\theta\cos 10^\circ + 7\sin\theta\sin 10^\circ$  A1 or equiv  
 Attempt division throughout by  $\cos\theta\cos 10^\circ$  M1 or by  $\cos\theta$  (or  $\cos 10^\circ$ ) only  
 Obtain  $3t + 3p = 7 + 7pt$  A1 or equiv  
 Obtain  $\frac{3p-7}{7p-3}$  A1 5 or equiv

**12**