### Stewart House 32 Russell Square London WC1B 5DN

### January 2004

### Advanced Subsidiary/Advanced Level

### General Certificate of Education

### Subject STATISTICS 6684

Ouestion number Scheme Marks 1. List of patients registered with the practice. Require 'list' or 'register' or database or similar **B1** (a) (1) **(b)** The patient(s) **B1** (1) (c) Adv: Quicker, cheaper, easier, used when testing results in destruction of item, quality of info about each sampling unit is often better. **B1** Any one Disady: Uncertainty due to natural variation, uncertainty due to bias, possible bias as sampling frame incomplete, bias due to subjective choice of sample, bias due to non-response . **B1** Any one (2) (**d**) Non-response due to patients registered with the practice but who have left the area **B1** (1) (Total 5 Marks) 2(a)  $P(R \ge 4) = 1 - P(R \le 3) = 0.6533$ Require 1 minus and correct inequality **M1A1** (2)**(b)**  $P(S \le 1) = P(S = 0) + P(S = 1) = e^{-2.71} + 2.71e^{-2.71} = 0.2469$ awrt 0.247 M1,A1,A1 (3)  $P(T \le 18) = P(Z \le \frac{18 - 25}{5}), = P(Z \le -1.4) = 0.0808$ (c) 4 dp, cc no marks M1,A1 (2)(Total 7 Marks)  $p = \frac{1}{2}$ 3(a) **B1** (1) **B1** Binomial distribution is symmetrical **(b)** (1) Since *n* is large and  $p \approx 0.5$  then use normal approximation, Can be implied below (c) **M1** np = 96 and npq = 49.92A1A1  $P(90 \le X < 105) \approx P(89.5 \le Y \le 104.5)$  where  $Y \square N(96,49.92)$  $\pm 0.5$  cc on both M1,  $\approx P\left(\frac{89.5 - 96}{\sqrt{49.92}} \le Z \le \frac{104.5 - 96}{\sqrt{49.92}}\right)$ Standardisation of both M1  $\approx P(-0.92 \le Z \le 1.20)$ awrt -0.92 & 1.20 A1 ≈ 0.7055-0.7070 4dp in range A1 (7) (Total 9 Marks)

### Stewart House 32 Russell Square London WC1B 5DN

### January 2004

### Advanced Subsidiary /Advanced Level

### General Certificate of Education

### Subject STATISTICS 6684

Ouestion number Scheme Marks 4 (a) n large, p small **B1,B1** (2) **(b)** Let *X* represent the number of people catching the virus,  $X \square B\left(12, \frac{1}{150}\right)$ Implied **B1**  $P(X = 2) = C_2^{12} \left(\frac{1}{150}\right)^2 \left(\frac{149}{150}\right)^{10}$ , = 0.0027 Use of Bin including  $C_2^{12}$ , 0.0027(4) only M1A1,A1 (4) (c)  $X \square \operatorname{Po}(np) = \operatorname{Po}(8)$ Poisson, 8 B1,B1  $P(X < 7) = P(X \le 6) = 0.3134$  $X \le 6$  for method, 0.3134 **M1A1** (4) (Total 10 Marks) 5(a) Vehicles pass at random / one at a time / independently / at a constant rate Any 2&context B1B1dep (2)**(b)** X is the number of vehicles passing in a 10 minute interval,  $X \square \operatorname{Po}\left(\frac{51}{60} \times 10\right) = \operatorname{Po}(8.5)$ Implied Po(8.5) B1  $P(X=6) = \frac{8.5^6 e^{-8.5}}{6!}$ , = 0.1066 (or 0.2562-0.1496=0.1066) Clear attempt using 6, 4dp M1A1 (3) (c)  $P(X \ge 9) = 1 - P(X \le 8) = 0.4769$ Require 1 minus and correct inequality **M1A1** (2) (**d**)  $H_0: \lambda = 8.5, H_1: \lambda < 8.5$ One tailed test only for alt hyp B1∫,B1∫  $P(X \le 4 | \lambda = 8.5) = 0.0744, > 0.05$  $X \leq 4$  for method, 0.0744 M1,A1 (Or P( $X \le 3 | \lambda = 8.5$ ) = 0.0301, < 0.05 so CR  $X \le 3$  correct CR M1,A1) Insufficient evidence to reject  $H_0$ , 'Accept' M1 so no evidence to suggest number of vehicles has decreased. A1∫ Context (6) (Total 13 Marks)

## Stewart House 32 Russell Square London WC1B 5DN

## January 2004

### Advanced Subsidiary /Advanced Level

General Certificate of Education

## Subject STATISTICS 6684

Question number	Scheme	Marks	
6 (a)	Let <i>X</i> represent the number of plant pots with defects, $X \square B(25,0.20)$ Implied $P(X \le 1) = 0.0274, P(X \ge 10) = 0.0173$ Clear attempt at both tails required, 4dp Critical region is $X \le 1, X \ge 10$	B1 M1A1A1 A1	(5)
(b)	Significance level = 0.0274+0.0173=0.0447 Accept % 4dp	<b>B1</b> cao	
( <b>c</b> )	$H_0: \lambda = 10, H_1: \lambda > 10 \text{ (or } H_0: \lambda = 60, H_1: \lambda > 60)$	B1B1	(1)
	Let <i>Y</i> represent the number sold in 6 weeks, under $H_0$ , $Y \square Po(60)$		
	$P(Y \ge 74) \approx P(W > 73.5)$ where $W \square N(60,60)$ ±0.5 for cc ,73.5		
	$\approx P(Z \ge \frac{73.5 - 60}{\sqrt{60}}) = P(Z > 1.74) =, 0.0407 - 0.0409 < 0.05$ Standardise using $60\sqrt{60}$	M1,A1	
	Evidence that rate of sales per week has increased. $A1\int$		
		Total 13 Ma	IFKS)

### Stewart House 32 Russell Square London WC1B 5DN

### January 2004

### Advanced Subsidiary /Advanced Level

### General Certificate of Education

### Subject STATISTICS 6684

Ouestion number Scheme Marks 7 (a)  $\int_{0}^{4} kx(5-x) \mathrm{d}x = 1$ Limits required **M1**  $k\left[\frac{5x^2}{2} - \frac{x^3}{3}\right]_0^4 = 1$  $\left|\frac{5x^2}{2} - \frac{x^3}{3}\right|$ A1 Sub in limits and solve to give \*\*\*\*  $k = \frac{3}{56}$  \*\*\*\* Correct solution A1 (3) **(b)**  $F(x) = \int_0^{x_0} f(x) dx = \int_0^{x_0} \frac{3}{56} x (5-x) dx = \frac{3}{56} \left[ \frac{5x^2}{2} - \frac{x^3}{3} \right]^{x_0}$  Variable upper limit required **M1**  $=\frac{x_0^2}{112}(15-2x_0)$ A1 0 x < 0 $0 \le x \le 4$  $F(x) = \frac{x^2}{112}(15 - 2x)$ Ends, middle. B1.B1.∫ (4) (c)  $E(x) = \int_{0}^{4} \frac{3}{56} x^{2} (5-x) dx = \frac{3}{56} \left[ \frac{5x^{3}}{3} - \frac{x^{4}}{4} \right]_{0}^{4} = 2.29 \int xf(x) dx, \left[ \frac{5x^{3}}{3} - \frac{x^{4}}{4} \right], 3sf(2\frac{2}{7}) M1A1A1$ (3) (**d**)  $f'(x) = \frac{3}{56}(5-2x) = 0 \implies Mode=2.5$ Attempt f'(x), (5-2x) = 0, 2.5 M1A1A1 (Or Sketch M1, x=0&5 A1, Mode=2.5 A1) (3) **(e)** F(2.3)=0.491, F(2.5)=0.558 Their F, awrt 0.491 & 0.558 or 0.984 & -6.5 M1.A1  $F(m)=0.5 \implies m$  lies between 2.3 and 2.5 cso A1 (3) (**f**) Mean (2.29)<Median (2.3-2.5)<Mode (2.5) **B1** Negative skew B1 dep (2)(Total 18 Marks)