

Mark Scheme 4723

June 2006

4723

Mark Scheme

June 2006

| | | | |
|-------|--|---|---|
| 1 | Differentiate to obtain $k(4x+1)^{-\frac{1}{2}}$ Obtain $2(4x+1)^{-\frac{1}{2}}$ Obtain $\frac{2}{3}$ for value of first derivative Attempt equation of tangent through (2, 3) Obtain $y = \frac{2}{3}x + \frac{5}{3}$ or $2x - 3y + 5 = 0$ | M1 A1 A1 M1 A1 | any non-zero constant k or equiv, perhaps unsimplified or unsimplified equiv using numerical value of first derivative provided derivative is of form $k'(4x+1)^n$ 5 or equiv involving 3 terms |
| <hr/> | | | |
| 2 | <u>Either:</u> Attempt to square both sides Obtain $3x^2 - 14x + 8 = 0$ Obtain correct values $\frac{2}{3}$ and 4 Attempt valid method for solving inequality Obtain $\frac{2}{3} < x < 4$ <u>Or:</u> Attempt solution of two linear equations or inequalities Obtain value $\frac{2}{3}$ Obtain value 4 Attempt valid method for solving inequality Obtain $\frac{2}{3} < x < 4$ | M1 A1 A1 M1 A1 M1 M1 A1 M1 M1 A1 A1 A1 M1 A1 (5) | producing 3 terms on each side or inequality involving $<$ or $>$ implied by correct answer or plausible incorrect answer 5 or correctly expressed equiv; allow \leq signs one eqn with signs of $2x$ and x the same, second eqn with signs different implied by correct answer or plausible incorrect answer implied by correct answer or plausible incorrect answer (5) or correctly expressed equiv; allow \leq signs |
| <hr/> | | | |
| 3 | (i) Attempt evaluation of cubic expression at 2 and 3 Obtain -11 and 31 Conclude by noting change of sign (ii) Obtain correct first iterate Attempt correct process to obtain at least 3 iterates Obtain 2.34 | M1 A1 A1 B1 M1 A1 | 3 or equiv; following any calculated values provided negative then positive using x_1 value such that $2 \leq x_1 \leq 3$ using any starting value now 3 answer required to 2 d.p. exactly; $2 \rightarrow 2.3811 \rightarrow 2.3354 \rightarrow 2.3410$; $2.5 \rightarrow 2.3208 \rightarrow 2.3428 \rightarrow 2.3401$; $3 \rightarrow 2.2572 \rightarrow 2.3505 \rightarrow 2.3392$ |

4723

Mark Scheme

June 2006

| | |
|--|---|
| <p>4 (i) State $\ln y = (x-1)\ln 5$</p> <p>Obtain $x = 1 + \frac{\ln y}{\ln 5}$</p> <p>(ii) Differentiate to obtain single term of form $\frac{k}{y}$</p> <p>Obtain $\frac{1}{y \ln 5}$</p> <p>(iii) Substitute for y and attempt reciprocal</p> <p>Obtain $25 \ln 5$</p> | <p>B1 whether following $\ln y = \ln 5^{x-1}$ or not; brackets needed</p> <p>B1 2 AG; correct working needed; missing brackets maybe now implied</p> <p>M1 any constant k</p> <p>A1 2 or equiv involving y</p> <p>M1 or equiv method for finding derivative without using part (ii)</p> <p>A1 2 or exact equiv</p> |
| <hr/> | |
| <p>5 (i) State $\sin 2\theta = 2 \sin \theta \cos \theta$</p> <p>(ii) Attempt to find exact value of $\cos \alpha$</p> <p>Obtain $\frac{1}{4}\sqrt{15}$</p> <p>Substitute to confirm $\frac{1}{8}\sqrt{15}$</p> <p>(iii) State or imply $\sec \beta = \frac{1}{\cos \beta}$</p> <p>Use identity to produce equation involving $\sin \beta$</p> <p>Obtain $\sin \beta = 0.3$ and hence 17.5</p> | <p>B1 1 or equiv; any letter acceptable here (and in parts (ii) and (iii))</p> <p>M1 using identity attempt or right-angled triangle</p> <p>A1 or exact equiv</p> <p>A1 3 AG</p> <p>B1</p> <p>M1</p> <p>A1 3 and no other values between 0 and 90; allow 17.4 or value rounding to 17.4 or 17.5</p> |
| <hr/> | |
| <p>6 (i) <u>Either</u>: Obtain $f(-3) = -7$</p> <p>Show correct process for compn of functions</p> <p>Obtain -47</p> <p><u>Or</u>: Show correct process for compn of functions</p> <p>Obtain $2 - (2 - x^2)^2$</p> <p>Obtain -47</p> <p>(ii) Attempt correct process for finding inverse</p> <p>Obtain either one of $x = \pm \sqrt{2-y}$ or both</p> <p>Obtain correct $-\sqrt{2-x}$</p> <p>(iii) Draw graph showing attempt at reflection in $y = x$</p> <p>Draw (more or less) correct graph</p> <p>Indicate coordinates 2 and $-\sqrt{2}$</p> | <p>B1 maybe implied</p> <p>M1</p> <p>A1 3</p> <p>M1 using algebraic approach</p> <p>A1 or equiv</p> <p>A1 (3)</p> <p>M1 as far as $x = \dots$ or equiv</p> <p>A1 or equiv perhaps involving x</p> <p>A1 3 or equiv; in terms of x now</p> <p>M1</p> <p>A1 with end-point on x-axis and no minimum point in third quadrant</p> <p>A1 3 accept -1.4 in place of $-\sqrt{2}$</p> |
| <p>7 (a) Obtain integral of form $k(4x-1)^{-1}$</p> | <p>M1 any non-zero constant k</p> |

4723

Mark Scheme

June 2006

| | | |
|---|----------------|--|
| Obtain $-\frac{1}{2}(4x-1)^{-1}$ | A1 | or equiv; allow + c |
| Substitute limits and attempt evaluation | M1 | for any expression of form $k'(4x-1)^n$ |
| Obtain $\frac{2}{21}$ | A1 4 | or exact equiv |
| (b) Integrate to obtain $\ln x$ | B1 | |
| Substitute limits to obtain $\ln 2a - \ln a$ | B1 | |
| Subtract integral attempt from attempt at area of appropriate rectangle | M1 | or equiv |
| Obtain $1 - (\ln 2a - \ln a)$ | A1 | or equiv |
| Show at least one relevant logarithm property | M1 | at any stage of solution |
| Obtain $1 - \ln 2$ and hence $\ln(\frac{1}{2}e)$ | A1 6 AG | full detail required |
| <hr/> | | |
| 8 (i) State $R = 13$ | B1 | or equiv |
| State at least one equation of form $R \cos \alpha = k$, $R \sin \alpha = k'$, $\tan \alpha = k''$ | M1 | or equiv; allow sin / cos muddles; implied by correct α |
| Obtain 67.4 | A1 3 | allow 67 or greater accuracy |
| (ii) Refer to translation and stretch | M1 | in either order; allow here equiv terms such as 'move', 'shift'; with both transformations involving constants |
| State translation in positive x direction by 67.4 | A1√ | or equiv; following their α ; using correct terminology now |
| State stretch in y direction by factor 13 | A1√ 3 | or equiv; following their R ; using correct terminology now |
| (iii) Attempt value of $\cos^{-1}(2 \div R)$ | M1 | |
| Obtain 81.15 | A1√ | following their R ; accept 81 |
| Obtain 148.5 as one solution | A1 | accept 148.5 or 148.6 or value rounding to either of these |
| Add their α value to second value correctly attempted | M1 | |
| Obtain 346.2 | A1 5 | accept 346.2 or 346.3 or value rounding to either of these; and no other solutions |

4723

Mark Scheme

June 2006

| | |
|---|--|
| <p>9 (i) Attempt to express x in terms of y Obtain $x = e^{\frac{1}{2}y} + 1$ State or imply volume involves $\int \pi x^2$ Attempt to express x^2 in terms of y Obtain $k \int (e^y + 2e^{\frac{1}{2}y} + 1) dy$ Integrate to obtain $k(e^y + 4e^{\frac{1}{2}y} + y)$ Use limits 0 and p Obtain $\pi(e^p + 4e^{\frac{1}{2}p} + p - 5)$</p> | <p>*M1 obtaining two terms A1 or equiv B1 *M1 dep *M; expanding to produce at least 3 terms A1 any constant k including 1; allow if dy absent A1 M1 dep *M *M; evidence of use of 0 needed A1 8 AG; necessary detail required</p> |
| <p>(ii) State or imply $\frac{dp}{dt} = 0.2$ Obtain $\pi(e^p + 2e^{\frac{1}{2}p} + 1)$ as derivative of V Attempt multiplication of values or expressions for $\frac{dp}{dt}$ and $\frac{dV}{dp}$ Obtain $0.2\pi(e^4 + 2e^2 + 1)$ Obtain 44</p> | <p>B1 maybe implied by use of 0.2 in product B1 M1 A1√ following their $\frac{dV}{dp}$ expression A1 5 or greater accuracy</p> |

4723

Mark Scheme

June 2006