

Question number	Scheme	Marks
<p>1. (a)</p>	<p>Take a (simple) random sample from (mutually exclusive) groups of the population 1g/1h Sample sizes within strata in strict proportion to numbers in each strata in the population Advantage: More accurate estimate of variance of population mean Individual estimates for strata available Disadvantage: Difficult if strata are large Definition of strata problematic (may overlap)</p>	<p>B1 B1 Any one B1 Any one B1</p>
<p>(b)</p>	<p>Non-random sampling from groups of the population Advantage: Representative sample can be achieved with small sample size Cheap (costs kept to a minimum) Administration relatively easy Disadvantage Not possible to estimate sampling errors due to lack of randomness Judgment of interviewer can affect choice of sample – bias OK Non-response not recorded Difficulties of defining controls e.g. social class</p>	<p>B1 B1dep Any one (not quick) B1 Any one B1</p>
<div style="border: 1px solid black; border-radius: 50%; width: 30px; height: 30px; display: flex; align-items: center; justify-content: center; margin: 0 auto;">8</div>		
<p>2. (a)</p>	<p>$X \sim N(124, 20^2)$ or $\bar{X} \sim (124, \frac{20^2}{30})$ or assume σ^2 estimated by s^2 or CLT, vals. $\bar{x} \pm 2.5758 \frac{\sigma}{\sqrt{n}} = 124 \pm 2.5758 \frac{20}{\sqrt{30}}$ $= 124 \pm 9.405$ $= (115, 133)$</p>	<p>B1, B1 B1M1A1 3 sf A1</p>
<p>(b)</p>	<p>140 is not in confidence interval Underweight apples chosen or Sample may not be representative/may be biased</p>	<p>M1 Any one A1f</p>
<div style="border: 1px solid black; border-radius: 50%; width: 30px; height: 30px; display: flex; align-items: center; justify-content: center; margin: 0 auto;">8</div>		

Question number	Scheme	Marks																															
<p>3. (a) (b) (c)</p>	<p>$E(X-Y)=20-10=10$ $Var(X-Y)=5+4=9$ $X-Y \sim N(10,9)$ $P(13 < X-Y < 16) = P(X-Y < 16) - P(X-Y < 13)$ $= P(Z < \frac{16-10}{3}) - P(Z < \frac{13-10}{3})$ $= P(Z < 2) - P(Z < 1)$ $= 0.9772 - 0.8413 = 0.1359$</p>	<p>Require minus, 10 M1A1 Require plus, 9 M1A1 Implied B1f Subtract M1 Standardise M1 2&1 A1 0.1359 A1 (5)</p>																															
<p>4.</p>	<p>H_0 : Taking drug and catching a cold are independent (not associated) H_1 : Taking drug and catching a cold are not independent (associated) (not ditto)</p> <table border="1" data-bbox="284 1003 1133 1133"> <tr> <td></td> <td>Cold</td> <td>Not Cold</td> <td></td> </tr> <tr> <td>Drug</td> <td>34 (39.5)</td> <td>66 (60.5)</td> <td>100</td> </tr> <tr> <td>Dummy</td> <td>45 (39.5)</td> <td>55 (60.5)</td> <td>100</td> </tr> <tr> <td></td> <td>79</td> <td>121</td> <td>200</td> </tr> </table> <table border="1" data-bbox="284 1155 922 1375"> <tr> <td>O</td> <td>E</td> <td>$\frac{(O - E)^2}{E}$</td> </tr> <tr> <td>34</td> <td>39.5</td> <td>0.766</td> </tr> <tr> <td>66</td> <td>60.5</td> <td>0.5</td> </tr> <tr> <td>45</td> <td>39.5</td> <td>0.765</td> </tr> <tr> <td>55</td> <td>60.5</td> <td>0.5</td> </tr> </table> <p>$\sum \frac{(O - E)^2}{E} = 2.53$ (NB with Yates 2.09) attempt & add, awrt 0.766 & 0.5 twice, awrt 2.53 M1A1A1 $\nu = 1, \chi^2_1(5\%) = 3.841 > 2.53$ 1, 3.841 B1,B1 No reason to believe that the chance of catching a cold is affected by taking the new drug A1f</p>		Cold	Not Cold		Drug	34 (39.5)	66 (60.5)	100	Dummy	45 (39.5)	55 (60.5)	100		79	121	200	O	E	$\frac{(O - E)^2}{E}$	34	39.5	0.766	66	60.5	0.5	45	39.5	0.765	55	60.5	0.5	<p>B1 both B1 All totals B1 $E = \frac{RT \times CT}{GT}$ M1A1A1 (11)</p>
	Cold	Not Cold																															
Drug	34 (39.5)	66 (60.5)	100																														
Dummy	45 (39.5)	55 (60.5)	100																														
	79	121	200																														
O	E	$\frac{(O - E)^2}{E}$																															
34	39.5	0.766																															
66	60.5	0.5																															
45	39.5	0.765																															
55	60.5	0.5																															

Question number	Scheme	Marks																																								
5	<p>μ_a and μ_b are mean weight of population after and before closure respectively.</p> <p>$H_0 : \mu_b = \mu_a$</p> <p>$H_1 : \mu_b > \mu_a$</p> $z = \frac{10 - 8}{\sqrt{\frac{2.64^2}{100} + \frac{1.94^2}{120}}}$ <p>Fraction, denom Ok alone</p> $z = \frac{2}{\sqrt{0.1011}} = 6.29$ <p>awrt 6.29</p> <p>Critical region is $z \geq 1.6449$, $6.29 > 1.6449$ or in critical region or Reject H_0 (or $P(Z \geq 6.29) = 0, 0 < 0.05$ or z in critical region or Reject H_0 B1M1)</p> <p>There is evidence that closing the factory has reduced mean river pollution</p>	<p>B1</p> <p>B1B1</p> <p>M1A1</p> <p>M1A1</p> <p>A1</p> <p>1.6449 B1, M1</p> <p>A1J</p> <p>(11)</p> <p style="text-align: center;">11</p>																																								
6 (a)	<table border="1" data-bbox="277 1003 1286 1137"> <tr><td>A</td><td>2</td><td>5</td><td>3</td><td>7</td><td>8</td><td>1</td><td>4</td><td>6</td><td></td></tr> <tr><td>B</td><td>3</td><td>2</td><td>6</td><td>5</td><td>7</td><td>4</td><td>1</td><td>8</td><td></td></tr> <tr><td> d </td><td>1</td><td>3</td><td>3</td><td>2</td><td>1</td><td>3</td><td>3</td><td>2</td><td></td></tr> <tr><td>d²</td><td>1</td><td>9</td><td>9</td><td>4</td><td>1</td><td>9</td><td>9</td><td>4</td><td>46</td></tr> </table> <p>$r_s = 1 - \frac{6 \times 46}{8 \times 63}$</p> <p>$r_s = 0.452$</p>	A	2	5	3	7	8	1	4	6		B	3	2	6	5	7	4	1	8		d	1	3	3	2	1	3	3	2		d ²	1	9	9	4	1	9	9	4	46	<p>d M1</p> <p>$\sum d^2$ M1A1</p> <p>M1A1J</p> <p>0.452 A1</p> <p>(6)</p>
A	2	5	3	7	8	1	4	6																																		
B	3	2	6	5	7	4	1	8																																		
d	1	3	3	2	1	3	3	2																																		
d ²	1	9	9	4	1	9	9	4	46																																	
(b)	<p>$H_0 : \rho = 0, H_1 : \rho \neq 0 (\rho > 0)$</p> <p>critical values are ± 0.7381 (0.6429)</p> <p>$0.452 < 0.7381$ ($0.452 < 0.6429$) or not sig or Insufficient evidence to reject H_0</p> <p>No agreement between the two judges.</p>	<p>B1B1</p> <p>0.7381(0.6429) B1</p> <p>M1</p> <p>Context A1J</p> <p>(5)</p> <p style="text-align: center;">11</p>																																								

Question number	Scheme	Marks														
7																
(a)	$\mu = 0.3 \times 50 + 0.2 \times 10 + 0.5 \times 2 = 18$ $\sigma^2 = (0.3 \times 50^2 + 0.2 \times 10^2 + 0.5 \times 2^2) - 18^2 = 448$	<p>M1A1</p> <p>M1A1</p> <p>(4)</p>														
(b)	<p>(50,50) or (50,50) without ordered pairs</p> <p>(10,2) (10,2)</p> <p>(2,10) (10,10)</p> <p>(10,10) (50,10)</p> <p>(50,10) (2,2)</p> <p>(10,50) (50,2)</p> <p>(2,2)</p> <p>(50,2)</p> <p>(2,50)</p> <p style="text-align: right;">either, -1 each missing pair</p>	<p>B2</p> <p>(2)</p>														
(c)	<table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td style="text-align: center;">\bar{x}</td> <td style="text-align: center;">2</td> <td style="text-align: center;">6</td> <td style="text-align: center;">10</td> <td style="text-align: center;">26</td> <td style="text-align: center;">30</td> <td style="text-align: center;">50</td> </tr> <tr> <td style="text-align: center;">$P(\bar{X} = \bar{x})$</td> <td style="text-align: center;">0.25</td> <td style="text-align: center;">0.2</td> <td style="text-align: center;">0.04</td> <td style="text-align: center;">0.3</td> <td style="text-align: center;">0.12</td> <td style="text-align: center;">0.09</td> </tr> </table>	\bar{x}	2	6	10	26	30	50	$P(\bar{X} = \bar{x})$	0.25	0.2	0.04	0.3	0.12	0.09	<p>All means, probabs multiplied, -1 each error</p> <p>B1 M1 A2</p> <p>(4)</p>
\bar{x}	2	6	10	26	30	50										
$P(\bar{X} = \bar{x})$	0.25	0.2	0.04	0.3	0.12	0.09										
(d)	$P(2 \leq \bar{X} < 7) = 0.25 + 0.2 = 0.45$	<p>Probabilities of 2 and 6 added, 0.45</p> <p>M1 A1J</p> <p>(2)</p>														
(e)	$E(\bar{X}) = 2 \times 0.25 + 6 \times 0.2 + \dots = 18$ $\text{Var}(\bar{X}) = 2^2 \times 0.25 + 6^2 \times 0.2 + \dots - 18^2 = 224$ $\sum x^2 P(X = x) - (\text{theirs})^2, 224$ <p>So $E(\bar{X}) = 18 = \mu$ and $\text{Var}(\bar{X}) = 224 = \frac{1}{2} \sigma^2$ as required.</p>	<p>$\sum xP(X = x)$ from table, 18</p> <p>M1 A1</p> <p>M1A1</p> <p>A1</p> <p>(5)</p>														