Version 1.0



General Certificate of Education (A-level) June 2013

Physics

Investigative Skills Assignment (ISA P)

PHY3/P13/mark

Written Test



Further copies of this Mark Scheme are available from: aqa.org.uk

Copyright $\ensuremath{\textcircled{O}}$ 2012 AQA and its licensors. All rights reserved.

Copyright

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

The Assessment and Qualifications Alliance (AQA) is a company limited by guarantee registered in England and Wales (company number 3644723) and a registered charity (registered charity number 1073334). Registered address: AQA, Devas Street, Manchester M15 6EX.

Marking Guidelines Explanatory Notes

The marking guidelines should be considered a working document. A version of the marking guidelines will be placed on the Secure Key Materials Website in September. This is to allow centre's to undertake ISA practicals as soon as they wish. Centres can use this version of the marking guidelines to mark candidates work. However this version of the marking guidelines may be subject to amendments. An updated version of the marking guidelines to be used during the present academic year will be placed on the Secure Key Materials Website by **31**st **October**. Examinations Officers must ensure that Teachers receive the final version of the marking guidelines. **Centres should ensure that their marking is in line with the updated version of the marking guidelines**.

The marking guidelines have been devised by a team of experienced examiners. They have tried to anticipate all possible responses worthy of credit. In order to establish consistency it is essential that all centres mark exactly to this scheme.

For ease of use the mark scheme has been presented in tabular form. Concise answers are given in the left-hand column. More detailed explanatory notes for some questions are included in the right-hand column.

Marking of Stage 1 of the ISA – student data and graph – should ideally be completed before the ISA written test to ensure that candidates do not change any data. (Alternatively, centres should take other steps to ensure that candidates do not change any information on their data script/graph). The marking of this section should be annotated with a red tick at the point where the mark has been awarded together with the letter referring to this mark scheme, eg ' \checkmark b'. No other comments or feedback should be written on the candidates' scripts. The total mark for this section should be written at the top of the paper. This will be transferred to the grid on the front page of the ISA test booklet.

Marking of the ISA test should be done using a red tick to represent each mark awarded. Further annotated comments **can** be added where necessary as an explanation as to why a particular point has been awarded which will greatly aid the moderation process. The total mark for each question should be entered on the grid on the front cover of the ISA booklet and the total mark calculated.

Assessment Advisers are allocated to each centre and they can advise on the marking process. You should receive the contact details for the Assessment Advisor through the post. If you have not received them, please contact the AQA subject team.

Stage 1		Mark	Additional guidance notes
(a)	Single clear table with columns correctly headed with both quantity and unit ✓	1	Headings can be in words or symbols. Units can be in words or the correct abbreviation. Standard notation for quantity and unit is expected e.g. $\frac{M}{\cos(\frac{\theta_m}{2})} / \text{kg}$ but accept units given inside brackets e.g. $\frac{M}{\cos(\frac{\theta_m}{2})}$ (g) Do not award this mark if any units are written in the body of the table
(b)	Six completed rows of data for $M = 100, 200, 300, 400, 500$ and 600 g. Each row must have four recorded values for length and two for angle θ . \checkmark	1	All values <i>M</i> should be given to the nearest gram. However allow 0.10 kg, etc but do not allow 0.1 kg, etc. All values of length must be given to the nearest mm and all values of θ to the nearest degree only.
(c)	Values of <i>L</i> and θ_m correctly calculated for $M = 200$ g and $M = 400$ g (i.e. 2 nd and 4 th readings).	1	No sf penalty
(d)	$\frac{M}{\cos\left(\frac{\theta_{\rm m}}{2}\right)}$ correctly calculated for $M = 200$ g and $M = 400$ g. \checkmark	1	No sf penalty but allow ecf from (c)
(e)	Suitably large graph scale (do not award if scale on either axis could have been doubled). Scale must have <u>sensible</u> divisions which can be easily read (e.g. not in multiples of 3, 6, 7, 9 etc.) and both scales must start at $zero$	1	The plotted points <i>plus the origin</i> should occupy at least half of each axis.

(f)	<i>L</i> plotted on the <i>y</i> -axis and $\frac{M}{\cos\left(\frac{\theta_{\rm m}}{2}\right)}$ on the <i>x</i> -axis with both axes correctly labelled with quantity and unit \checkmark	1	Allow error carried forward for incorrect unit(s) from the table, but otherwise no unit: no mark . <i>However, do not penalise here if the unit is missing both on the graph</i> <i>and in the table.</i>
(g)	First and fifth points from the <i>y</i> -axis accurately plotted to within 1 mm√	1	This mark is independent of mark (e), i.e. candidates who have used an unsuitable scale can still achieve the mark for accurate plotting.
(h)	Suitable straight line of best fit drawn√	1	A straight line with a positive gradient and a positive intercept on the L axis is expected, but credit any well drawn straight line that fits the trend and cuts the L axis if there is an even scatter of points about the line Points which are obviously anomalous should not unduly influence the line drawn.
	Total	8	

Stage 2 Section A		Mark	Additional guidance notes
1(a)(i)	Graph supports prediction because: Graph has a <u>positive</u> gradient√ Graph has a <u>positive</u> intercept on the L-axis (or y-axis) √ OR Graph does not support prediction because: Graph is not a straight line or graph does not have a positive gradient or graph does not have a positive intercept on the L-axis	2	One mark only can be awarded for the explanation if the graph does not support the theory.
1(a)(ii)	Answer is reliable because the points are close to the line of best fit ✓ OR Answer is not reliable because (some of) the points are not very close to the line of best fit	1	
1(a)(iii)	Value of B correctly found from intercept on the <i>L</i> -axis: value (probably between 4 and 6 cm) \checkmark 2 or 3 sf with unit \checkmark	2	
1(a)(iv)	Length of spring when $M = 0 \checkmark$	1	Accept: the unstretched/original/natural length of the spring
1(b)(i)	Half the range or <u>+</u> 1 mm if all values the same \checkmark	1	Allow missing \pm but a <u>correct unit</u> is required No sf penalty
1(b)(ii)	Half the range or <u>+</u> 1 degree if all values the same \checkmark	1	Allow missing <u>+</u> No sf or unit penalty
1(b)(iii)	Both percentage uncertainties shown correctly calculated, together with a correct statement as to which is the larger (or if they are the same, a statement to that effect) \checkmark	1	Allow ecf from (b)(i) and/or (b)(ii)

1(b)(iv)	Statements to the effect that: % uncertainty in <i>L</i> is smaller with the larger mass \checkmark % uncertainty in θ_m is greater with the larger mass \checkmark Consistent conclusion \checkmark	3	Consistent conclusion:Ifin part (b)(iii) $\delta L \% < \text{or} = \delta \theta_m \%$ thenthe difference would increase and $\delta \theta_m \%$ would (still) be the largerIfin part (b)(iii) $\delta L \% > \delta \theta_m \%$ thenit is likely that $\delta \theta_m \%$ would now be larger than $\delta L \%$
	Total	12	

Stage 2 Section B		Mark	Additional guidance notes
2(a)(i)	$W = 2mg\cos\phi$ $\therefore m = W/(2g\cos\phi)$	1	The question says <i>show that,</i> so the candidates must write down both steps.
2(a)(ii)	2.411 0.793 2.635 0.856	2	One mark for each column: exact answers only.
2(a)(iii)	Both points correctly plotted to the nearest mm Well drawn straight line of best fit.	2	The line should follow the trend of the points with an even scatter of points on either side of the line.
2(b)(i)	Triangle drawn with smallest side at least 8 cm in length. \checkmark Correct readings taken from the line for the triangle \checkmark Gradient in the range 0.45 to 0.49 (0.445 to 0.494) quoted to 2 or 3 significant figures \checkmark	3	The size of the triangle can be identified from readings taken from the line. The third mark is independent of the other two: error carried forward for incorrect readings (or for a poor line of best fit) which give a gradient out of range is not allowed.
2(b)(ii)	Candidate's answer for gradient in 2(b)(i) correctly multiplied by g (expected answer 4.6) \checkmark	2	No s.f. penalty. The second mark is for the unit and can be awarded if the numerical answer is incorrect.
2(c)	$\delta x\% = 0.2 \text{ and } \delta y\% = 0.5\checkmark$ $\delta(x/y)\% = \delta x\% + \delta y\% = 0.2 + 0.5 = 0.7\checkmark$ Use of $\delta(x/y)^2\% = 2 \times \delta(x/y)(\%)\checkmark$	3	Final answer is (<u>+</u>) 1.4 (%) which automatically gains all three marks Otherwise Accept only 1 s.f. for 1^{st} and/or 2^{nd} marks. The third mark is for the method, not the final answer

2(d)(i)	Systematic errors in measurements are errors which show a pattern or a bias or a trend ✓	1	 Some acceptable alternatives A systematic error is one which deviates by a fixed amount from the true value of a measurement An error which has the same value in all readings A difference between the true value of a quantity and the indicated value caused by a fault in the measuring device Accept a good example of systematic error.
2(d)(ii)	<i>y</i> would be larger \checkmark because angle θ would be smaller or because friction would be opposing the increasing weight of $m \checkmark$	2	
	Total	16	

Stage 2 Section B		Mark	Additional guidance notes
3	 (a) Weigh the box and its contents <u>each time</u> (b) (With θ = 0) place the box on the slope and lift the end slowly until the box is just about to slide down the slope and measure angle θ. (c) Measure angle θ with a protractor. or Determine angle θ using trigonometry. (d) take <u>at least three</u> readings for θ for each value of W and then average them. (e) Repeat step (b) <u>at least four</u> more times with a larger weight inside the box each time. (f) Plot a graph of θ (or tan θ) against W. (h) Award this mark for good practical technique 	5	 Please note, next to your tick, put the letter corresponding to the marking point being awarded (e.g. √a). Some acceptable answers for mark (h) With the box in the same position as before, repeat step (b) Distribute the extra weight evenly inside the box Raise the ramp with a lab-jack for stability/to measure θ accurately Clamp slope when θ found before measuring it
	Total	5	