

Centre Number						Candidate Number					
Surname						Other Names					
<b>Notice to Candidate.</b> The work you submit for assessment must be your own. If you copy from someone else or allow another candidate to copy from you, or if you cheat in any other way, you may be disqualified.											
<b>Candidate Declaration.</b> I have read and understood the Notice to Candidate and can confirm that I have produced the attached work without assistance other than that which is acceptable under the scheme of assessment.											
Candidate Signature						Date					

For Teacher's Use	
Section	Mark
PSA	
Task	
Section A	
Section B	
TOTAL (max 50)	



General Certificate of Education  
Advanced Level Examination  
June 2011

# Chemistry

# CHM6T/P11/test

## Unit 6T A2 Investigative Skills Assignment

For submission by 15 May 2011

<b>For this paper you must have:</b> <ul style="list-style-type: none"> <li>the Periodic Table/Data Sheet provided at the end of this paper</li> <li>the Task Sheet and your Candidate Results Sheet</li> <li>a ruler with millimetre measurements</li> <li>a calculator.</li> </ul>	<b>Time allowed</b> <ul style="list-style-type: none"> <li>1 hour</li> </ul>
<b>Instructions:</b> <ul style="list-style-type: none"> <li>Use black ink or black ball-point pen.</li> <li>Fill in the boxes at the top of this page.</li> <li>Answer <b>all</b> questions.</li> <li>You must answer the questions in the space provided. Do not write outside the box around each page or on blank pages.</li> <li>Do all rough work in this book. Cross through any work you do not want to be marked.</li> </ul>	<b>Information</b> <ul style="list-style-type: none"> <li>The marks for questions are shown in brackets.</li> <li>The maximum mark for this paper is 30.</li> <li>You will be marked on your ability to:               <ul style="list-style-type: none"> <li>organise information clearly</li> <li>use scientific terminology accurately.</li> </ul> </li> </ul>
<b>Details of additional assistance (if any).</b> Did the candidate receive any help or information in the production of this work? If you answer yes give the details below or on a separate page.  Yes <input type="checkbox"/> No <input type="checkbox"/>	

**Teacher Declaration:**

I confirm that the candidate's work was conducted under the conditions laid out by the specification. I have authenticated the candidate's work and am satisfied that to the best of my knowledge the work produced is solely that of the candidate.

Signature of teacher ..... Date .....

As part of AQA's commitment to assist students, AQA may make your coursework available on a strictly anonymous basis to teachers, examining staff and students in paper form or electronically, through the Internet or other means, for the purpose of indicating a typical mark or for other educational purposes. In the unlikely event that your coursework is made available for the purposes stated above, you may object to this at any time and we will remove the work on reasonable notice. If you have any concerns please contact AQA.

To see how AQA complies with the Data Protection Act 1988 please see our Privacy Statement at [aqa.org.uk](http://aqa.org.uk).


**Section A**

These questions are about the task, Determination of an equilibrium constant.  
You should use your Task Sheet and your Candidate Results Sheet to answer them.

Answer **all** questions in the spaces provided.

- 1** Record the average titre from your Candidate Results Sheet.
- .....
- (1 mark)
- 2** The concentration of sodium hydroxide solution was  $0.100 \text{ mol dm}^{-3}$ .
- Use your answer from Question **1** to calculate the amount, in moles, of sodium hydroxide required to neutralise all of the acid in  $25.0 \text{ cm}^3$  of the diluted equilibrium mixture.
- .....
- .....
- (1 mark)
- 3** Use your answer to Question **2** to calculate the total amount, in moles, of  $\text{H}^+$  ions in the whole sample of undiluted equilibrium mixture ( $250 \text{ cm}^3$  of diluted equilibrium mixture).
- .....
- .....
- (1 mark)
- 4** The mixture you prepared and allowed to reach equilibrium contained  $2.00 \text{ cm}^3$  of  $1.00 \text{ mol dm}^{-3}$  sulfuric acid as a catalyst.
- 4 (a)** Calculate the amount, in moles, of  $\text{H}^+$  ions in the sulfuric acid. Assume that the sulfuric acid is completely dissociated.
- .....
- .....
- .....
- (1 mark)
- 4 (b)** Use your answers to Question **3** and Question **4 (a)** to calculate the amount, in moles, of propanoic acid in the whole sample of undiluted equilibrium mixture.
- .....
- (1 mark)

5 The equation for the reaction occurring in the mixture is given below.



5 (a) The original 7.00 cm<sup>3</sup> sample of propanoic acid used to prepare the equilibrium mixture contained 0.0940 mol of CH<sub>3</sub>CH<sub>2</sub>COOH

Use your answer to Question 4 (b), to calculate the amount, in moles, of propanoic acid that has reacted at equilibrium.

(If you were unable to complete the calculation in Question 4 (b) assume that the amount of propanoic acid in the whole sample of undiluted equilibrium mixture was 0.0258 mol. This is **not** the correct answer.)

.....  
..... (1 mark)

5 (b) The original 7.00 cm<sup>3</sup> sample of ethanol used to prepare the equilibrium mixture contained 0.102 mol of CH<sub>3</sub>CH<sub>2</sub>OH

Use this information and your answer to Question 5 (a) to calculate the amount, in moles, of ethanol remaining at equilibrium.

.....  
..... (1 mark)

5 (c) Use your answer to Question 5 (a) to deduce the amount, in moles, of CH<sub>3</sub>CH<sub>2</sub>COOCH<sub>2</sub>CH<sub>3</sub> and the amount, in moles, of H<sub>2</sub>O formed at equilibrium.

CH<sub>3</sub>CH<sub>2</sub>COOCH<sub>2</sub>CH<sub>3</sub> .....  
.....  
.....  
H<sub>2</sub>O .....  
.....  
..... (2 marks)

6 Suggest the main reason why the actual amount of water in the equilibrium mixture is greater than that deduced in Question 5 (c).

.....  
..... (1 mark)

Turn over ►

7 In another experiment, at a fixed temperature, a different equilibrium mixture contained the following amounts, in moles, of each component.

$\text{CH}_3\text{CH}_2\text{COOH}$	$\text{CH}_3\text{CH}_2\text{OH}$	$\text{CH}_3\text{CH}_2\text{COOCH}_2\text{CH}_3$	$\text{H}_2\text{O}$
0.0424	0.0525	0.0745	0.0813

7 (a) Write an expression for the equilibrium constant,  $K_c$ , for the reaction given in Question 5.

.....  
 .....  
 .....

(1 mark)

7 (b) Use the data in the table above to calculate a value for the equilibrium constant,  $K_c$ , at this fixed temperature.  
 Record your answer to the appropriate precision.

.....  
 .....  
 .....

(2 marks)

8 If the mixture is uncovered during the time it is left to reach equilibrium, some of the ester formed will evaporate.  
 Explain why a smaller volume of sodium hydroxide would then be required in the titration compared with the volume for the covered mixture.

.....  
 .....  
 .....

(2 marks)

**Section B**

Answer **all** questions in the spaces provided.

**9** Propanoic acid can be made from propan-1-ol by oxidation using acidified potassium dichromate(VI). Propanal is formed as an intermediate during this oxidation.

**9 (a)** State the colour of the chromium species after the potassium dichromate(VI) has reacted.

.....  
(1 mark)

**9 (b)** Describe the experimental conditions and the practical method used to ensure that the acid is obtained in a high yield. Draw a diagram of the assembled apparatus you would use.

Conditions .....

.....

Apparatus

(4 marks)

**9 (c)** Describe the different experimental conditions necessary to produce propanal in high yield rather than propanoic acid.

.....  
.....  
(2 marks)

Turn over ►

**10** Propan-1-ol is a volatile, flammable liquid.  
Give **one** safety precaution that should be used during the reaction to minimise this hazard.

.....

(1 mark)

**11** A student followed the progress of the oxidation of propan-1-ol to propanoic acid by extracting the organic compounds from one sample of reaction mixture.

**11 (a)** Give a chemical reagent which would enable the student to confirm the presence of propanal in the extracted compounds.  
State what you would observe when propanal reacts with this reagent.

Reagent .....

Observation .....

.....

(2 marks)

**11 (b)** Give a chemical reagent that would enable the student to confirm the presence of propanoic acid in the extracted compounds.  
State what you would observe when propanoic acid reacts with this reagent.

Reagent .....

Observation .....

.....

(2 marks)

**12** Predict which **one** of the compounds, propan-1-ol, propanal and propanoic acid will have the highest boiling point. Explain your answer.

Prediction .....

Explanation .....

.....

.....

.....

.....

(3 marks)

**END OF QUESTIONS**

## GCE Chemistry Data Sheet

Table 1

Infrared absorption data

Bond	Wavenumber /cm <sup>-1</sup>
N-H (amines)	3300 – 3500
O-H (alcohols)	3230 – 3550
C-H	2850 – 3300
O-H (acids)	2500 – 3000
C≡N	2220 – 2260
C=O	1680 – 1750
C=C	1620 – 1680
C-O	1000 – 1300
C-C	750 – 1100

Table 2

<sup>1</sup>H n.m.r. chemical shift data

Type of proton	δ/ppm
ROH	0.5–5.0
RCH <sub>3</sub>	0.7–1.2
RNH <sub>2</sub>	1.0–4.5
R <sub>2</sub> CH <sub>2</sub>	1.2–1.4
R <sub>3</sub> CH	1.4–1.6
	2.1–2.6
	3.1–3.9
RCH <sub>2</sub> Cl or Br	3.1–4.2
	3.7–4.1
	4.5–6.0
	9.0–10.0
	10.0–12.0

Table 3

<sup>13</sup>C n.m.r. chemical shift data

Type of carbon	δ/ppm
	5–40
	10–70
	20–50
	25–60
	alcohols, ethers or esters 50–90
	90–150
	110–125
	110–160
	esters or acids 160–185
	aldehydes or ketones 190–220

Turn over ►

# The Periodic Table of the Elements

	1	2	3	4	5	6	7	0
(1)	6.9 <b>Li</b> lithium 3	9.0 <b>Be</b> beryllium 4	10.8 <b>B</b> boron 5	12.0 <b>C</b> carbon 6	14.0 <b>N</b> nitrogen 7	16.0 <b>O</b> oxygen 8	19.0 <b>F</b> fluorine 9	20.2 <b>Ne</b> neon 10
(2)	23.0 <b>Na</b> sodium 11	24.3 <b>Mg</b> magnesium 12	27.0 <b>Al</b> aluminium 13	28.1 <b>Si</b> silicon 14	31.0 <b>P</b> phosphorus 15	32.1 <b>S</b> sulfur 16	35.5 <b>Cl</b> chlorine 17	39.9 <b>Ar</b> argon 18
(3)	39.1 <b>K</b> potassium 19	40.1 <b>Ca</b> calcium 20	45.0 <b>Sc</b> scandium 21	47.9 <b>Ti</b> titanium 22	50.9 <b>V</b> vanadium 23	52.0 <b>Cr</b> chromium 24	58.9 <b>Co</b> cobalt 27	79.9 <b>Br</b> bromine 35
(4)	85.5 <b>Rb</b> rubidium 37	87.6 <b>Sr</b> strontium 38	88.9 <b>Y</b> yttrium 39	91.2 <b>Zr</b> zirconium 40	92.9 <b>Nb</b> niobium 41	96.0 <b>Mo</b> molybdenum 42	102.9 <b>Rh</b> rhodium 45	126.9 <b>I</b> iodine 53
(5)	132.9 <b>Cs</b> caesium 55	137.3 <b>Ba</b> barium 56	138.9 <b>La *</b> lanthanum 57	178.5 <b>Hf</b> hafnium 72	180.9 <b>Ta</b> tantalum 73	183.8 <b>W</b> tungsten 74	192.2 <b>Ir</b> iridium 77	209.0 <b>Po</b> polonium 84
(6)	[223] <b>Fr</b> francium 87	[226] <b>Ra</b> radium 88	[227] <b>Ac †</b> actinium 89	[267] <b>Rf</b> rutherfordium 104	[268] <b>Db</b> dubnium 105	[271] <b>Sg</b> seaborgium 106	[276] <b>Mt</b> meitnerium 109	[222] <b>Rn</b> radon 86
(7)								
(8)	1.0 <b>H</b> hydrogen 1							
(9)								
(10)								
(11)								
(12)								
(13)								
(14)								
(15)								
(16)								
(17)								
(18)								

Key	
relative atomic mass	
<b>symbol</b>	
name	
atomic (proton) number	

Elements with atomic numbers 112-116 have been reported but not fully authenticated	
204.4 <b>Tl</b> thallium 81	207.2 <b>Pb</b> lead 82
200.6 <b>Hg</b> mercury 80	209.0 <b>Bi</b> bismuth 83
197.0 <b>Au</b> gold 79	209.0 <b>Po</b> polonium 84
195.1 <b>Pt</b> platinum 78	209.0 <b>Po</b> polonium 84
192.2 <b>Ir</b> iridium 77	209.0 <b>Po</b> polonium 84
190.2 <b>Os</b> osmium 76	209.0 <b>Po</b> polonium 84
186.2 <b>Re</b> rhenium 75	209.0 <b>Po</b> polonium 84
183.8 <b>W</b> tungsten 74	209.0 <b>Po</b> polonium 84
180.9 <b>Ta</b> tantalum 73	209.0 <b>Po</b> polonium 84
178.5 <b>Hf</b> hafnium 72	209.0 <b>Po</b> polonium 84
174.1 <b>Rf</b> rutherfordium 104	209.0 <b>Po</b> polonium 84
173.1 <b>Yb</b> ytterbium 70	209.0 <b>Po</b> polonium 84
171.9 <b>Lu</b> lutetium 71	209.0 <b>Po</b> polonium 84
170.9 <b>Tm</b> thulium 69	209.0 <b>Po</b> polonium 84
168.9 <b>Er</b> erbium 68	209.0 <b>Po</b> polonium 84
167.3 <b>Dy</b> dysprosium 66	209.0 <b>Po</b> polonium 84
164.9 <b>Ho</b> holmium 67	209.0 <b>Po</b> polonium 84
162.5 <b>Gd</b> gadolinium 64	209.0 <b>Po</b> polonium 84
158.9 <b>Tb</b> terbium 65	209.0 <b>Po</b> polonium 84
157.3 <b>Gd</b> gadolinium 64	209.0 <b>Po</b> polonium 84
152.0 <b>Eu</b> europium 63	209.0 <b>Po</b> polonium 84
150.4 <b>Sm</b> samarium 62	209.0 <b>Po</b> polonium 84
145 <b>Pm</b> promethium 61	209.0 <b>Po</b> polonium 84
144.2 <b>Nd</b> neodymium 60	209.0 <b>Po</b> polonium 84
140.9 <b>Pr</b> praseodymium 59	209.0 <b>Po</b> polonium 84
140.1 <b>Ce</b> cerium 58	209.0 <b>Po</b> polonium 84
137.3 <b>Ba</b> barium 56	209.0 <b>Po</b> polonium 84
132.9 <b>Cs</b> caesium 55	209.0 <b>Po</b> polonium 84
131.3 <b>Xe</b> xenon 54	209.0 <b>Po</b> polonium 84
126.9 <b>I</b> iodine 53	209.0 <b>Po</b> polonium 84
127.6 <b>Te</b> tellurium 52	209.0 <b>Po</b> polonium 84
121.8 <b>Sb</b> antimony 51	209.0 <b>Po</b> polonium 84
118.7 <b>Sn</b> tin 50	209.0 <b>Po</b> polonium 84
114.8 <b>In</b> indium 49	209.0 <b>Po</b> polonium 84
112.4 <b>Cd</b> cadmium 48	209.0 <b>Po</b> polonium 84
107.9 <b>Ag</b> silver 47	209.0 <b>Po</b> polonium 84
106.4 <b>Pd</b> palladium 46	209.0 <b>Po</b> polonium 84
101.1 <b>Ru</b> ruthenium 44	209.0 <b>Po</b> polonium 84
101.1 <b>Ru</b> ruthenium 44	209.0 <b>Po</b> polonium 84
55.8 <b>Fe</b> iron 26	209.0 <b>Po</b> polonium 84
58.9 <b>Co</b> cobalt 27	209.0 <b>Po</b> polonium 84
58.9 <b>Co</b> cobalt 27	209.0 <b>Po</b> polonium 84
54.9 <b>Mn</b> manganese 25	209.0 <b>Po</b> polonium 84
54.9 <b>Mn</b> manganese 25	209.0 <b>Po</b> polonium 84
52.0 <b>Cr</b> chromium 24	209.0 <b>Po</b> polonium 84
52.0 <b>Cr</b> chromium 24	209.0 <b>Po</b> polonium 84
50.9 <b>V</b> vanadium 23	209.0 <b>Po</b> polonium 84
50.9 <b>V</b> vanadium 23	209.0 <b>Po</b> polonium 84
47.9 <b>Ti</b> titanium 22	209.0 <b>Po</b> polonium 84
47.9 <b>Ti</b> titanium 22	209.0 <b>Po</b> polonium 84
45.0 <b>Sc</b> scandium 21	209.0 <b>Po</b> polonium 84
45.0 <b>Sc</b> scandium 21	209.0 <b>Po</b> polonium 84
40.1 <b>Ca</b> calcium 20	209.0 <b>Po</b> polonium 84
40.1 <b>Ca</b> calcium 20	209.0 <b>Po</b> polonium 84
39.1 <b>K</b> potassium 19	209.0 <b>Po</b> polonium 84
39.1 <b>K</b> potassium 19	209.0 <b>Po</b> polonium 84
38.9 <b>Y</b> yttrium 39	209.0 <b>Po</b> polonium 84
38.9 <b>Y</b> yttrium 39	209.0 <b>Po</b> polonium 84
37 <b>Rb</b> rubidium 37	209.0 <b>Po</b> polonium 84
37 <b>Rb</b> rubidium 37	209.0 <b>Po</b> polonium 84
36 <b>Kr</b> krypton 36	209.0 <b>Po</b> polonium 84
36 <b>Kr</b> krypton 36	209.0 <b>Po</b> polonium 84
35.5 <b>Cl</b> chlorine 17	209.0 <b>Po</b> polonium 84
35.5 <b>Cl</b> chlorine 17	209.0 <b>Po</b> polonium 84
32.1 <b>S</b> sulfur 16	209.0 <b>Po</b> polonium 84
32.1 <b>S</b> sulfur 16	209.0 <b>Po</b> polonium 84
31.0 <b>P</b> phosphorus 15	209.0 <b>Po</b> polonium 84
31.0 <b>P</b> phosphorus 15	209.0 <b>Po</b> polonium 84
28.1 <b>Si</b> silicon 14	209.0 <b>Po</b> polonium 84
28.1 <b>Si</b> silicon 14	209.0 <b>Po</b> polonium 84
27.0 <b>Al</b> aluminium 13	209.0 <b>Po</b> polonium 84
27.0 <b>Al</b> aluminium 13	209.0 <b>Po</b> polonium 84
20.2 <b>Ne</b> neon 10	209.0 <b>Po</b> polonium 84
20.2 <b>Ne</b> neon 10	209.0 <b>Po</b> polonium 84
19.0 <b>F</b> fluorine 9	209.0 <b>Po</b> polonium 84
19.0 <b>F</b> fluorine 9	209.0 <b>Po</b> polonium 84
16.0 <b>O</b> oxygen 8	209.0 <b>Po</b> polonium 84
16.0 <b>O</b> oxygen 8	209.0 <b>Po</b> polonium 84
14.0 <b>N</b> nitrogen 7	209.0 <b>Po</b> polonium 84
14.0 <b>N</b> nitrogen 7	209.0 <b>Po</b> polonium 84
12.0 <b>C</b> carbon 6	209.0 <b>Po</b> polonium 84
12.0 <b>C</b> carbon 6	209.0 <b>Po</b> polonium 84
10.8 <b>B</b> boron 5	209.0 <b>Po</b> polonium 84
10.8 <b>B</b> boron 5	209.0 <b>Po</b> polonium 84
9.0 <b>Be</b> beryllium 4	209.0 <b>Po</b> polonium 84
9.0 <b>Be</b> beryllium 4	209.0 <b>Po</b> polonium 84
6.9 <b>Li</b> lithium 3	209.0 <b>Po</b> polonium 84
6.9 <b>Li</b> lithium 3	209.0 <b>Po</b> polonium 84

* 58 – 71 Lanthanides	† 90 – 103 Actinides
173.1 <b>Yb</b> ytterbium 70	175.0 <b>Lu</b> lutetium 71
173.1 <b>Yb</b> ytterbium 70	[262] <b>Lr</b> lawrencium 103
168.9 <b>Tm</b> thulium 69	[258] <b>Md</b> mendelevium 101
168.9 <b>Tm</b> thulium 69	[258] <b>Md</b> mendelevium 101
167.3 <b>Er</b> erbium 68	[257] <b>Fm</b> fermium 100
167.3 <b>Er</b> erbium 68	[257] <b>Fm</b> fermium 100
164.9 <b>Ho</b> holmium 67	[252] <b>Es</b> einsteinium 99
164.9 <b>Ho</b> holmium 67	[252] <b>Es</b> einsteinium 99
162.5 <b>Dy</b> dysprosium 66	[251] <b>Cf</b> californium 98
162.5 <b>Dy</b> dysprosium 66	[251] <b>Cf</b> californium 98
158.9 <b>Tb</b> terbium 65	[247] <b>Bk</b> berkelium 97
158.9 <b>Tb</b> terbium 65	[247] <b>Bk</b> berkelium 97
157.3 <b>Gd</b> gadolinium 64	[247] <b>Cm</b> curium 96
157.3 <b>Gd</b> gadolinium 64	[247] <b>Cm</b> curium 96
152.0 <b>Eu</b> europium 63	[243] <b>Am</b> americium 95
152.0 <b>Eu</b> europium 63	[243] <b>Am</b> americium 95
150.4 <b>Sm</b> samarium 62	[244] <b>Pu</b> plutonium 94
150.4 <b>Sm</b> samarium 62	[244] <b>Pu</b> plutonium 94
[145] <b>Pm</b> promethium 61	[237] <b>Np</b> neptunium 93
[145] <b>Pm</b> promethium 61	[237] <b>Np</b> neptunium 93
144.2 <b>Nd</b> neodymium 60	238.0 <b>U</b> uranium 92
144.2 <b>Nd</b> neodymium 60	238.0 <b>U</b> uranium 92
140.9 <b>Pr</b> praseodymium 59	231.0 <b>Pa</b> protactinium 91
140.9 <b>Pr</b> praseodymium 59	231.0 <b>Pa</b> protactinium 91
140.1 <b>Ce</b> cerium 58	232.0 <b>Th</b> thorium 90
140.1 <b>Ce</b> cerium 58	232.0 <b>Th</b> thorium 90