


PH1

Question		Marking details	Marks Available																			
1	(a)	Rate of change [accept: increase] of velocity [not speed]. accept: $\frac{v-u}{t}$ or $\frac{\Delta v}{t}$ or $\frac{\Delta v}{\Delta t}$ (not $\frac{v}{t}$)	1																			
	(b)	(i) Both ΣF calculated correctly (20 N and 4 N) (1) Use of $a = \frac{\Sigma F}{m}$ (1) Accelerations = 10 m s ⁻² and 2 m s ⁻² (e.c.f.) (1) [Accept answers based upon calculating resultant acceleration]	3																			
		(ii) Diagram with forces shown in opposition (1) and horizontal (1) [B.o.d. on plan-view forces unless clearly incorrect]	2																			
	(c)	<table border="1"> <thead> <tr> <th>Statement</th> <th>Must be true</th> <th>Could be true</th> <th>Cannot be true</th> </tr> </thead> <tbody> <tr> <td>XXXXXXXXXX</td> <td></td> <td></td> <td>✓</td> </tr> <tr> <td>XXXXXXXXXX</td> <td></td> <td>✓</td> <td></td> </tr> <tr> <td>XXXXXXXXXX</td> <td></td> <td>✓</td> <td></td> </tr> <tr> <td>XXXXXXXXXX</td> <td></td> <td></td> <td>✓</td> </tr> </tbody> </table>	Statement	Must be true	Could be true	Cannot be true	XXXXXXXXXX			✓	XXXXXXXXXX		✓		XXXXXXXXXX		✓		XXXXXXXXXX			✓
Statement	Must be true	Could be true	Cannot be true																			
XXXXXXXXXX			✓																			
XXXXXXXXXX		✓																				
XXXXXXXXXX		✓																				
XXXXXXXXXX			✓																			
			[10]																			
2.	(a)	Same p.d. in both branches (1) 2 × resistance in upper branch (or converse) (1) } Correct qualitative ½ current in upper branch (or converse) (1) } answer → 1 mark Currents add up to 0.12 A (1)	4																			
	(b)	(i) 4.8 V (ii) 9.6 V	1 1																			
	(c)	(i) Correct use of $R = \frac{V}{I}$ (1) [or by implication] = 120 Ω (1) [e.g. $\frac{4.8}{0.04}$ ✓, $\frac{9.6}{0.08}$ (✓ b.o.d.) or $\frac{9.6}{0.4}$ ✓ not $\frac{9.6}{1.2}$ ×]	2																			
		(ii) $\frac{1}{R} = \frac{1}{R_A} + \frac{1}{R_B}$ [or equiv.] and use of correct values of R_A and R_B . [or by impl.](1) $R = 80 \Omega$ (1) or $R = \frac{9.6}{0.12}$ (1) = 80 Ω (1)	2																			
			[10]																			

Question		Marking details	Marks Available
3	(a)	(i) 2.4 V (ii) $\frac{V}{V_{\text{TOTAL}}} = \frac{R}{R_{\text{TOTAL}}}$ or $V_{\text{OUT}} = \frac{R_1}{R_1 + R_2} V_{\text{IN}}$, selected [or by impl.] Substitution, e.g. $3.6 = \frac{225}{225 + R_2} 6.0$ (1) [or by impl.] Manipulation (1); $R_2 = 150 \Omega$ (1) Or – from 1 st principles: $\frac{3.6}{225}$ (1) = 0.016 A (1); $\frac{2.4}{0.016}$ (1) = 150Ω (1) [or $\frac{6.0}{0.016} = 375 \Omega$ then $375 - 225 = 150 \Omega$]	1 4
	(b)	(i) R_2 changed to 1500Ω (ii) Initial current = $\frac{6.0}{375 \text{ e.c.f.}}$ (1) = 0.016 A (1) Final current = 0.0016 A (1) [or accept answer based upon a good qualitative argument] N.B. Calculation of final current → only 2 unless current previously calculated]	1 3 [9]
4.	(a)	Collisions between electrons and lattice [or atoms of the conductor or [metal] ions] (1) Kinetic energy of electrons transferred to lattice (1) <u>No collisions</u> (and ∴ no energy transfer) in superconductors) (1) [accept: interactions instead of collisions]	3
	(b)	(i) -190°C (ii) A→B [accept $-250 - -190$, or “temperatures below -190 ”] [full region required]	1 1
	(c)	liquid nitrogen [accept: liquid helium]	1 [6]

Question		Marking details	Marks Available
5.	(a)	Flow of charge	1
	(b)	<div style="display: flex; align-items: center;"> <div style="margin-right: 20px;">(i)</div>  <div style="margin-left: 20px;"> <p>n [free] electrons per unit volume [or electron density] [or in written answer]</p> </div> </div> <p>Diagram (1) $\left. \begin{array}{l} \text{Volume of conductor} = Al \\ \text{Number of free electrons} = nAl \end{array} \right\} \text{ (1)}$ Total charge flowing within $l = nAle$ [Accept without [] if diagram is clear]</p> <p>$I = \frac{nAle}{t} \text{ (1) and } v = \frac{l}{t} \text{ (1)}$</p> <p>(ii) $v = \frac{I}{nAe}$ $= \frac{3.0}{5.0 \times 10^{28} \times 2 \times 10^{-6} \text{ (1)} \times 1.6 \times 10^{-19}} = 1.9 \times 10^{-4} \text{ m s}^{-1} \text{ ((unit))(1)}$</p> <p>(iii) I.the same as II.half.....</p>	4 2 1 1 [9]
6	(a)	Units of LHS = N = kg m s ⁻² Units of RHS = (kg m ⁻³ · m ²) (+ manip.)(1) × (m ² s ⁻²) (1)	3
		$v^2 = \frac{2.8 \times 10^4}{1.2 \times 15 \times 4.2} \text{ (1) [or by impl.]}$ $v = 19.2 \text{ m s}^{-1} \text{ (1)}$	2
	(b)	(i) Centre of gravity	1
		(ii) Bottom of near-side wheel labelled as ‘pivot’	1
		$F_{\text{wind}} \times 2.1 \text{ (1)} = 1.0 \times 10^5 \times 1.4 \text{ (1) [or by impl.]}$ $\therefore F_{\text{wind}} = 67 \text{ kN (1) [accept 66 kN } \checkmark \text{ b.o.d.]}$	3
			[10]

Question		Marking details	Marks Available
7.	(a)	(i) Ball is seen to stay directly in front of the passenger [or clearly implied by 2 nd statement]. (1) No [horizontal] forces on ball [so horizontal speed is constant, with the same value as the train] (1)	2
		(ii) Observer sees the ball moving in the same direction as the train [with the same speed]. [Accept: "moving with the train."]	1
	(b)	Passenger sees the ball accelerating [or moving] 'backwards' [or towards the rear of the train]. Observer sees the ball moving in the same direction as the train with decreasing speed. (1) Net [horizontal] force on ball [due to air resistance] towards the back of the train. (1)	3
		(i) The graph is symmetrical / up time = down time.	1
		(ii) $x = 11 \text{ m}; t = 1.5 \text{ s}$ (1) $x = \frac{u+v}{2}t$, or $v^2 = u^2 + 2ax$ or $x = ut + \frac{1}{2}at^2$, or $11 = \frac{u}{2} \times 1.5$ (1) $0 = u^2 + 2 \times 9.81 \times 11$ $11 = 1.5u + \frac{1}{2} \times 9.81 \times 1.5^2$ (1) (1) $\therefore u = 14.7 \text{ m s}^{-1}$ (1) [accept $v = u + at$ with $v = 0$ and $t = 1.5 \text{ s}$]	3
		(iii) Graph: v axis – 20 to + 20 e.c.f. (1) Intercept on v axis 14.7 m s^{-1} e.c.f. (1) Straight line graph (1) to intercept time axis of 1.5 s (1) Graph continued straight beyond 1.5 s to negative values of v (1)	5
	[15]		

Question		Marking details	Marks Available
8	(a)	Energy cannot be created or destroyed (1) ...[only] changed from one form into another (1) [Accept: total energy [in the Universe] is constant for 1 st mark]	2
	(b)	(i) Area under graph = energy stored (1) [or by impl.] $x = 70$ m chosen (1) [or by impl.] Elastic potential energy [= $\frac{1}{2} \times 1600 \times 70$] = 56 kJ (1) Alternative: calculation of k [22.9 N m ⁻¹] or left as, e.g. $\frac{1600}{70}$ ✓ Use of $x = 70$ m to calculate energy ✓; Energy stored = 56 kJ ✓	3 3
		(ii) 70 kJ – 56 kJ (e.c.f.) (1) = mgh (1) [Or $E_p(\text{grav})$ lost = $60 \times 9.81 \times 96$ (1) [= 56 kJ]. $\therefore 14$ kJ = mgh (1)] $\therefore h = 23.8$ m (1) Alternative: 7.0×10^4 J = mgh ✓ $\rightarrow h = 118.92$ m ✓ Then subtract 96 m $\rightarrow 22.92$ m ✓	3
		(iii) Tension in 'bungee' = weight of Jumper = 60×9.81 [= 589 N] (1) From graph $x = 26$ [± 1] m (1) [or from $k \rightarrow 25.8$ m] $\therefore d = 52$ m. (1)	3 [11]