



ADVANCED GCE
MATHEMATICS
Probability & Statistics 3

4734

Candidates answer on the answer booklet.

OCR supplied materials:

- 8 page answer booklet (sent with general stationery)
- List of Formulae (MF1)

Other materials required:

- Scientific or graphical calculator

Thursday 23 June 2011
Morning

Duration: 1 hour 30 minutes



INSTRUCTIONS TO CANDIDATES

- Write your name, centre number and candidate number in the spaces provided on the answer booklet. Please write clearly and in capital letters.
- Use black ink. Pencil may be used for graphs and diagrams only.
- Read each question carefully. Make sure you know what you have to do before starting your answer.
- Answer **all** the questions.
- Do **not** write in the bar codes.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- You are permitted to use a scientific or graphical calculator in this paper.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- **You are reminded of the need for clear presentation in your answers.**
- The total number of marks for this paper is **72**.
- This document consists of **4** pages. Any blank pages are indicated.

2

- 1 The random variables X and Y are independent with $X \sim \text{Po}(5)$ and $Y \sim \text{Po}(4)$. S denotes the sum of 2 observations of X and 3 observations of Y .

(i) Find $E(S)$ and $\text{Var}(S)$. [2]

(ii) The random variable T is defined by $\frac{1}{2}X - \frac{1}{4}Y$. Show that $E(T) = \text{Var}(T)$. [3]

(iii) State which of S and T (if either) does not have a Poisson distribution, giving a reason for your answer. [2]

- 2 The population proportion of all men with red-green colour blindness is denoted by p . Each of a random sample of 80 men was tested and it was found that 6 had red-green colour blindness.

(i) Calculate an approximate 95% confidence interval for p . [4]

(ii) For a different random sample of men, the proportion with red-green colour blindness is denoted by p_s . Estimate the sample size required in order that $|p_s - p| \leq 0.05$ with probability 95%. [3]

(iii) Give one reason why the calculated sample size is an estimate. [1]

- 3 The monthly demand for a product, X thousand units, is modelled by the random variable X with probability density function given by

$$f(x) = \begin{cases} ax & 0 \leq x \leq 1, \\ a(x-2)^2 & 1 < x \leq 2, \\ 0 & \text{otherwise,} \end{cases}$$

where a is a positive constant. Find

(i) the value of a , [4]

(ii) the probability that the monthly demand is at most 1500 units, [2]

(iii) the expected monthly demand. [3]

- 4 An experiment by Lord Rutherford at Cambridge in 1909 involved measuring the numbers of α -particles emitted during radioactive decay. The following table shows emissions during 2608 intervals of 7.5 seconds.

Number of particles emitted, x	0	1	2	3	4	5	6	7	8	9	10	≥ 11
Frequency	57	203	383	525	532	408	273	139	45	27	10	6

It is given that the mean number of particles emitted per interval, calculated from the data, is 3.87, correct to 3 significant figures.

(i) Find the contribution to the χ^2 value of the frequency of 273 corresponding to $x = 6$ in a goodness of fit test for a Poisson distribution. [4]

(ii) Given that no cells need to be combined, state why the number of degrees of freedom is 10. [1]

(iii) Given also that the calculated value of χ^2 is 13.0, correct to 3 significant figures, carry out the test at the 10% significance level. [4]

3

- 5 The continuous random variable X has (cumulative) distribution function given by

$$F(x) = \begin{cases} 0 & x < 1, \\ \frac{4}{3} \left(1 - \frac{1}{x^2}\right) & 1 \leq x \leq 2, \\ 1 & x > 2. \end{cases}$$

- (i) Find the median value of X . [2]

- (ii) Find the (cumulative) distribution function of Y , where $Y = \frac{1}{X^2}$, and hence find the probability density function of Y . [6]

- (iii) Evaluate $E\left(2 - \frac{2}{X^2}\right)$. [3]

- 6 The Body Mass Index (BMI) of each of a random sample of 100 army recruits from a large intake in 2008 was measured. The results are summarised by

$$\Sigma x = 2605.0, \quad \Sigma x^2 = 68\,636.41.$$

It may be assumed that BMI has a normal distribution.

- (i) Find a 98% confidence interval for the mean BMI of all recruits in 2008. [5]

- (ii) Estimate the percentage of the intake with a BMI greater than 30.0. [3]

- (iii) The BMIs of two randomly chosen recruits are denoted by B_1 and B_2 . Estimate $P(B_1 - B_2 < 5)$. [4]

- (iv) State, giving a reason, for which of the above calculations the normality assumption is unnecessary. [1]

- 7 In order to improve their mathematics results 10 students attended an intensive Summer School course. Each student took a test at the start of the course and a similar test at the end of the course. The table shows the scores achieved in each test.

Student	1	2	3	4	5	6	7	8	9	10
First test score	37	27	38	47	54	27	52	39	62	23
Second test score	47	29	50	44	72	37	63	45	76	32

It is desired to test whether there has been an increase in the population mean score.

- (i) Explain why a two-sample t -test would not be appropriate. [1]

- (ii) Stating any necessary assumptions, carry out a suitable t -test at the $\frac{1}{2}\%$ significance level. [10]

- (iii) The Summer School director claims that after taking the course the population mean score increases by more than 5. Is there sufficient evidence for this claim? [4]

**Copyright Information**

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.