

**PH2 Mark scheme – January 2011**

Question			Marking details	Marks Available
1	(a)	(i)	0.20 m	1
		(ii)	I. 10 m s <sup>-1</sup> [e.c.f.]	1
			II. 0.02 s	1
		III. Displaced wave drawn with same amp and wavelength (1) As 1 <sup>st</sup> marking point with displacement 0.05 m to right (1)	2	
		(iii)	Direction of [particle] oscillation [accept <u>particle</u> movement] and direction of travel [or direction of energy propagation] (1) at right angles (1).	2
	(b)	(i)	Progressive waves transfer energy through medium; stationary waves do not.	1
(ii)		For progressive waves the amplitude doesn't change [or falls gradually] (1) For stationary waves the amplitude increases, decreases and increases (1) [or drops to zero at equally spaced points / nodes]	2	
				<b>[10]</b>
2.	(a)	(i)	Spreads out [or equiv. but not just “bends”]	1
		(ii)	constant phase relationship (1) [between light from slits / sources]	1
	(b)	re-arrangement of formula at any stage (1) [or by impl.] answer correct except, perhaps, for powers of 10 (1) 1.9 m (1)	3	
	(c)	Dark fringes caused by destructive interference (1). With one slit closed, light from the other slit not cancelled [or equiv.](1)	2	
				<b>[7]</b>

Question		Marking details	Marks Available
3	(a)	(i) Formula correctly transposed at any stage (1). $n = 2$ (1); $d = 2.2 \mu\text{m}$ (1)	3
		(ii) Uncertainty [accept error] in measuring angle makes lower uncertainty [accept error] in $d$ .	1
	(b)	(i) $2\lambda = 2.2 \times 10^{-6} \sin 35.1^\circ$ [e.c.f.] (1) [or by impl.] $\lambda = 633 \text{ nm}$ (1)	2
		(ii) <b>Either</b> $\frac{d}{\lambda} = 3.5$ [or $< 4$ ] <b>or</b> $\frac{3\lambda}{d}$ and $\frac{4\lambda}{d}$ evaluated [in an attempt to find $\sin \theta$ ]. (1) [e.c.f. on $d$ or $\lambda$ ] $3^{\text{rd}}$ order deduced by valid reasoning (1).	2
			<b>[8]</b>
4.	(a)	$n_{\text{clad}} \sin 90^\circ = 1.540 \sin 77^\circ$ <b>or</b> $n_{\text{clad}} = 1.540 \sin 77^\circ$ [or by impl.] (1) $n_{\text{clad}} = 1.50$ [1] [accept 1.5] (1)	2
	(b)	(i) speed = $\frac{3.00 \times 10^8}{1.54}$ (1) time = $\frac{\text{distance}}{\text{speed}}$ (1) [transposed at any stage] $= 1.027 \times 10^{-5} \text{ s}$ (1) [omission of 1.54 loses just 1 mark]	3
		(ii) I. $AB = \frac{AC}{\sin 77^\circ}$ <b>or</b> $AB = \frac{AC}{\cos 13^\circ}$ <b>or</b> equiv. (1) II. Zigzag time = $1.027 \times 10^{-5} \times 1.026 \text{ s}$ (1) [ <b>or</b> Extra time = $1.027 \times 10^{-5} \times 0.026$ or by impl.] Extra time = $2.7 \times 10^{-7} \text{ s}$ [e.c.f. on speed] (1)	1 2
	(iii)	Bit of data arrives spread out over a period of time [accept: data smeared or multimode dispersion] (1). Data bits could overlap on arrival / can't distinguish (1)	2

Question		Marking details	Marks Available	
5.	(a)	[minimum] energy needed to eject an electron [from surface]	1	
	(b)	(i) $hf_{\min} = \phi$ [or equiv. or by impl.] (1) $f_{\min} = 5.7 \times 10^{14}$ Hz (1)	2	
		(ii) $E_{k \max} = 6.63 \times 10^{-34} \times 7.0 \times 10^{14} - 3.8 \times 10^{-19}$ [or equiv or by impl.] (1) $= 8.4 \times 10^{-20}$ J (1)	2	
	(c)	(i) Increasing intensity increases number of photons per second [or “photons cannot co-operate”]. (1) But individual photon energy unchanged [or “frequency unchanged”] (1).	2	
		(ii) No. of emitted electrons per second [accept current].	1	
(d)	Increase p.d. from zero (1) until ammeter reads zero (1). Take voltmeter reading, $V$ . (1) Evaluate $eV$ . (1)	4		
			<b>[12]</b>	
6	(a)	(i) $\lambda = \frac{hc}{E}$ [any orientation] [ <b>or</b> $E = hf$ <b>and</b> $f = \frac{c}{\lambda}$ ] (1) $\lambda = 6.33 \times 10^{-7}$ m (( <b>unit</b> ))(1)	2	
		(ii) Red or orange.	1	
		(iii) Arrow shown from top energy level to middle level	1	
	(b)	(i) [Incident or passing] photon (1) of energy $3.14 \times 10^{-19}$ J [or equiv. but not just “of the right energy”] (1)	1	
		(ii) Any 2 × 1 of: <ul style="list-style-type: none"> <li>• coherent ✓</li> <li>• beam nearly parallel ✓</li> <li>• [almost] monochromatic [or same frequency] ✓</li> <li>• polarised ✓</li> </ul>	2	
	(c)	(i) [photons reflected by $M_2$ per second =] $6.3 \times 10^{15}$ [ $s^{-1}$ ] <b>and</b> [photons transmitted per second =] $0.7 \times 10^{15}$ [ $s^{-1}$ ]	1	
		(ii) $0.7 \times 10^{15} s^{-1} \times 3.14 \times 10^{-19}$ J [or by impl.] (1) $= 0.22$ mW (( <b>unit</b> ))(1) [1 mark lost if wrong number of photons used]	2	
		(iii) Stimulated emission event gives 2 photons out for 1 photon in. (1) Many such events as photons traverse amplifying medium [twice] (1) [ <b>or</b> other true <b>and</b> relevant observation]	2	
				<b>[13]</b>

Question			Marking details	Marks Available
7.	(a)	(i)	LHS: lepton number [= 0 + 0] = 0 (1) RHS: lepton number = [0] - 1 + 1 (1) [= 0]	2
		(ii)	I. 4 → 3 II. 2 → 3	1 1
	(b)	weak (1) because of neutrino involvement [ <b>or</b> change in quark flavour] (1)	2	
	(c)	takes place in the Sun (1) first stage in fusion chain [ <b>or</b> ultimately leads to sunshine] (1) <b>Alternatively:</b> <u>has</u> taken place in stars (✓) leading to the formation of heavy elements (✓)	2	
(d)	electro-magnetic	1		
				<b>[9]</b>
8	(a)	(i)	Power = intensity × 4πr <sup>2</sup> (1) = 3.8[5] × 10 <sup>26</sup> W (1) [1 mark lost for factors of 2, 3 or 10 <sup>n</sup> adrift]	2
		(ii)	absorption by atmosphere.	1
	(b)	(i)	$A = \frac{3.85 \times 10^{26}}{5.67 \times 10^{-8} \times 5780^4} \text{ m}^2$ [e.c.f.] (1) = 6.1 × 10 <sup>18</sup> m <sup>2</sup> (1) [6.08 × 10 <sup>18</sup> m <sup>2</sup> ]	2
		(ii)	<b>Either</b> $d = 2\sqrt{\frac{A}{4\pi}}$ [or equiv.] (1) = 1.39 × 10 <sup>9</sup> m (1) <b>Or</b> $A = 4\pi \left[\frac{d}{2}\right]^2$ (1) = 6.15 × 10 <sup>18</sup> m <sup>2</sup> (1)	2
(c)	$\lambda_{\text{max}} = \frac{W}{T} = \frac{2.90 \times 10^{-3} \text{ mK}}{5780 \text{ K}}$ (1) = 500 nm [which is in the visible] (1) Sketch graph of correct general shape (1) with peak at 500 nm [e.c.f.] (1)	4		
				<b>[11]</b>