



# GCE

## Chemistry A

Advanced GCE F325

Equilibria, Energetics and Elements

# Mark Scheme for June 2010

---

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of pupils of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, OCR Nationals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by Examiners. It does not indicate the details of the discussions which took place at an Examiners' meeting before marking commenced.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the Report on the Examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2010

Any enquiries about publications should be addressed to:

OCR Publications  
PO Box 5050  
Annesley  
NOTTINGHAM  
NG15 0DL

Telephone: 0870 770 6622  
Facsimile: 01223 552610  
E-mail: [publications@ocr.org.uk](mailto:publications@ocr.org.uk)

F325

Mark Scheme

June 2010

Question		Expected Answers	Marks	Additional Guidance
1	a	<p>F B G E D</p> <p>FIVE correct ✓✓✓ FOUR correct ✓✓ THREE correct ✓</p>	3	<p><b>ALLOW</b></p> <p>1450 736 76 -642</p> <p>G</p>
	b	<p>Correct calculation  <math>-642 - (+76 + (2 \times 150) + 736 + 1450 + (2 \times -349))</math> ✓  <math>-642 - 1864</math>  <math>= -2506</math> ✓ (kJ mol<sup>-1</sup>)</p>	2	<p><b>ALLOW</b> for 1 mark:  -2705 (2 × 150 and 2 × 349 not used for Cl)  -2356 (2 × 150 not used for Cl)  -2855 (2 × 349 not used for Cl)  +2506 (wrong sign)  <b>DO NOT ALLOW</b> any other answers</p>
	c	<p>Magnesium ion <b>OR</b> Mg<sup>2+</sup>  has greater charge (than sodium ion <b>OR</b> Na<sup>+</sup>)  <b>OR</b> Mg<sup>2+</sup> has greater charge density ✓</p> <p>Magnesium ion <b>OR</b> Mg<sup>2+</sup> is smaller ✓</p> <p>Mg<sup>2+</sup> has a stronger attraction (than Na<sup>+</sup>) to Cl<sup>-</sup> ion  <b>OR</b>  Greater attraction between oppositely charged ions ✓</p>	3	<p><b>ANNOTATIONS MUST BE USED</b></p> <p><b>ALLOW</b> magnesium/Mg is 2+ but sodium/Na is 1+  <b>DO NOT ALLOW</b> Mg atom is 2+ but Na atom is 1+  <b>ALLOW</b> 'charge density' here <b>only</b></p> <p><b>ALLOW</b> Mg <b>OR</b> magnesium is smaller  <b>DO NOT ALLOW</b> Mg<sup>2+</sup> has a smaller <b>atomic</b> radius</p> <p><b>ALLOW</b> anion <b>OR</b> negative ion for Cl<sup>-</sup>  <b>DO NOT ALLOW</b> chlorine ions  <b>DO NOT ALLOW</b> Mg has greater attraction</p> <p><b>ALLOW</b> 'attracts with more force' for greater attraction  but <b>DO NOT ALLOW</b> 'greater force' (could be repulsion)</p> <p><b>ALLOW</b> reverse argument throughout in terms of Na<sup>+</sup></p>
<b>Total</b>			<b>8</b>	



F325

Mark Scheme

June 2010

Question	Expected Answers	Marks	Additional Guidance
	<p><b>Calculation of rate constant (3 marks)</b></p> $k = \frac{\text{rate}}{[\text{BrO}_3^-][\text{Br}^-][\text{H}^+]^2}$ <p><b>OR</b></p> $\frac{1.19 \times 10^{-5}}{(5.0 \times 10^{-2})(1.5 \times 10^{-1})(3.1 \times 10^{-1})^2} \checkmark$ <p><math>= 1.7 \times 10^{-2}</math> <b>OR</b> <math>1.65 \times 10^{-2} \checkmark</math> <math>\text{dm}^9 \text{mol}^{-3} \text{s}^{-1} \checkmark</math></p>	3	<p><b>ANNOTATIONS MUST BE USED</b></p> <p>Calculation can be from any of the experimental runs – they all give the same value of <math>k</math></p> <p><b>ALLOW</b> <math>\text{mol}^{-3} \text{dm}^9 \text{s}^{-1}</math></p> <p><b>ALLOW</b> <math>1.6510579 \times 10^{-2}</math> and correct rounding to <math>1.7 \times 10^{-2}</math></p> <p><b>Correct numerical answer subsumes previous marking point</b></p> <p><b>DO NOT ALLOW</b> fraction: <math>\frac{238}{14415}</math></p> <p>-----</p> <p><b>ALLOW ECF from incorrect rate equation.</b> Examples are given below for 1st line of initial rates data. IF other rows have been used, then calculate the rate constant from data chosen.</p> <p><b>Example 1:</b> 1st order with respect to <math>\text{H}^+</math>  <math>\text{rate} = k [\text{BrO}_3^-] [\text{Br}^-] [\text{H}^+]</math>  <math>k = \frac{\text{rate}}{[\text{BrO}_3^-][\text{Br}^-][\text{H}^+]}</math>  <b>OR</b> <math>\frac{1.19 \times 10^{-5}}{(5.0 \times 10^{-2})(1.5 \times 10^{-1})(3.1 \times 10^{-1})} \checkmark</math>  <math>= 5.1 \times 10^{-3}</math> <b>OR</b> <math>5.12 \times 10^{-3} \checkmark</math> <math>\text{dm}^6 \text{mol}^{-2} \text{s}^{-1} \checkmark</math>  <b>ALLOW</b> <math>5.11827957 \times 10^{-3}</math> and correct rounding to <math>5.1 \times 10^{-3}</math></p> <p>-----</p> <p><b>Example 2:</b> Zero order with respect to <math>\text{BrO}_3^-</math>  <math>\text{rate} = k [\text{Br}^-] [\text{H}^+]^2</math>  <math>k = \frac{\text{rate}}{[\text{Br}^-][\text{H}^+]^2}</math>  <b>OR</b> <math>\frac{1.19 \times 10^{-5}}{(1.5 \times 10^{-1})(3.1 \times 10^{-1})^2} \checkmark</math>  <math>= 8.3 \times 10^{-4}</math> <b>OR</b> <math>8.26 \times 10^{-4} \checkmark</math> <math>\text{dm}^6 \text{mol}^{-2} \text{s}^{-1} \checkmark</math>  <b>ALLOW</b> <math>8.255289629 \times 10^{-4}</math> and correct rounding to <math>8.3 \times 10^{-4}</math></p>
	<b>Total</b>	<b>10</b>	

F325

Mark Scheme

June 2010

Question		Expected Answers	Marks	Additional Guidance
3	a	<p>measured pH &gt; 1 <b>OR</b> <math>[H^+] &lt; 0.1</math> (mol dm<sup>-3</sup>) ✓</p> <p><math>[H^+] = 10^{-pH}</math> ✓</p> <p><math>K_a = \frac{[H^+][CH_3CH_2COO^-]}{[CH_3CH_2COOH]}</math> <b>OR</b> <math>\frac{[H^+]^2}{[CH_3CH_2COOH]}</math> ✓</p> <p>Calculate <math>K_a</math> from <math>\frac{[H^+]^2}{0.100}</math> ✓</p>	4	<p><b>ALLOW</b> C<sub>2</sub>H<sub>5</sub> throughout question</p> <p><b>ALLOW</b> <math>[H^+] &lt; [CH_3CH_2COOH]</math> <b>OR</b> <math>[H^+] &lt; [HA]</math></p> <p><b>ALLOW</b> measured pH is higher than expected</p> <p><b>ALLOW</b> measured pH is not as acidic as expected</p> <p><b>ALLOW</b> a quoted pH value or range &gt; 1 and &lt; 7</p> <p><b>OR</b> between 1 and 7</p> <p><b>ALLOW</b> <math>[H^+] = \text{antilog } -pH</math> <b>OR</b> <math>[H^+] = \text{inverse log } -pH</math></p> <p><b>ALLOW</b> <math>\frac{[H^+][A^-]}{[HA]}</math> <b>OR</b> <math>\frac{[H^+]^2}{[HA]}</math></p> <p><b>IF</b> <math>K_a</math> is <b>NOT</b> given and <math>K_a = \frac{[H^+]^2}{0.100}</math> is shown, award mark for <math>K_a</math> also</p> <p>(i.e. <math>K_a = \frac{[H^+]^2}{0.100}</math> is automatically awarded the last 2 marks)</p>
	b	<p><b>Marks are for correctly calculated values.</b></p> <p><b>Working shows how values have been derived.</b></p> <p><math>[H^+] = 10^{-13.46} = 3.47 \times 10^{-14}</math> (mol dm<sup>-3</sup>) ✓</p> <p><math>[OH^-] = \frac{1.0 \times 10^{-14}}{3.47 \times 10^{-14}} = 0.29</math> (mol dm<sup>-3</sup>) ✓</p>	2	<p><b>ALLOW</b> <math>3.467368505 \times 10^{-14}</math> and correct rounding to <math>3.5 \times 10^{-14}</math></p> <p><b>ALLOW</b> 0.28840315 and correct rounding to 0.29, i.e. <b>ALLOW</b> 0.288</p> <p><b>ALLOW</b> alternative approach using pOH:</p> <p>pOH = 14 – 13.46 = 0.54 ✓</p> <p><math>[OH^-] = 10^{-0.54} = 0.29</math> (mol dm<sup>-3</sup>) ✓</p> <p>Correct answer gets <b>BOTH</b> marks</p>



F325

Mark Scheme

June 2010

Question		Expected Answers	Marks	Additional Guidance	
	d	$\text{HNO}_3 + \text{CH}_3\text{CH}_2\text{COOH} \rightleftharpoons \text{CH}_3\text{CH}_2\text{COOH}_2^+ + \text{NO}_3^- \checkmark$ <p style="text-align: center;">acid 1      base 2                  acid 2                  base 1    <math>\checkmark</math></p>	2	<p>State symbols <b>NOT</b> required</p> <p><b>ALLOW 1 AND 2</b> labels the other way around.</p> <p><b>ALLOW</b> 'just acid' and 'base' labels throughout if linked by lines so that it is clear what the acid–base pairs are.</p> <p><b>IF</b> proton transfer is wrong way around then <b>ALLOW</b> 2nd mark for idea of acid–base pairs, i.e.</p> $\text{HNO}_3 + \text{CH}_3\text{CH}_2\text{COOH} \rightleftharpoons \text{CH}_3\text{CH}_2\text{COO}^- + \text{H}_2\text{NO}_3^+ \times$ <p style="text-align: center;">base 2      acid 1                  base 1      acid 2    <math>\checkmark</math></p>	
	e	i	$2\text{CH}_3\text{CH}_2\text{COOH} + \text{Mg} \rightarrow (\text{CH}_3\text{CH}_2\text{COO})_2\text{Mg} + \text{H}_2 \checkmark$	1	<p><b>IGNORE</b> state symbols</p> <p><b>ALLOW</b> ionic equation: <math>2\text{H}^+ + \text{Mg} \rightarrow \text{Mg}^{2+} + \text{H}_2</math></p> <p><b>IGNORE</b> any random charges in formula of <math>(\text{CH}_3\text{CH}_2\text{COO})_2\text{Mg}</math> as long as the charges are <b>correct (charges are treated as working)</b> i.e. <math>(\text{CH}_3\text{COO}^-)_2\text{Mg}</math> <b>OR</b> <math>(\text{CH}_3\text{COO})_2^- \text{Mg}</math> should <b>not</b> be penalised However, <math>\text{Mg}^{2+}</math> instead of Mg on the left side of equation is obviously wrong</p>
		ii	$2\text{H}^+ + \text{CO}_3^{2-} \longrightarrow \text{H}_2\text{O} + \text{CO}_2$ <p><b>OR</b> <math>2\text{H}^+ + \text{CO}_3^{2-} \longrightarrow \text{H}_2\text{CO}_3</math></p> <p><b>OR</b> <math>\text{H}^+ + \text{CO}_3^{2-} \longrightarrow \text{HCO}_3^- \checkmark</math></p>	1	State symbols <b>NOT</b> required
<b>Total</b>			<b>17</b>		



F325

Mark Scheme

June 2010

Question			Expected Answers	Marks	Additional Guidance
4	a	i	Complete circuit (with voltmeter) and salt bridge linking two half-cells ✓ Pt electrode in solution of Fe <sup>2+</sup> /Fe <sup>3+</sup> ✓ Ag in solution of Ag <sup>+</sup> ✓	3	<b>DO NOT ALLOW</b> 'solution of a silver halide', e.g. AgCl (as these are insoluble) but <b>DO ALLOW</b> any solution of any other silver salt (whether insoluble or not)  <b>IF</b> candidate has used incorrect redox systems, then mark ECF as follows: <b>(i) each</b> incorrect system will cost the candidate <b>one</b> mark <b>(ii) ECF</b> if species have been quoted (see Additional Guidance below) <b>(iii) ECF</b> for equation <b>(iv) ECF</b> for cell potential <b>YOU MAY NEED TO WORK OUT THESE ECF RESPONSES YOURSELF DEPENDING ON THE INCORRECT REDOX SYSTEMS CHOSEN</b>
		ii	electrons <b>AND</b> ions ✓	1	For electrons, <b>ALLOW</b> e <sup>-</sup> For 'ions', <b>ALLOW</b> formula of an ion in one of the half-cells or salt bridge, e.g. Ag <sup>+</sup> , Fe <sup>2+</sup> , Fe <sup>3+</sup> <b>ALLOW ECF</b> as in (i)
		iii	Ag + Fe <sup>3+</sup> → Ag <sup>+</sup> + Fe <sup>2+</sup> ✓	1	<b>ALLOW ECF</b> as in (i) <b>ALLOW</b> equilibrium sign
		iv	0.43 V ✓	1	<b>ALLOW ECF</b> as in (i)
	b	i	Cl <sub>2</sub> <b>OR</b> O <sub>2</sub> <b>AND</b> H <sup>+</sup> ✓	1	<b>ALLOW</b> chlorine <b>ALLOW</b> O <sub>2</sub> <b>AND</b> 4H <sup>+</sup> <b>ALLOW</b> O <sub>2</sub> <b>AND</b> acid <b>DO NOT ALLOW</b> O <sub>2</sub> alone <b>DO NOT ALLOW</b> equation or equilibrium
		ii	I <sup>-</sup> ✓	1	<b>ALLOW</b> 2I <sup>-</sup> <b>OR</b> iodide <b>DO NOT ALLOW</b> equation or equilibrium

F325

Mark Scheme

June 2010

Question	Expected Answers	Marks	Additional Guidance
c	<p>A fuel cell converts energy from reaction of a fuel with oxygen into a voltage/electrical energy ✓</p> <p><math>2\text{H}_2 + \text{O}_2 \rightarrow 2\text{H}_2\text{O}</math> ✓</p> <p>Two from:</p> <ul style="list-style-type: none"> <li>• under pressure <b>OR</b> at low temperature <b>OR</b> as a liquid</li> <li>• adsorbed on solid</li> <li>• absorbed within solid</li> </ul> <p style="text-align: right;">✓✓</p> <p>Energy is needed to make the hydrogen <b>OR</b> energy is needed to make fuel cell ✓</p>	5	<p><b>ANNOTATIONS MUST BE USED</b></p> <p><b>ALLOW</b> combustion for reaction of fuel with oxygen/reactants</p> <p><b>ALLOW</b> a fuel cell requires constant supply of fuel</p> <p><b>OR</b> operates continuously as long as a fuel (and oxygen) are added</p> <p><b>ALLOW</b> multiples, e.g. <math>\text{H}_2 + \frac{1}{2}\text{O}_2 \rightarrow \text{H}_2\text{O}</math></p> <p><b>IGNORE</b> state symbols</p> <p><b>ALLOW</b> 'material' <b>OR</b> metal for solid</p> <p><b>ALLOW</b> as a metal hydride</p>
	<b>Total</b>	<b>13</b>	

F325

Mark Scheme

June 2010

Question			Expected Answers	Marks	Additional Guidance
5	a	i	$(K_c =) \frac{[\text{NH}_3]^2}{[\text{N}_2][\text{H}_2]^3} \checkmark$	1	Must be square brackets
		ii	$\text{dm}^6 \text{mol}^{-2} \checkmark$	1	<b>ALLOW</b> $\text{mol}^{-2} \text{dm}^6$ <b>ALLOW ECF</b> from incorrect $K_c$ expression
	b		<p><b>Unless otherwise stated, marks are for correctly calculated values. Working shows how values have been derived.</b></p> <p><math>[\text{N}_2] = \frac{7.2}{6.0}</math> <b>OR</b> <math>1.2 \text{ (mol dm}^{-3}\text{)}</math></p> <p><b>AND</b> <math>[\text{H}_2] = \frac{12}{6.0}</math> <b>OR</b> <math>2.0 \text{ (mol dm}^{-3}\text{)}</math> <math>\checkmark</math></p> <p><math>[\text{NH}_3] = \sqrt{(K_c \times [\text{N}_2] \times [\text{H}_2]^3)}</math></p> <p><b>OR</b> <math>\sqrt{(8.00 \times 10^{-2} \times 1.2 \times 2.0^3)}</math> <math>\checkmark</math></p> <p><math>= 0.876</math> <b>OR</b> <math>0.88 \text{ (mol dm}^{-3}\text{)}</math> <math>\checkmark</math></p> <p>amount <math>\text{NH}_3 = 0.876 \times 6 = 5.26</math> <b>OR</b> <math>5.3 \text{ (mol)}</math> <math>\checkmark</math></p>	4	<p><b>ANNOTATIONS MUST BE USED</b></p> <p>For <b>all</b> parts, <b>ALLOW</b> numerical answers from 2 significant figures up to the calculator value</p> <p>1st mark is for realising that concentrations need to be calculated.</p> <p><b>Correct numerical answer with no working would score all previous calculation marks</b></p> <p><b>ALLOW</b> calculator value: 0.876356092 down to 0.88, correctly rounded</p> <p><b>ALLOW</b> calculator value down to 5.3, correctly rounded</p>

F325

Mark Scheme

June 2010

Question	Expected Answers	Marks	Additional Guidance
b	<p><b>EXAMPLES OF INCORRECT RESPONSES IN (b) THAT MAY BE WORTHY OF CREDIT</b></p>		<p>-----</p> <p><b>ALLOW ECF from incorrect concentrations (3 marks)</b> For example, If concentrations <b>not</b> calculated at start, then</p> $[\text{NH}_3] = \sqrt{(8.00 \times 10^{-2} \times 7.2 \times 12.0^3)} \checkmark$ $= 31.5 \text{ mol dm}^{-3} \checkmark$ <p>Equilibrium amount of <math>\text{NH}_3 = 31.5 \times 6 = 189.6 \text{ (mol)} \checkmark</math></p> <p>-----</p> <p><b>IF candidate has <math>K_c</math> expression upside down, then all 4 marks are available in (b) by ECF</b></p> <p>Correct <math>[\text{N}_2]</math> AND <math>[\text{H}_2] \checkmark</math></p> $[\text{NH}_3] = \sqrt{\frac{[\text{N}_2][\text{H}_2]^3}{K_c}} = \sqrt{\frac{1.2 \times 2^3}{8.00 \times 10^{-2}}} \checkmark$ $= 11.0 \text{ mol dm}^{-3} \checkmark$ <p>Equilibrium amount of <math>\text{NH}_3 = 11.0 \times 6 = 66.0 \text{ (mol)} \checkmark</math></p> <p>-----</p> <p><b>IF candidate has used <math>K_c</math> value of <math>8.00 \times 10^{-2}</math> AND values for <math>\text{N}_2</math> AND <math>\text{H}_2</math> with powers wrong, mark by ECF from calculated as below (3 max in (b))</b></p> <p>Correct <math>[\text{N}_2]</math> AND <math>[\text{H}_2] \checkmark</math></p> <p><math>[\text{NH}_3]</math> expression ✗</p> <p><b>ECF:</b> Calculated <math>[\text{NH}_3] \checkmark</math></p> <p><b>ECF:</b> Equilibrium amount of <math>\text{NH}_3 \checkmark</math></p>

F325

Mark Scheme

June 2010

Question		Expected Answers	Marks	Additional Guidance
	c i	Equilibrium shifts to right OR Equilibrium towards ammonia ✓  Right hand side has fewer number of (gaseous) moles ✓	2	<b>ALLOW</b> 'moves right' OR 'goes right' OR 'favours right' OR 'goes forwards'  <b>ALLOW</b> 'ammonia side' has fewer moles <b>ALLOW</b> 'there are more (gaseous) moles on left'
	ii	$K_c$ does not change ✓  Increased pressure increases concentration terms on bottom of $K_c$ expression more than the top <b>OR</b> system is now no longer in equilibrium ✓  top of $K_c$ expression increases and bottom decreases until $K_c$ is reached ✓	3	<b>ANNOTATIONS MUST BE USED</b> Any response in terms of $K_c$ changing scores <b>ZERO</b> for Part (ii) <b>ALLOW</b> $K_c$ is temperature dependent only OR $K_c$ does not change with pressure  <b>ALLOW</b> $\frac{[\text{NH}_3]^2}{[\text{N}_2][\text{H}_2]^3}$ no longer equal to $K_c$
	d i	$\text{CH}_4 + \text{H}_2\text{O} \longrightarrow 3\text{H}_2 + \text{CO}$ ✓	1	State symbols <b>NOT</b> required <b>ALLOW</b> : $\text{CH}_4 + \text{H}_2\text{O} \longrightarrow \text{CH}_3\text{OH} + \text{H}_2$ $\text{CH}_4 + 2\text{H}_2\text{O} \longrightarrow 4\text{H}_2 + \text{CO}_2$ $\text{CH}_4 + \text{H}_2\text{O} \longrightarrow 2\text{H}_2 + \text{HCHO}$ $\text{CH}_4 + 2\text{H}_2\text{O} \longrightarrow 3\text{H}_2 + \text{HCOOH}$
	ii	Electrolysis of water OR $\text{H}_2\text{O} \longrightarrow \text{H}_2 + \frac{1}{2}\text{O}_2$ ✓	1	<b>ALLOW</b> electrolysis of brine <b>DO NOT ALLOW</b> reforming <b>DO NOT ALLOW</b> cracking <b>DO NOT ALLOW</b> reaction of metal with acid

F325

Mark Scheme

June 2010

Question		Expected Answers	Marks	Additional Guidance
	e i	<p><b>Unless otherwise stated, marks are for correctly calculated values. Working shows how values have been derived.</b></p> $\Delta S = \Sigma S(\text{products}) - \Sigma S(\text{reactants}) /$ $= (2 \times 192) - (191 + 3 \times 131) \checkmark$ $= -200 \text{ (J K}^{-1} \text{ mol}^{-1}) \text{ OR } -0.200 \text{ (kJ K}^{-1} \text{ mol}^{-1}) \checkmark$ <p>Use of 298 K (could be within <math>\Delta G</math> expression below) <math>\checkmark</math></p> $\Delta G = \Delta H - T\Delta S$ <p><b>OR</b></p> $\Delta G = -92 - (298 \times -0.200)$ <p><b>OR</b></p> $\Delta G = -92000 - (298 \times -200) \checkmark$ $= -32.4 \text{ kJ mol}^{-1} \text{ OR } -32400 \text{ J mol}^{-1} \checkmark$ <p><b>(Units must be shown)</b></p> <p>For feasibility, <math>\Delta G &lt; 0</math> <b>OR</b> <math>\Delta G</math> is negative <math>\checkmark</math></p>	<p>5</p> <p>1</p>	<p><b>ANNOTATIONS MUST BE USED</b></p> <p>See <b>Appendix 1</b> for extra guidance for marking <b>5e(i)</b> and <b>5e(ii)</b></p> <p><b>NO UNITS</b> required at this stage <b>IGNORE</b> units</p> <p><b>ALLOW</b> <math>-32.4 \text{ kJ}</math> <b>OR</b> <math>-32400 \text{ J}</math> (<b>Units must be shown</b>) Award all 5 marks <b>above</b> for correct answer with no working</p> <p><b>IF</b> 25 °C has been used instead of 298 K, correctly calculated <math>\Delta G</math> values are = <math>-87 \text{ kJ mol}^{-1}</math> <b>OR</b> <math>-87000 \text{ J mol}^{-1}</math> <b>4 marks</b> are still available up to this point and maximum possible from <b>(e)(i)</b> is 5 marks</p>
	ii	<p>As the temperature increases, <math>T\Delta S</math> becomes more negative <b>OR</b> <math>T\Delta S</math> becomes more negative than <math>\Delta H</math> <b>OR</b> <math>T\Delta S</math> becomes more significant <math>\checkmark</math></p> <p>Eventually <math>\Delta H - T\Delta S</math> becomes positive <math>\checkmark</math></p>	<p>2</p>	<p><b>ALLOW</b> <math>T\Delta S &gt; \Delta H</math> (i.e. assume no sign at this stage) <b>ALLOW</b> 'entropy term' as alternative for <math>T\Delta S</math> <b>ALLOW</b> <math>-T\Delta S</math> becomes more positive <b>ALLOW</b> <math>-T\Delta S</math> decreases</p> <p><b>ALLOW</b> <math>\Delta G</math> becomes positive <b>OR</b> <math>\Delta G &gt; 0</math></p>

F325

Mark Scheme

June 2010

Question		Expected Answers	Marks	Additional Guidance
	iii	Activation energy is too high <b>OR</b> reaction too slow ✓	1	<b>ALLOW</b> increases the rate <b>OR</b> more molecules exceed activation energy <b>OR</b> more successful collisions <b>ALLOW</b> rate constant increases <b>IGNORE</b> comments on yield
<b>Total</b>			<b>22</b>	

F325

Mark Scheme

June 2010




Question			Expected Answers	Marks	Additional Guidance
6	a	i	$1s^2 2s^2 2p^6 3s^2 3p^6 3d^5 4s^1$ ✓	1	<b>ALLOW</b> $1s^2 2s^2 2p^6 3s^2 3p^6 4s^1 3d^5$ (i.e. 4s before 3d) <b>ALLOW</b> $[Ar]4s^1 3d^5$ <b>OR</b> $[Ar]3d^5 4s^1$
		ii	$1s^2 2s^2 2p^6 3s^2 3p^6 3d^3$ ✓	1	<b>ALLOW</b> $[Ar]3d^3$ <b>ALLOW</b> $1s^2 2s^2 2p^6 3s^2 3p^6 3d^3 4s^0$ <b>OR</b> $[Ar]3d^3 4s^0$
	b		$Zn \longrightarrow Zn^{2+} + 2e^-$ ✓ $Cr_2O_7^{2-} + 14H^+ + 8e^- \longrightarrow 2Cr^{2+} + 7H_2O$ ✓  $4Zn + Cr_2O_7^{2-} + 14H^+ \longrightarrow 4Zn^{2+} + 2Cr^{2+} + 7H_2O$ ✓	3	<b>ALLOW</b> multiples  <b>WATCH</b> for balancing of the equations printed on paper <b>IF</b> printed equations and answer lines have different balancing numbers <b>OR</b> electrons, <b>IGNORE</b> numbers on printed equations (i.e. treat these as working) and mark responses on answer lines <b>only</b>  <b>NO ECF</b> for overall equation i.e. the expected answer is the <b>ONLY</b> acceptable answer
	c	i	Ligand substitution ✓	1	<b>ALLOW</b> ligand exchange
		ii	$[Cr(H_2O)_6]^{3+} + 6NH_3 \longrightarrow [Cr(NH_3)_6]^{3+} + 6H_2O$ ✓ ✓	2	1 mark is awarded for each side of equation <b>ALLOW</b> equilibrium sign <b>ALLOW</b> 1 mark for 2+ shown instead of 3+ on both sides of equation <b>ALLOW</b> 1 mark for substitution of 4 $NH_3$ : $[Cr(H_2O)_6]^{3+} + 4NH_3 \longrightarrow [Cr(NH_3)_4(H_2O)_2]^{3+} + 4H_2O$
	d	i	Donates an electron pair to a metal ion <b>OR</b> forms a coordinate bond to a metal ion ✓	1	<b>ALLOW</b> donates an electron pair to a metal <b>ALLOW</b> dative (covalent) bond for coordinate bond
		ii	Donates <b>two</b> electron pairs <b>OR</b> forms <b>two</b> coordinate bonds ✓  Lone pairs on two O atoms ✓	2	First mark is for the idea of two coordinate bonds  <b>ALLOW</b> lone pair on O and N <b>DO NOT ALLOW</b> lone pairs on $COO^-$ (could involve C)  Second mark is for the atoms that donate the electron pairs Look for the atoms with lone pairs also on response to <b>(d)(iii)</b> and credit here if not described in <b>(d)(ii)</b>



F325

Mark Scheme

June 2010

Question	Expected Answers	Marks	Additional Guidance
iii	<p>Forms two optical isomers <b>OR</b> two enantiomers  <b>OR</b> two non-superimposable mirror images ✓</p>  <p>✓✓ For each structure</p>	3	<p><b>IGNORE</b> any charges shown</p> <p><b>ALLOW</b> any attempt to show bidentate ligand.  Bottom line is the diagram on the left.</p> <p>1 mark for 3D diagram with ligands attached for <b>ONE</b> stereoisomer.  Must contain 2 out wedges, 2 in wedges and 2 lines in plane of paper:</p>  <p>2nd mark for reflected diagram of <b>SECOND</b> stereoisomer.  The diagram below would score the 2nd mark but not the first</p> 

F325

Mark Scheme

June 2010

Question	Expected Answers	Marks	Additional Guidance
e	<p style="text-align: center;">N : H : Cr : O 11.1/14 : 3.17/1 : 41.27/52 : 44.45/16</p> <p><b>OR</b> 0.793 : 3.17 : 0.794 : 2.78 ✓</p> <p><b>A:</b> <math>\text{N}_2\text{H}_8\text{Cr}_2\text{O}_7</math> ✓</p> <p>Ions: <math>\text{NH}_4^+</math> ✓ <math>\text{Cr}_2\text{O}_7^{2-}</math> ✓</p> <p><b>B:</b> <math>\text{Cr}_2\text{O}_3</math> ✓</p> <p>Correctly calculates molar mass of <b>C</b> <math>= 1.17 \times 24.0 = 28.08 \text{ (g mol}^{-1}\text{)}</math> ✓</p> <p><b>C:</b> <math>\text{N}_2</math> ✓</p> <p>Equation: <math>(\text{NH}_4)_2\text{Cr}_2\text{O}_7 \longrightarrow \text{Cr}_2\text{O}_3 + 4\text{H}_2\text{O} + \text{N}_2</math> ✓</p>	8	<p><b>ANNOTATIONS MUST BE USED</b></p> <p><b>ALLOW A:</b> <math>(\text{NH}_4)_2\text{Cr}_2\text{O}_7</math></p> <p><b>IF</b> candidate has obtained <math>\text{NH}_4\text{CrO}_4</math> for A, <b>ALLOW</b> <math>\text{NH}_4^+</math> <b>DO NOT ALLOW</b> <math>\text{CrO}_4^-</math></p> <p><b>ALLOW:</b> (relative) molecular mass <b>ALLOW:</b> 28 <b>ALLOW:</b> 'C is 28'</p> <p><b>ALLOW</b> <math>\text{N}_2\text{H}_8\text{Cr}_2\text{O}_7</math> in equation.</p>
<b>Total</b>		<b>22</b>	

F325

Mark Scheme

June 2010

Question	Expected Answers	Marks	Additional Guidance
7 a i	$\text{H}_2\text{O}_2 \longrightarrow \text{O}_2 + 2\text{H}^+ + 2\text{e}^- \checkmark\checkmark$	2	All other multiples score 1 mark e.g. $\frac{1}{2} \text{H}_2\text{O}_2 \longrightarrow \frac{1}{2} \text{O}_2 + \text{H}^+ + \text{e}^-$ $5\text{H}_2\text{O}_2 \longrightarrow 5\text{O}_2 + 10\text{H}^+ + 10\text{e}^-$
b	<p><b>Marks are for correctly calculated values. Working shows how values have been derived.</b></p> $n(\text{KMnO}_4) = \frac{0.0200 \times 23.45}{1000} = 4.69 \times 10^{-4} \text{ (mol) } \checkmark$ $n(\text{H}_2\text{O}_2) = 5/2 \times 4.69 \times 10^{-4} = 1.1725 \times 10^{-3} \text{ (mol) } \checkmark$ $n(\text{H}_2\text{O}_2) \text{ in } 250 \text{ cm}^3 \text{ solution}$ $= 10 \times 1.1725 \times 10^{-3} = 1.1725 \times 10^{-2} \text{ (mol) } \checkmark$ $\text{concentration in g dm}^{-3} \text{ of original H}_2\text{O}_2$ $= 40 \times 1.1725 \times 10^{-2} \times 34 = 15.9 \text{ (g dm}^{-3}) \checkmark$ $n(\text{O}_2) = 5/2 \times 4.69 \times 10^{-4} = 1.1725 \times 10^{-3} \text{ (mol) } \checkmark$ $\text{volume O}_2 = 24.0 \times 1.1725 \times 10^{-3} = 0.0281 \text{ dm}^3 \checkmark$	<p>4</p> <p>2</p>	<p><b>ANNOTATIONS MUST BE USED</b></p> <p><b>DO NOT ALLOW</b> <math>4.7 \times 10^{-4}</math></p> <p><b>ALLOW</b> <math>1.173 \times 10^{-3}</math> OR <math>1.17 \times 10^{-3}</math> (i.e. 3 significant figures upwards)</p> <p><b>ALLOW</b> by <b>ECF</b>: <math>5/2 \times</math> ans above</p> <p><b>ALLOW</b> by <b>ECF</b> <math>10 \times</math> ans above</p> <p><b>ALLOW</b> concentration <math>\text{H}_2\text{O}_2 = 0.0469 \text{ mol dm}^{-3}</math></p> <p><b>ALLOW</b> by <b>ECF</b> <math>40 \times n(\text{H}_2\text{O}_2) \times 34</math></p> <p><b>ALLOW</b> <math>0.0469 \times 10 \times 34 = 15.9 \text{ g dm}^{-3} \checkmark</math></p> <p><b>ALLOW</b> two significant figures, 16 (<math>\text{g dm}^{-3}</math>) up to calculator value of 15.946 <math>\text{g dm}^{-3}</math></p> <p><b>ALLOW</b> <math>0.028 \text{ dm}^3</math> OR <math>0.02814 \text{ dm}^3</math> <b>ALLOW</b> <math>28 \text{ cm}^3</math> OR <math>28.14 \text{ cm}^3</math> Value <b>AND</b> units required <b>DO NOT ALLOW</b> <math>0.03 \text{ dm}^3</math></p> <p><b>ALLOW</b> by <b>ECF</b>: <math>24.0 \times</math> calculated moles of <math>\text{O}_2</math> (2 significant figures up to calculator value)</p>
<b>Total</b>		<b>8</b>	

F325

Mark Scheme

June 2010

Appendix 1

Extra guidance for marking atypical responses to **5e(i)** and **5e(ii)**

Question	Expected Answer	Mark	Additional Guidance
5 e i	<p><b>TOTAL ENTROPY APPROACH:</b>  <b>ALL MARKS AVAILABLE</b>                      Unless otherwise stated, marks are for correctly calculated values.                      Working shows how values have been derived.</p> $\Delta S = \Sigma S(\text{products}) - \Sigma S(\text{reactants}) /$ $= (2 \times 192) - (191 + 3 \times 131) \checkmark$ $= -200 (\text{J K}^{-1} \text{mol}^{-1}) \text{ OR } -0.200 (\text{kJ K}^{-1} \text{mol}^{-1}) \checkmark$ <p>Use of 298 K (could be within expression below) <math>\checkmark</math></p> $\Delta S_{\text{total}} = \Delta S_{\text{system}} + \Delta S_{\text{surroundings}}$ $\Delta S_{\text{surroundings}} = - \frac{\Delta H}{T}$ <p><b>OR</b> <math display="block">\Delta S_{\text{total}} = \Delta S_{\text{system}} - \frac{\Delta H}{T}</math></p> <p><b>OR</b> <math display="block">\Delta S_{\text{total}} = -0.200 - \frac{-92}{298}</math></p> <p><b>OR</b> <math display="block">\Delta S_{\text{total}} = -200 - \frac{-92000}{298} \checkmark</math></p> $= 0.109 \text{ kJ (K}^{-1} \text{mol}^{-1}) \text{ OR } 109 \text{ J (K}^{-1} \text{mol}^{-1}) \checkmark$ <p>Feasible when <math>\Delta S_{\text{total}} &gt; 0 \checkmark</math></p>	<p>5</p> <p>1</p>	<p><b>ANNOTATIONS MUST BE USED</b></p> <p><b>NO UNITS</b> required at this stage  <b>IGNORE</b> units</p> <p><b>ALLOW</b> 0.109 kJ <b>OR</b> 109 J  <b>IF</b> 25°C has been used instead of 298 K, correctly calculated <math>\Delta S_{\text{total}}</math> values are = 3.48 kJ K<sup>-1</sup> mol<sup>-1</sup> <b>OR</b> 3,480 J K<sup>-1</sup> mol<sup>-1</sup></p>

F325

Mark Scheme

June 2010

Question	Expected Answer	Mark	Additional Guidance
5 e i	<p><b>MAX/MIN TEMPERATURE APPROACH: 5 MARKS MAX AVAILABLE</b></p> <p>Unless otherwise stated, marks are for correctly calculated values. Working shows how values have been derived.</p> $\Delta S = \Sigma S(\text{products}) - \Sigma S(\text{reactants}) /$ $= (2 \times 192) - (191 + 3 \times 131) \checkmark$ $= -200 \text{ (J K}^{-1} \text{ mol}^{-1}) \text{ OR } -0.200 \text{ (kJ K}^{-1} \text{ mol}^{-1}) \checkmark$ <p>Use of 298 K (could be within <math>\Delta G</math> expression below) <math>\checkmark</math></p> $\Delta G = \Delta H - T\Delta S$ <p>OR When <math>\Delta G = 0</math>, <math>0 = \Delta H - T\Delta S</math>;</p> <p>OR <math>T = \frac{\Delta H}{\Delta S} = \frac{-92}{-0.200}</math></p> <p>OR <math>T = \frac{\Delta H}{\Delta S} = \frac{-92000}{-200} \checkmark</math></p> $= 460 \text{ K } \checkmark$ $= 187 \text{ }^\circ\text{C (use of 298) } \checkmark$ <p>The condition <math>\Delta G = 0</math> because temperature at which <math>\Delta G = 0</math> is the maximum temperature for feasibility <b>AND</b> justification for the being the maximum <math>\checkmark</math></p>		<p><b>ANNOTATIONS MUST BE USED</b></p> <p>This candidate has not answered the question but many marks are still available.</p> <p><b>NO UNITS</b> required at this stage <b>IGNORE</b> units</p> <p>By this approach, the calculated temperature is the switchover between feasibility and non-feasibility but it cannot be assumed that this is the maximum temperature</p>

F325

Mark Scheme

June 2010

Question			Expected Answer	Mark	Additional Guidance
5	e	ii	As the temperature increases, $\Delta H/T$ becomes <b>less</b> negative <b>OR</b> $\Delta H/T$ becomes <b>more</b> negative than $\Delta S(\text{system})$ <b>OR</b> $\Delta H/T$ becomes <b>less</b> significant <b>OR</b> $\Delta S(\text{surroundings})$ becomes <b>less</b> significant <b>OR</b> $\Delta S(\text{system}) > \Delta H/T$ <b>OR</b> $\Delta S(\text{system}) > \Delta S(\text{surroundings})$ ✓  Eventually $\Delta S(\text{total})$ becomes <b>negative</b> ✓	2	<b>ALLOW</b> $\Delta H/T > \Delta S_{\text{system}}$ (i.e. assume no sign at this stage)  <b>ALLOW</b> $-\Delta H/T$ becomes more positive <b>ALLOW</b> $-\Delta H/T$ increases

**OCR (Oxford Cambridge and RSA Examinations)**  
**1 Hills Road**  
**Cambridge**  
**CB1 2EU**

**OCR Customer Contact Centre**

**14 – 19 Qualifications (General)**

Telephone: 01223 553998

Facsimile: 01223 552627

Email: [general.qualifications@ocr.org.uk](mailto:general.qualifications@ocr.org.uk)

**[www.ocr.org.uk](http://www.ocr.org.uk)**

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

**Oxford Cambridge and RSA Examinations**  
**is a Company Limited by Guarantee**  
**Registered in England**  
**Registered Office; 1 Hills Road, Cambridge, CB1 2EU**  
**Registered Company Number: 3484466**  
**OCR is an exempt Charity**

**OCR (Oxford Cambridge and RSA Examinations)**  
**Head office**  
**Telephone: 01223 552552**  
**Facsimile: 01223 552553**

© OCR 2010

