

## **GCE A level**

1324/01

# PHYSICS – PH4 Oscillations and Fields

P.M. MONDAY, 20 January 2014

1 hour 30 minutes

For Examiner's use only					
Question	Maximum Mark	Mark Awarded			
1.	8				
2.	8				
3.	16				
4.	14				
5.	11				
6.	13				
7.	10				
Total	80				

#### **ADDITIONAL MATERIALS**

In addition to this examination paper, you will require a calculator and a Data Booklet.

### **INSTRUCTIONS TO CANDIDATES**

Use black ink or black ball-point pen.

Write your name, centre number and candidate number in the spaces at the top of this page.

Answer all questions.

Write your answers in the spaces provided in this booklet.

#### INFORMATION FOR CANDIDATES

The total number of marks available for this paper is 80.

The number of marks is given in brackets at the end of each question or part question.

You are reminded of the necessity for good English and orderly presentation in your answers.


You are reminded to show all working. Credit is given for correct working even when the final answer given is incorrect.

## Answer all questions.

Examiner only

1. (a) State the principle of conservation of momentum. [2]

A nickel nucleus emits a  $\gamma$  ray photon as shown.



(b) Show that the  $\gamma$  ray photon has a momentum of approximately  $7 \times 10^{-22} \, \text{kg} \, \text{m} \, \text{s}^{-1}$ . [3]

(c)	Calculate the final speed (v) of the nickel nucleus ( $m_{\text{nickel}} = 9.95 \times 10^{-26} \text{kg}$ ).	[3]
•••••		
•••••		······································
*********		
•••••		•••••••••••••••••••••••••••••••••••••••

Examiner only

**2.** (a) Using the equations:

$$p = \frac{1}{3}\rho \overline{c^2}$$

$$pV = nRT$$

show that the mean kinetic energy of an individual (monatomic) gas particle of mass m is given by: [3]

$$\frac{1}{2}m\overline{c^2} = \frac{3}{2}kT$$

•••••		
***************************************	• • • • • • • • • • • • • • • • • • • •	
•••••		
•••••		
(b)	(i)	Calculate the temperature at which argon atoms have a rms speed of 630 m s <sup>-1</sup> . (Relative molecular mass of argon = 38.)
••••••		
•••••		


(ii)	Calculate the rms speed of argon atoms when this temperature is doubled.	[2

© WJEC CBAC Ltd.


Examiner only

3. A mass oscillates vertically on a spring as shown.

Examiner only



(a) The graph below shows the variation of acceleration with displacement of the mass on the spring.



(i) Explain how the graph verifies that the mass will perform simple harmonic motion. [2]


(ii) Use the graph to show that the frequency of oscillation of the mass on the spring is approximately 1 Hz. [3]

0.05

(iii)	The amplitude of oscillation of the mass on the spring is 2.8 cm. Writ (or calculate) the maximum acceleration of the mass.	e down [1]
(iv)	Calculate the maximum kinetic energy of the 0.20 kg mass.	[3]
(v)	If the mass is moving upwards at its maximum speed when $t = 0$ s, calculate time that the mass moves upwards with a speed of $0.100 \mathrm{ms^{-1}}$ .	the first [3]

Examiner only


(b) When damping is introduced the following graph of displacement against time is obtained. Explain how the principle of conservation of energy applies during the cycles shown. [4]



•••••	 	 	•••••••	 	 	 
•••••	 	 		 	 	 

4. An ideal monatomic gas is taken through the closed cycle ABCA as shown.

Examiner only



(a)	There are 15.6 mol of gas	s. Show that the temperatures of points <b>A</b> , <b>B</b> and	nd <b>C</b> are 309 K
	471 K and 160 K respective	ely.	[3]

(b) Calculate the **change** in internal energy for the three processes. [3]

(i) AB .....

(ii) BC

(iii) CA

	ulate the work done by the gas for each of the three processes.	
(i)	AB (one mark is available for the accuracy of your estimate)	[3]
•••••		
*******		
(ii)	BC	[1]
(iii)	CA	[1]
(i)	Calculate the heat supplied to the gas for process <b>AB</b> .	[2]
•••••		
(ii)	Process <b>AB</b> is in fact a very rapid compression. Explain why the answer to should be a low value.	(d)(i) [1]

5.	(a)	The	mass of the planet Mercury is $3.30 \times 10^{23}  \text{kg}$ and its radius is 2440 km.	
		(i)	Calculate the gravitational field strength on the surface of Mercury.	[2]
		**********		
		•••••		
		(ii)	Calculate the gravitational potential on the surface of Mercury.	[2]
		•••••		
		**********		
		(iii)	Explain briefly why the potential is negative.	[1]

A proveloc	jectile of mass 0.454 kg is fired upwards from Mercury's surface with an initial vertical ity of 1700 m s <sup>-1</sup> .
	Initial vertical velocity 1700 m s <sup>-1</sup>
(i)	Calculate the total energy of the projectile as it is being launched. [3]
(ii)	Use the principle of conservation of energy to calculate the maximum height that the projectile reaches (Mercury has no atmosphere so that air resistance is negligible). [3]


© WJEC CBAC Ltd. (1324-01) Turn over.

Examiner only

12

**6.** Three negative charges are arranged as shown.

(a) Draw arrows at P to represent the three electric fields due to the three charges. [2]



(b) Show that the resultant electric field at **P** is zero.

	(You may like to use the approximatio	$n \frac{1}{4\pi\varepsilon_0} = 9 \times 10^9 \text{F}^{-1}\text{m.}$	[5]
•••••			
•••••			
•••••			
••••••			
•••••			

© WJEC CBAC Ltd. (1324-01)


(c)	Calculate the electric potential at <b>P</b> .	3]	
••••			
(d)	An electron starts to accelerate from rest from point <b>P</b> (in a direction out of the plan of the paper). Calculate its speed when it arrives at another point where the potential –200 V.	ne is 3]	
•			
•••••			
•••••			

Examiner only

Turn over.

7. A star and planet travel in circular orbits about their mutual centre of mass.

Examiner only



(a)	Calculate the period of orbit of the planet-star system and convert your answer to (your answer will need to be accurate to 3 significant figures for full marks).	(3)
(b)	Calculate the radius of:  (i) the orbit of the star;	[3]
	(ii) the orbit of the planet.	

(c)	(i)	Show that the orbital speed of the star is around 600 m s <sup>-1</sup> . [2]	Examiner only
	(ii)	Calculate the Doppler shift for a wavelength of $1.875\mu m$ due to the orbital speed of the star (assuming that the system is viewed edge-on). [2]	:

## **END OF PAPER**