

3. A truck of mass of 300 kg moves along a straight horizontal road with a constant speed of 10 m s^{-1} . The resistance to motion of the truck has magnitude 120 N.

(a) Find the rate at which the engine of the truck is working. (2)

On another occasion the truck moves at a constant speed up a hill inclined at θ to the horizontal, where $\sin \theta = \frac{1}{14}$. The resistance to motion of the truck from non-gravitational forces remains of magnitude 120 N. The rate at which the engine works is the same as in part (a).

(b) Find the speed of the truck. (4)



Leave
blank

Question 4 continued

Handwriting practice lines consisting of 28 horizontal lines.



5.




Figure 2

A shop sign $ABCDEFG$ is modelled as a uniform lamina, as illustrated in Figure 2. $ABCD$ is a rectangle with $BC = 120$ cm and $DC = 90$ cm. The shape EFG is an isosceles triangle with $EG = 60$ cm and height 60 cm. The mid-point of AD and the mid-point of EG coincide.

- (a) Find the distance of the centre of mass of the sign from the side AD . (5)

The sign is freely suspended from A and hangs at rest.

- (b) Find the size of the angle between AB and the vertical. (4)



Leave blank

Question 5 continued

Lined writing area for the answer to Question 5.



6.




Figure 3

A child playing cricket on horizontal ground hits the ball towards a fence 10 m away. The ball moves in a vertical plane which is perpendicular to the fence. The ball just passes over the top of the fence, which is 2 m above the ground, as shown in Figure 3.

The ball is modelled as a particle projected with initial speed $u \text{ m s}^{-1}$ from point O on the ground at an angle α to the ground.

- (a) By writing down expressions for the horizontal and vertical distances, from O of the ball t seconds after it was hit, show that

$$2 = 10 \tan \alpha - \frac{50g}{u^2 \cos^2 \alpha} \tag{6}$$

Given that $\alpha = 45^\circ$,

- (b) find the speed of the ball as it passes over the fence. (6)



8. Particles A , B and C of masses $4m$, $3m$ and m respectively, lie at rest in a straight line on a smooth horizontal plane with B between A and C . Particles A and B are projected towards each other with speeds u m s⁻¹ and v m s⁻¹ respectively, and collide directly.

As a result of the collision, A is brought to rest and B rebounds with speed kv m s⁻¹. The coefficient of restitution between A and B is $\frac{3}{4}$.

- (a) Show that $u = 3v$. (6)

- (b) Find the value of k . (2)

Immediately after the collision between A and B , particle C is projected with speed $2v$ m s⁻¹ towards B so that B and C collide directly.

- (c) Show that there is no further collision between A and B . (4)



