

OXFORD CAMBRIDGE AND RSA EXAMINATIONS**Advanced Subsidiary General Certificate of Education
Advanced General Certificate of Education****MATHEMATICS****4728****Mechanics 1**Thursday **16 JUNE 2005** Afternoon 1 hour 30 minutesAdditional materials:
Answer booklet
Graph paper
List of Formulae (MF1)**TIME** 1 hour 30 minutes**INSTRUCTIONS TO CANDIDATES**

- Write your name, centre number and candidate number in the spaces provided on the answer booklet.
- Answer **all** the questions.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- The acceleration due to gravity is denoted by $g \text{ m s}^{-2}$. Unless otherwise instructed, when a numerical value is needed, use $g = 9.8$.
- You are permitted to use a graphical calculator in this paper.


INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- The total number of marks for this paper is 72.
- Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.
- **You are reminded of the need for clear presentation in your answers.**

This question paper consists of 4 printed pages.

2


1



A light inextensible string has its ends attached to two fixed points A and B . The point A is vertically above B . A smooth ring R of mass m kg is threaded on the string and is pulled by a force of magnitude 1.6 N acting upwards at 45° to the horizontal. The section AR of the string makes an angle of 30° with the downward vertical and the section BR is horizontal (see diagram). The ring is in equilibrium with the string taut.

- (i) Give a reason why the tension in the part AR of the string is the same as that in the part BR . [1]
- (ii) Show that the tension in the string is 0.754 N, correct to 3 significant figures. [3]
- (iii) Find the value of m . [3]

2



Particles A and B , of masses 0.2 kg and 0.3 kg respectively, are attached to the ends of a light inextensible string. Particle A is held at rest at a fixed point and B hangs vertically below A . Particle A is now released. As the particles fall the air resistance acting on A is 0.4 N and the air resistance acting on B is 0.25 N (see diagram). The downward acceleration of each of the particles is a m s^{-2} and the tension in the string is T N.

- (i) Write down two equations in a and T obtained by applying Newton's second law to A and to B . [4]
- (ii) Find the values of a and T . [3]

3

3 Two small spheres P and Q have masses 0.1 kg and 0.2 kg respectively. The spheres are moving directly towards each other on a horizontal plane and collide. Immediately before the collision P has speed 4 m s^{-1} and Q has speed 3 m s^{-1} . Immediately after the collision the spheres move away from each other, P with speed $u\text{ m s}^{-1}$ and Q with speed $(3.5 - u)\text{ m s}^{-1}$.

(i) Find the value of u . [4]

After the collision the spheres both move with deceleration of magnitude 5 m s^{-2} until they come to rest on the plane.


(ii) Find the distance PQ when both P and Q are at rest. [4]

4 A particle moves downwards on a smooth plane inclined at an angle α to the horizontal. The particle passes through the point P with speed $u\text{ m s}^{-1}$. The particle travels 2 m during the first 0.8 s after passing through P , then a further 6 m in the next 1.2 s . Find

(i) the value of u and the acceleration of the particle, [7]

(ii) the value of α in degrees. [2]

5



Two small rings A and B are attached to opposite ends of a light inextensible string. The rings are threaded on a rough wire which is fixed vertically. A is above B . A horizontal force is applied to a point P of the string. Both parts AP and BP of the string are taut. The system is in equilibrium with angle $BAP = \alpha$ and angle $ABP = \beta$ (see diagram). The weight of A is 2 N and the tensions in the parts AP and BP of the string are 7 N and $T\text{ N}$ respectively. It is given that $\cos \alpha = 0.28$ and $\sin \alpha = 0.96$, and that A is in limiting equilibrium.

(i) Find the coefficient of friction between the wire and the ring A . [7]


(ii) By considering the forces acting at P , show that $T \cos \beta = 1.96$. [2]

(iii) Given that there is no frictional force acting on B , find the mass of B . [3]

4

- 6 A particle of mass 0.04 kg is acted on by a force of magnitude $P \text{ N}$ in a direction at an angle α to the upward vertical.
- (i) The resultant of the weight of the particle and the force applied to the particle acts horizontally. Given that $\alpha = 20^\circ$ find
- the value of P , [3]
 - the magnitude of the resultant, [2]
 - the magnitude of the acceleration of the particle. [2]
- (ii) It is given instead that $P = 0.08$ and $\alpha = 90^\circ$. Find the magnitude and direction of the resultant force on the particle. [5]

7



A car P starts from rest and travels along a straight road for 600 s . The (t, v) graph for the journey is shown in the diagram. This graph consists of three straight line segments. Find

- the distance travelled by P , [3]
- the deceleration of P during the interval $500 < t < 600$. [2]

Another car Q starts from rest at the same instant as P and travels in the same direction along the same road for 600 s . At time $t \text{ s}$ after starting the velocity of Q is $(600t^2 - t^3) \times 10^{-6} \text{ m s}^{-1}$.

- Find an expression in terms of t for the acceleration of Q . [2]
- Find how much less Q 's deceleration is than P 's when $t = 550$. [2]
- Show that Q has its maximum velocity when $t = 400$. [2]
- Find how much further Q has travelled than P when $t = 400$. [6]