

**FP2 Paper \*adapted 2008**

1. Solve the differential equation  $\frac{dy}{dx} - 3y = x$

to obtain  $y$  as a function of  $x$ . (Total 5 marks)

2. (a) Simplify the expression  $\frac{(x+3)(x+9)}{x-1} - (3x-5)$ , giving your answer in the form  $\frac{a(x+b)(x+c)}{x-1}$ , where  $a, b$  and  $c$  are integers. (4)

(b) Hence, or otherwise, solve the inequality  $\frac{(x+3)(x+9)}{x-1} > 3x-5$  (4)(Total 8 marks)


3. (a) Find the general solution of the differential equation  $3\frac{d^2y}{dx^2} - \frac{dy}{dx} - 2y = x^2$  (8)

(b) Find the particular solution for which, at  $x = 0, y = 2$  and  $\frac{dy}{dx} = 3$ . (6)(Total 14 marks)

4. The diagram above shows the curve  $C_1$  which has polar equation  $r = a(3 + 2 \cos \theta), 0 \leq \theta < 2\pi$  and the circle  $C_2$  with equation  $r = 4a, 0 \leq \theta < 2\pi$ , where  $a$  is a positive constant.

(a) Find, in terms of  $a$ , the polar coordinates of the points where the curve  $C_1$  meets the circle  $C_2$ . (4)

The regions enclosed by the curves  $C_1$  and  $C_2$  overlap and this common region  $R$  is shaded in the figure.



(b) Find, in terms of  $a$ , an exact expression for the area of the region  $R$ . (8)

(c) In a single diagram, copy the two curves in the diagram above and also sketch the curve  $C_3$  with polar equation  $r = 2a \cos \theta, 0 \leq \theta < 2\pi$  Show clearly the coordinates of the points of intersection of  $C_1, C_2$  and  $C_3$  with the initial line,  $\theta = 0$ . (3)(Total 15 marks)

5. (a) Find, in terms of  $k$ , the general solution of the differential equation

$$\frac{d^2x}{dt^2} + 4\frac{dx}{dt} + 3x = kt + 5, \text{ where } k \text{ is a constant and } t > 0. (7)$$

For large values of  $t$ , this general solution may be approximated by a linear function.

(b) Given that  $k = 6$ , find the equation of this linear function. (2)(Total 9 marks)

6. (a) Find, in the simplest surd form where appropriate, the exact values of  $x$  for which

$$\frac{x}{2} + 3 = \left| \frac{4}{x} \right|. \quad (5)$$

- (b) Sketch, on the same axes, the line with equation  $y = \frac{x}{2} + 3$  and the graph of

$$y = \left| \frac{4}{x} \right|, \quad x \neq 0. \quad (3)$$

- (c) Find the set of values of  $x$  for which  $\frac{x}{2} + 3 > \left| \frac{4}{x} \right|$ . (2)(Total 10 marks)

7. (a) Show that the substitution  $y = vx$  transforms the differential equation

$$\frac{dy}{dx} = \frac{x}{y} + \frac{3y}{x}, \quad x > 0, \quad y > 0 \quad (I)$$

into the differential equation  $x \frac{dv}{dx} = 2v + \frac{1}{v}$ . (II) (3)

- (b) By solving differential equation (II), find a general solution of differential equation (I) in the form  $y = f(x)$ . (7)

Given that  $y = 3$  at  $x = 1$ , (c) find the particular solution of differential equation (I). (2)

8. The curve  $C$  shown in the diagram above has polar equation

$$r = 4(1 - \cos \theta), \quad 0 \leq \theta \leq \frac{\pi}{2}.$$

At the point  $P$  on  $C$ , the tangent to  $C$  is parallel to the line  $\theta = \frac{\pi}{2}$ .

- (a) Show that  $P$  has polar coordinates  $\left(2, \frac{\pi}{3}\right)$ . (5)

The curve  $C$  meets the line  $\theta = \frac{\pi}{2}$  at the point  $A$ . The tangent to  $C$  at the initial line at the point  $N$ . The finite region  $R$ , shown shaded in the diagram above, is bounded by the initial line, the line  $\theta = \frac{\pi}{2}$ , the arc  $AP$  of  $C$  and the line  $PN$ .


- (b) Calculate the exact area of  $R$ .

(8)

- 10.

$$(x^2 + 1) \frac{d^2y}{dx^2} = 2y^2 + (1 - 2x) \frac{dy}{dx} \quad (I)$$

- (a) By differentiating equation (I) with respect to  $x$ , show that



$$(x^2 + 1)\frac{d^3y}{dx^3} = (1 - 4x)\frac{d^2y}{dx^2} + (4y - 2)\frac{dy}{dx}. \quad (3)$$

Given that  $y = 1$  and  $\frac{dy}{dx} = 1$  at  $x = 0$ ,

(b) find the series solution for  $y$ , in ascending powers of  $x$ , up to and including the term in  $x_3$ .(4)

(c) Use your series to estimate the value of  $y$  at  $x = -0.5$ , giving your answer to two decimal places.(1)

**11.** The point  $P$  represents a complex number  $z$  on an Argand diagram such that

$$|z - 3| = 2|z|.$$

(a) Show that, as  $z$  varies, the locus of  $P$  is a circle, and give the coordinates of the centre and the radius of the circle.(5)

The point  $Q$  represents a complex number  $z$  on an Argand diagram such that

$$|z + 3| = |z - i\sqrt{3}|.$$

(b) Sketch, on the same Argand diagram, the locus of  $P$  and the locus of  $Q$  as  $z$  varies.(5)

(c) On your diagram shade the region which satisfies

$$|z - 3| \geq 2|z| \text{ and } |z + 3| \geq |z - i\sqrt{3}|. \quad (2)$$

**12.** De Moivre's theorem states that  $(\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta$  for  $n \in \mathfrak{R}$

(a) Use induction to prove de Moivre's theorem for  $n \in \mathbb{Z}^+$ . (5)

(b) Show that  $\cos 5\theta = 16\cos^5\theta - 20\cos^3\theta + 5\cos\theta$  (5)

(c) Hence show that  $2\cos\frac{\pi}{10}$  is a root of the equation

$$x^4 - 5x^2 + 5 = 0$$

(3)