



# **GCE MARKING SCHEME**


**CHEMISTRY  
AS/Advanced**

**SUMMER 2011**


### CHEMISTRY - CH5

Q.1 (a) Reacts with both acids **and** bases / behaves as an acid **and** a base. [1]

(b) Chromium atom, Cr [1]



Chromium(III) ion, Cr<sup>3+</sup> [1]



(c) (i) Orange → yellow [1]


(ii) Cr +6 (1) in both reactant and product - do not accept 6+  
no change in oxidation states so not a redox reaction. (1) [2]

(d) Add sodium hydroxide solution dropwise until there is an **excess** / small volume at a time until **excess**. [1]

White precipitate forms with Mg but doesn't dissolve again (therefore not amphoteric). [1]

White precipitate forms with Al then dissolves in excess NaOH (therefore amphoteric). [1]

(e) (i)



(co-ordinate bonds can be shown as lines but are incorrect if shown as arrows from Al to Cl)

Al is electron deficient - do not accept 'AlCl<sub>3</sub> is electron deficient' [1]

Cl has lone pairs [1]

(ii) Tetrahedral (1); four electron pairs and no lone pairs/ four bonding pairs (1) [2]

**Total [14]**

- Q.2** (a) (i)  $\text{H}_2 + \frac{1}{2} \text{O}_2 \rightarrow \text{H}_2\text{O}$  [1]
- (ii) Higher efficiency / no carbon dioxide emissions / water only / no greenhouse gases / can use renewable energy resources. [1]  
Too vague - do not accept clean / no polluting gases / no global warming.
- (iii) A = Salt bridge (1)  
B = High resistance voltmeter /potentiometer (1)  
C = Platinum electrodes (1) [3]
- (b) (i)  $\Delta H = 2 \times \Delta H (\text{H}_2\text{O}) + \Delta H (\text{CO}_2) - \Delta H(\text{CH}_3\text{OH})$   
 $= 2 \times -286 + (-394) - (-239)$  (1)  
 $= -727 \text{ kJ mol}^{-1}$  (1) [2]
- (ii) Entropy of (methanol) gas is higher than liquid (1)  
So entropy change will be more negative (1) [2]
- (iii)  $\Delta G = -727000 - (298 \times -81) = -703 \text{ kJ mol}^{-1}$  (1) *Allow ECF*  
Negative  $\Delta G$  means reaction is feasible. (1) [2]

**Total [11]**

**Q.3** (a) Any 2 for (1) each from:

- Measure pressure (at constant volume) over time
- Measure volume (at constant pressure) over time
- Colorimetry/ measuring colour over time

*1 mark allowed if time not mentioned*

[2]

(b) (i) When concentration doubles, rate doubles (1)

Therefore first order or rate is proportional to concentration (*must give reason to obtain this mark*) (1) [2]

Credit possible by alternative methods:

Calculate k for each and show that all values are the same;  
Calculate k for one concentration and use to calculate other values.

(ii)  $k = \text{Rate} \div [\text{N}_2\text{O}_5]$  e.g.  $k = 3.00 \times 10^{-5} \div 4.00 \times 10^{-3}$  (1)  
 $= 7.50 \times 10^{-3}$  (1) *must be 3 significant figures*

Units =  $\text{s}^{-1}$  (1)

[3]

(iii) Rate determining step must have one  $\text{N}_2\text{O}_5$  molecule as reactant. (1)  
Mechanism A matches this rate equation (1) *need reason to get this mark*

Accept reverse argument.

[2]

(c) (i)  $K_p = \frac{P_{\text{N}_2\text{O}_4}}{P_{\text{NO}_2}^2}$

[1]

(ii) Increasing temp shifts equilibrium to left / favours endothermic reaction (1) so value of  $K_p$  is decreased. (1)

[2]

(iii)  $P_{\text{N}_2\text{O}_4} = 9.5 \times 10^3 \text{ Pa}$  (1)  
 $K_p = 9.5 \times 10^3 \div (2.81 \times 10^5)^2 = 1.20 \times 10^{-7}$  (1) *Allow ECF*  
Units =  $\text{Pa}^{-1}$  (1) Mark consequentially on answer to (c)(i)

[3]

**Total [15]**

- Q.4** (a) (i) Transition metals have partially filled *d*-orbitals (in atom or ion) [1]
- (ii) Iron and copper have partially filled *d*-orbitals in their **ions**, zinc does not [1]
- (b) *QWC: organisation of information clearly and coherently; use of specialist vocabulary where appropriate.* (1)  
*QWC: selection of a form and style of writing appropriate to purpose and to complexity of subject matter.* (1) [2]
- Ligands cause *d*-orbitals to split
  - into 2 higher energy/ 3 lower energy
  - Electrons absorb light (frequencies) to move to higher energy level
  - Colour seen is colour transmitted/reflected/not absorbed
  - Copper(II) complexes absorb red /orange/yellow/all colours except blue.  
 [MAX 4 marks from points above]
  - Different ligands cause different splittings / different  $\Delta E$ .
  - Copper(I) ion has full *d*-orbitals.
  - So electrons cannot move to upper energy levels.
- [OVERALL MAX 6]
- (c) (i)  $\text{Fe}_2\text{O}_3 + 3\text{CO} \rightarrow 2\text{Fe} + 3\text{CO}_2$  [1]
- (ii) Fe oxidation state goes from +3 to 0 (1) / so it is reduced (1)  
 OR C (not CO) oxidation state goes from +2 to +4 (1)/ so it is being oxidised. (1) *Allow ECF* [2]
- (iii) Stable oxidation state of (C is +4 whilst) Pb is +2 (1)  
 Due to inert pair effect becoming more significant down the group. (1) [2]
- (d) (i)  $6\text{Fe}^{2+} + \text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ \rightarrow 6\text{Fe}^{3+} + 2\text{Cr}^{3+} + 7\text{H}_2\text{O}$  [1]
- (ii) Moles  $\text{Cr}_2\text{O}_7^{2-} = 23.80 \times 0.0200 \div 1000 = 4.76 \times 10^{-4}$  moles (1)  
 Moles  $\text{Fe}^{2+} = 4.76 \times 10^{-4} \times 6 = 2.86 \times 10^{-3}$  moles (1) [2]
- (iii) Mass Fe in sample =  $2.86 \times 10^{-3} \times 10 \times 55.8 = 1.59$  g (1)  
 Percentage Iron =  $1.59 \div 1.870 \times 100 = 85.2\%$  (1) [2]

**Total [20]**

- Q.5** (a) Named compound examples, need both name and use for (1)
- Sodium chlorate(I) = bleach
  - Sodium chlorate(V) = weedkiller
  - PVC = windows frames/guttering/pipes/insulation for electrical wires
  - Dichloromethane – solvent / paintstripper
  - CFCs = refrigerants / aerosol propellants
  - Aldrin / Dieldrin / DDT = Insecticides [1]
- (b) (i)  $\text{Cl}_2 + 2\text{Br}^- \rightarrow \text{Br}_2 + 2\text{Cl}^-$  [1]
- (ii) • Emf for reaction of bromide with chlorine is +0.27 V /  $E^\ominus$  for chlorine is more positive than for bromine. (1)
- Emf for reaction of bromide with iodine is -0.55 V /  $E^\ominus$  for iodine is less positive than for bromine. (1)
- Reactions are only feasible if Emf is positive / if  $E^\ominus$  for oxidising agent is more positive than for species being oxidised. (1) [3]
- (c) (i) White precipitate with (sodium) chloride, yellow precipitate with (sodium) iodide [1]
- (ii) *QWC: legibility of text; accuracy of spelling, punctuation and grammar; clarity of meaning. (1)* [1]
- NaCl: Steamy gas / bubbles (1)
  - NaI: Steamy gas / smell of rotten eggs / purple vapour or brown solution or black solid / yellow solid (1 mark for 2 observations)
  - NaCl:  $\text{NaHSO}_4$ , HCl / NaI:  $\text{NaHSO}_4$  / HI /  $\text{I}_2$  /  $\text{H}_2\text{S}$  /  $\text{SO}_2$  / S /  $\text{H}_2\text{O}$  (1 mark for 2 products; 2 marks for 4 products)
  - Iodide is easier to oxidise / iodide is a stronger reducing agent than chloride (1) [5]
- (d) (i) (Almost) completely dissociates to release  $\text{H}^+$ . [1]
- (ii) 
$$K_a = \frac{[\text{H}^+][\text{OCl}^-]}{[\text{HOCl}]}$$
 [1]
- (iii)  $[\text{H}^+] = 10^{-\text{pH}}$  OR  $\text{pH} = -\log [\text{H}^+]$  (1)
- $[\text{H}^+] = 5.88 \times 10^{-5} \text{ mol dm}^{-3}$  (1) [2]
- (iv) 
$$K_a = \frac{[\text{H}^+][\text{OCl}^-]}{[\text{HOCl}]} = \frac{(5.88 \times 10^{-5})^2}{0.100}$$
 (1) =  $3.47 \times 10^{-8} \text{ (mol dm}^{-3})$  (1)
- (allow consequential answers) [2]
- (v) pH above 7 (up to 10) (1)
- $\text{OCl}^-$  in equilibrium with  $\text{HOCl}$  /  $\text{OCl}^-$  will remove  $\text{H}^+$  from solution (1) [2]

**Total [20]**