



Friday 25 January 2013 – Afternoon

## AS GCE MATHEMATICS

4728/01 Mechanics 1

### QUESTION PAPER

Candidates answer on the Printed Answer Book.

**OCR supplied materials:**

- Printed Answer Book 4728/01
- List of Formulae (MF1)

**Other materials required:**

- Scientific or graphical calculator

**Duration:** 1 hour 30 minutes



### INSTRUCTIONS TO CANDIDATES

These instructions are the same on the Printed Answer Book and the Question Paper.

- The Question Paper will be found in the centre of the Printed Answer Book.
- Write your name, centre number and candidate number in the spaces provided on the Printed Answer Book. Please write clearly and in capital letters.
- **Write your answer to each question in the space provided in the Printed Answer Book.** Additional paper may be used if necessary but you must clearly show your candidate number, centre number and question number(s).
- Use black ink. HB pencil may be used for graphs and diagrams only.
- Answer **all** the questions.
- Read each question carefully. Make sure you know what you have to do before starting your answer.
- Do **not** write in the bar codes.
- You are permitted to use a scientific or graphical calculator in this paper.
- Give non-exact numerical answers correct to 3 significant figures unless a different degree of accuracy is specified in the question or is clearly appropriate.
- The acceleration due to gravity is denoted by  $g \text{ ms}^{-2}$ . Unless otherwise instructed, when a numerical value is needed, use  $g = 9.8$ .

### INFORMATION FOR CANDIDATES

This information is the same on the Printed Answer Book and the Question Paper.

- The number of marks is given in brackets [ ] at the end of each question or part question on the Question Paper.
- **You are reminded of the need for clear presentation in your answers.**
- The total number of marks for this paper is **72**.
- The Printed Answer Book consists of **16** pages. The Question Paper consists of **4** pages. Any blank pages are indicated.

### INSTRUCTION TO EXAMS OFFICER/INVIGILATOR

- Do not send this Question Paper for marking; it should be retained in the centre or recycled. Please contact OCR Copyright should you wish to re-use this document.

## 2


1 Three horizontal forces, acting at a single point, have magnitudes 12 N, 14 N and 5 N and act along bearings  $000^\circ$ ,  $090^\circ$  and  $270^\circ$  respectively. Find the magnitude and bearing of their resultant. [5]

2 A particle  $P$  moves in a straight line. The displacement of  $P$  from a fixed point on the line is  $(t^4 - 2t^3 + 5)$  m, where  $t$  is the time in seconds. Show that, when  $t = 1.5$ ,

(i)  $P$  is at instantaneous rest, [3]

(ii) the acceleration of  $P$  is  $9 \text{ m s}^{-2}$ . [3]

## 3



A particle  $P$  of mass  $0.25 \text{ kg}$  moves upwards with constant speed  $u \text{ m s}^{-1}$  along a line of greatest slope on a smooth plane inclined at  $30^\circ$  to the horizontal. The pulling force acting on  $P$  has magnitude  $T \text{ N}$  and acts at an angle of  $20^\circ$  to the line of greatest slope (see diagram). Calculate

(i) the value of  $T$ , [3]

(ii) the magnitude of the contact force exerted on  $P$  by the plane. [3]

The pulling force  $T \text{ N}$  acting on  $P$  is suddenly removed, and  $P$  comes to instantaneous rest  $0.4 \text{ s}$  later.

(iii) Calculate  $u$ . [4]

4 The acceleration of a particle  $P$  moving in a straight line is  $(t^2 - 9t + 18) \text{ m s}^{-2}$ , where  $t$  is the time in seconds.


(i) Find the values of  $t$  for which the acceleration is zero. [2]

(ii) It is given that when  $t = 3$  the velocity of  $P$  is  $9 \text{ m s}^{-1}$ . Find the velocity of  $P$  when  $t = 0$ . [4]

(iii) Show that the direction of motion of  $P$  changes before  $t = 1$ . [2]

3

5



A small smooth pulley is suspended from a fixed point by a light chain. A light inextensible string passes over the pulley. Particles  $P$  and  $Q$ , of masses  $0.3\text{ kg}$  and  $m\text{ kg}$  respectively, are attached to the opposite ends of the string. The particles are released from rest at a height of  $0.2\text{ m}$  above horizontal ground with the string taut; the portions of the string not in contact with the pulley are vertical (see diagram).  $P$  strikes the ground with speed  $1.4\text{ m s}^{-1}$ . Subsequently  $P$  remains on the ground, and  $Q$  does not reach the pulley.

- (i) Calculate the acceleration of  $P$  while it is in motion and the corresponding tension in the string. [4]
- (ii) Find the value of  $m$ . [3]
- (iii) Calculate the greatest height of  $Q$  above the ground. [4]
- (iv) It is given that the mass of the pulley is  $0.5\text{ kg}$ . State the magnitude of the tension in the chain which supports the pulley
- (a) when  $P$  is in motion, [2]
- (b) when  $P$  is at rest on the ground and  $Q$  is moving upwards. [1]
- 6 Particle  $P$  of mass  $0.3\text{ kg}$  and particle  $Q$  of mass  $0.2\text{ kg}$  are  $3.6\text{ m}$  apart on a smooth horizontal surface.  $P$  and  $Q$  are simultaneously projected directly towards each other along a straight line. Before the particles collide  $P$  has speed  $4\text{ m s}^{-1}$  and  $Q$  has speed  $5\text{ m s}^{-1}$ .
- (i) Given that the particles coalesce in the collision, calculate their common speed after they collide. [3]
- (ii) It is given instead that one particle is at rest immediately after the collision.
- (a) State which particle is in motion after the collision and find the speed of this particle. [4]
- (b) Find the time taken after the collision for the moving particle to return to its initial position. [4]
- (c) On a single diagram sketch the  $(t, v)$  graphs for the two particles, with  $t = 0$  as the instant of their initial projection. [4]

## 4

- 7  $A$  and  $B$  are two points on a line of greatest slope of a plane inclined at  $45^\circ$  to the horizontal and  $AB = 2$  m. A particle  $P$  of mass  $0.4$  kg is projected from  $A$  towards  $B$  with speed  $5 \text{ m s}^{-1}$ . The coefficient of friction between the plane and  $P$  is  $0.2$ .
- (i) Given that the level of  $A$  is above the level of  $B$ , calculate the speed of  $P$  when it passes through the point  $B$ , and the time taken to travel from  $A$  to  $B$ . [7]
- (ii) Given instead that the level of  $A$  is below the level of  $B$ ,
- (a) show that  $P$  does not reach  $B$ , [3]
- (b) calculate the difference in the momentum of  $P$  for the two occasions when it is at  $A$ . [4]

**Copyright Information**

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website ([www.ocr.org.uk](http://www.ocr.org.uk)) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.