

Centre No.							Paper Reference					Surname	Initial(s)		
Candidate No.							<b>6</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>/</b>	<b>0</b>	<b>1</b>	Signature	

Paper Reference(s)

**6678/01**

# Edexcel GCE

## Mechanics M2

### Advanced/Advanced Subsidiary

Thursday 31 May 2012 – Morning

Time: 1 hour 30 minutes

Examiner's use only

--	--	--

Team Leader's use only

--	--	--

Question Number	Leave Blank
1	
2	
3	
4	
5	
6	
7	
Total	

Materials required for examination

Mathematical Formulae (Pink)

Items included with question papers

Nil

**Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation or symbolic differentiation/integration, or have retrievable mathematical formulae stored in them.**

**Instructions to Candidates**

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper. Answer ALL the questions. You must write your answer to each question in the space following the question. Whenever a numerical value of *g* is required, take  $g = 9.8 \text{ m s}^{-2}$ . When a calculator is used, the answer should be given to an appropriate degree of accuracy.

**Information for Candidates**

A booklet 'Mathematical Formulae and Statistical Tables' is provided. Full marks may be obtained for answers to ALL questions. The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2). There are 7 questions in this question paper. The total mark for this paper is 75. There are 24 pages in this question paper. Any blank pages are indicated.

**Advice to Candidates**

You must ensure that your answers to parts of questions are clearly labelled. You must show sufficient working to make your methods clear to the examiner. Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Pearson Education Ltd copyright policy.  
©2012 Pearson Education Ltd.



**Turn over**



2. A particle  $P$  of mass  $3m$  is moving with speed  $2u$  in a straight line on a smooth horizontal plane. The particle  $P$  collides directly with a particle  $Q$  of mass  $4m$  moving on the plane with speed  $u$  in the opposite direction to  $P$ . The coefficient of restitution between  $P$  and  $Q$  is  $e$ .

(a) Find the speed of  $Q$  immediately after the collision. (6)

Given that the direction of motion of  $P$  is reversed by the collision,

(b) find the range of possible values of  $e$ . (5)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---


---







4.



**Figure 2**

A uniform circular disc has centre  $O$  and radius  $4a$ . The lines  $PQ$  and  $ST$  are perpendicular diameters of the disc. A circular hole of radius  $2a$  is made in the disc, with the centre of the hole at the point  $R$  on  $OP$  where  $OR = 2a$ , to form the lamina  $L$ , shown shaded in Figure 2.

(a) Show that the distance of the centre of mass of  $L$  from  $P$  is  $\frac{14a}{3}$ . (4)

The mass of  $L$  is  $m$  and a particle of mass  $km$  is now fixed to  $L$  at the point  $P$ . The system is now suspended from the point  $S$  and hangs freely in equilibrium. The diameter  $ST$  makes an angle  $\alpha$  with the downward vertical through  $S$ , where  $\tan \alpha = \frac{5}{6}$ .

(b) Find the value of  $k$ . (5)

---

---

---

---

---

---

---

---

---


---

---

---



5.



**Figure 3**

A small ball  $B$  of mass  $0.25 \text{ kg}$  is moving in a straight line with speed  $30 \text{ m s}^{-1}$  on a smooth horizontal plane when it is given an impulse. The impulse has magnitude  $12.5 \text{ N s}$  and is applied in a horizontal direction making an angle of  $(90^\circ + \alpha)$ , where  $\tan \alpha = \frac{3}{4}$ , with the initial direction of motion of the ball, as shown in Figure 3.

- (i) Find the speed of  $B$  immediately after the impulse is applied.
- (ii) Find the direction of motion of  $B$  immediately after the impulse is applied. (6)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---












7.



**Figure 4**

A small stone is projected from a point  $O$  at the top of a vertical cliff  $OA$ . The point  $O$  is 52.5 m above the sea. The stone rises to a maximum height of 10 m above the level of  $O$  before hitting the sea at the point  $B$ , where  $AB = 50$  m, as shown in Figure 4. The stone is modelled as a particle moving freely under gravity.

- (a) Show that the vertical component of the velocity of projection of the stone is  $14 \text{ m s}^{-1}$ . **(3)**
- (b) Find the speed of projection. **(9)**
- (c) Find the time after projection when the stone is moving parallel to  $OB$ . **(5)**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



