



General Certificate of Education  
Advanced Level Examination  
January 2013

## Mathematics

## MFP3

### Unit Further Pure 3

Friday 25 January 2013 1.30 pm to 3.00 pm

**For this paper you must have:**

- the blue AQA booklet of formulae and statistical tables.
- You may use a graphics calculator.

**Time allowed**

- 1 hour 30 minutes

**Instructions**

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- Write the question part reference (eg (a), (b)(i) etc) in the left-hand margin.
- You must answer each question in the space provided for that question. If you require extra space, use an AQA supplementary answer book; do **not** use the space provided for a different question.
- Do not write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.

**Information**

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 75.

**Advice**

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.

## 2

- 1 It is given that  $y(x)$  satisfies the differential equation

$$\frac{dy}{dx} = f(x, y)$$

where  $f(x, y) = \sqrt{2x + y}$

and  $y(3) = 5$

- (a) Use the Euler formula

$$y_{r+1} = y_r + hf(x_r, y_r)$$

with  $h = 0.2$ , to obtain an approximation to  $y(3.2)$ , giving your answer to four decimal places. (3 marks)

- (b) Use the formula

$$y_{r+1} = y_{r-1} + 2hf(x_r, y_r)$$

with your answer to part (a), to obtain an approximation to  $y(3.4)$ , giving your answer to three decimal places. (3 marks)

---

- 2 (a) Write down the expansion of  $e^{3x}$  in ascending powers of  $x$  up to and including the term in  $x^2$ . (1 mark)

- (b) Hence, or otherwise, find the term in  $x^2$  in the expansion, in ascending powers of  $x$ , of  $e^{3x}(1 + 2x)^{-\frac{3}{2}}$ . (4 marks)
- 

- 3 It is given that the general solution of the differential equation

$$\frac{d^2y}{dx^2} - 2\frac{dy}{dx} + y = 0$$

is  $y = e^x(Ax + B)$ . Hence find the general solution of the differential equation

$$\frac{d^2y}{dx^2} - 2\frac{dy}{dx} + y = 6e^x \quad (5 \text{ marks})$$



**4 (a)** Explain why  $\int_0^1 x^4 \ln x \, dx$  is an improper integral. (1 mark)

**(b)** Evaluate  $\int_0^1 x^4 \ln x \, dx$ , showing the limiting process used. (6 marks)

---

**5 (a)** Show that  $\tan x$  is an integrating factor for the differential equation

$$\frac{dy}{dx} + \frac{\sec^2 x}{\tan x} y = \tan x \quad (2 \text{ marks})$$

**(b)** Hence solve this differential equation, given that  $y = 3$  when  $x = \frac{\pi}{4}$ . (6 marks)

---

**6 (a)** It is given that  $y = \ln(e^{3x} \cos x)$ .

**(i)** Show that  $\frac{dy}{dx} = 3 - \tan x$ . (3 marks)

**(ii)** Find  $\frac{d^4 y}{dx^4}$ . (3 marks)

**(b)** Hence use Maclaurin's theorem to show that the first three non-zero terms in the expansion, in ascending powers of  $x$ , of  $\ln(e^{3x} \cos x)$  are  $3x - \frac{1}{2}x^2 - \frac{1}{12}x^4$ . (3 marks)

**(c)** Write down the expansion of  $\ln(1 + px)$ , where  $p$  is a constant, in ascending powers of  $x$  up to and including the term in  $x^2$ . (1 mark)

**(d) (i)** Find the value of  $p$  for which  $\lim_{x \rightarrow 0} \left[ \frac{1}{x^2} \ln \left( \frac{e^{3x} \cos x}{1 + px} \right) \right]$  exists.

**(ii)** Hence find the value of  $\lim_{x \rightarrow 0} \left[ \frac{1}{x^2} \ln \left( \frac{e^{3x} \cos x}{1 + px} \right) \right]$  when  $p$  takes the value found in part **(d)(i)**. (4 marks)

Turn over ►



- 7 (a) Find the general solution of the differential equation

$$\frac{d^2y}{dt^2} - 6\frac{dy}{dt} + 10y = e^{2t}$$

giving your answer in the form  $y = f(t)$ . (6 marks)

- (b) Given that  $x = t^{\frac{1}{2}}$ ,  $x > 0$ ,  $t > 0$  and  $y$  is a function of  $x$ , show that

$$\frac{d^2y}{dx^2} = 4t \frac{d^2y}{dt^2} + 2 \frac{dy}{dt} \quad (5 \text{ marks})$$

- (c) Hence show that the substitution  $x = t^{\frac{1}{2}}$  transforms the differential equation

$$x \frac{d^2y}{dx^2} - (12x^2 + 1) \frac{dy}{dx} + 40x^3y = 4x^3e^{2x^2}$$

into


$$\frac{d^2y}{dt^2} - 6\frac{dy}{dt} + 10y = e^{2t} \quad (2 \text{ marks})$$

- (d) Hence **write down** the general solution of the differential equation

$$x \frac{d^2y}{dx^2} - (12x^2 + 1) \frac{dy}{dx} + 40x^3y = 4x^3e^{2x^2} \quad (1 \text{ mark})$$



- 8 The diagram shows a sketch of a curve.



The polar equation of the curve is

$$r = \sin 2\theta \sqrt{\left(2 + \frac{1}{2} \cos \theta\right)}, \quad 0 \leq \theta \leq \frac{\pi}{2}$$

The point  $P$  is the point of the curve at which  $\theta = \frac{\pi}{3}$ .

The perpendicular from  $P$  to the initial line meets the initial line at the point  $N$ .

- (a) (i) Find the exact value of  $r$  when  $\theta = \frac{\pi}{3}$ . (2 marks)
- (ii) Show that the polar equation of the line  $PN$  is  $r = \frac{3\sqrt{3}}{8} \sec \theta$ . (2 marks)
- (iii) Find the area of triangle  $ONP$  in the form  $\frac{k\sqrt{3}}{128}$ , where  $k$  is an integer. (2 marks)
- (b) (i) Using the substitution  $u = \sin \theta$ , or otherwise, find  $\int \sin^n \theta \cos \theta \, d\theta$ , where  $n \geq 2$ . (2 marks)
- (ii) Find the area of the shaded region bounded by the line  $OP$  and the arc  $OP$  of the curve. Give your answer in the form  $a\pi + b\sqrt{3} + c$ , where  $a$ ,  $b$  and  $c$  are constants. (8 marks)

