

Write your name here

Surname					Other names				
Centre Number					Candidate Number				
<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>					<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>				

Edexcel GCE

Chemistry
Advanced Subsidiary
Unit 3B: Chemistry Laboratory Skills I Alternative

Monday 11 January 2010 – Afternoon Time: 1 hour 15 minutes	Paper Reference 6CH07/01
--	------------------------------------

Candidates may use a calculator.

Total Marks

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided – *there may be more space than you need.*

Information

- The total mark for this paper is 50.
- The marks for **each** question are shown in brackets – *use this as a guide as to how much time to spend on each question.*
- You will be assessed on your ability to organise and present information, ideas, descriptions and arguments clearly and logically, including your use of grammar, punctuation and spelling.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Keep an eye on the time.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

N34104A

©2010 Edexcel Limited.

7/7/5/3/



edexcel 
 advancing learning, changing lives

Answer ALL the questions. Write your answers in the spaces provided.

1 Compound **A** is a white solid that contains one cation and one anion.

(a) A flame test is carried out on compound **A** by mixing the solid with concentrated hydrochloric acid and using a piece of wire to place some of the solution formed in a Bunsen flame. The flame is coloured yellow.

(i) Name a material from which the wire is made.

Suggest ONE reason why this material is used.

(2)

Name.....

Reason.....

(ii) Identify, by name or formula, the cation present in compound **A**.

(1)

(b) When aqueous silver nitrate solution is added to a solution of compound **A**, a cream precipitate forms. The cream precipitate dissolves in concentrated aqueous ammonia solution.

(i) **Name** the cream precipitate formed when aqueous silver nitrate is added to the solution of compound **A**.

(1)

(ii) Give the **formula** of the anion in compound **A**.

(1)

(iii) Describe what you would see if the cream precipitate in (b)(i) was left in sunlight.

(1)



(c) Concentrated sulfuric acid is added to compound A in a test tube. Steamy fumes are seen at the mouth of the test tube.

After a few minutes, the contents of the test tube turn brown. A gas is given off which is tested with a piece of filter paper soaked in a solution of aqueous acidified potassium dichromate(VI). The paper turns green.

(i) Identify, by name or formula, the steamy fumes formed initially. (1)

(ii) Describe a further test you could carry out to confirm the identity of the steamy fumes. Give the result of your test. (2)

Test

Result

(iii) Identify, by name or formula, the substance responsible for the brown colour in the test tube. (1)

(iv) Name the gas which turned the filter paper green. Suggest the type of reaction by which this gas was formed from sulfuric acid. (2)

Gas

Type of reaction

(Total for Question 1 = 12 marks)



2 This question is about two isomeric alcohols, **X** and **Y**, each with molar mass 60 g mol^{-1} .

A solution of potassium dichromate(VI) in dilute sulfuric acid is added to each alcohol. Both alcohols cause the same colour change of the mixture on heating.

(a) A colourless liquid, **B**, is distilled from the mixture containing alcohol **X**.

The liquid **B** forms a red precipitate when it is boiled with Benedict's or Fehling's solution.

Give the displayed formula of liquid **B**, and the name of alcohol **X**.

(2)

Liquid **B**

Name of alcohol **X**

(b) A colourless liquid, **C**, is distilled from the mixture containing alcohol **Y**.

C does **not** react when it is boiled with Benedict's or Fehling's solution.

Give the **names** of liquid **C** and alcohol **Y**.

(2)

Liquid **C**

Alcohol **Y**

(Total for Question 2 = 4 marks)



3 This question is about calcium hydroxide, $\text{Ca}(\text{OH})_2$.

The solubility of calcium hydroxide in water can be found by titrating a saturated solution of calcium hydroxide with hydrochloric acid of known concentration.

- (a) Describe how you would make a saturated solution of calcium hydroxide suitable for use in this titration. Do not describe the subsequent titration procedure.

(2)

.....

.....

.....

.....

- (b) 10.0 cm^3 portions of the saturated solution are placed in a conical flask and titrated with $0.0500 \text{ mol dm}^{-3}$ hydrochloric acid added from a burette.

- (i) Name the apparatus used to measure the 10.0 cm^3 portions.

(1)

.....

- (ii) Suggest a suitable indicator for this titration and state the colour change you would expect to see at the end-point.

(2)

Indicator.....

Colour change from to



(c) The following results were obtained:

Titration numbers	1	2	3
Final burette reading / cm^3	19.20	28.05	37.10
Initial burette reading / cm^3	10.00	19.20	28.15
Titre / cm^3	9.20	8.85	

(i) Fill in the third titre value in the table. (1)

(ii) Suggest why the first titre should be disregarded. (1)


(iii) Calculate the mean titre. (1)

..... cm^3

(iv) Calculate the number of moles of hydrochloric acid in the mean titre. (1)



(v) The equation for the reaction is



Calculate the number of moles of calcium hydroxide in a 10.0cm^3 portion of the saturated solution.

(1)

(vi) Calculate the concentration of calcium hydroxide in mol dm^{-3} .

(1)

(vii) Calculate the solubility of calcium hydroxide in g dm^{-3} .

(1)



(d) The standard enthalpy change for the reaction of calcium hydroxide with hydrochloric acid was found by reacting 0.0100 mol of solid calcium hydroxide with 50.0 cm³ of a 1.00 mol dm⁻³ solution of hydrochloric acid (an excess), in a polystyrene cup. The temperature rose from 21.2 °C to 26.7 °C.

(i) Calculate the energy, in joules, transferred in the reaction. Use the expression

Energy transferred = mass × specific heat capacity × temperature change

[Assume density of solution = 1.0 g cm⁻³,
specific heat capacity of solution = 4.18 J g⁻¹ °C⁻¹]

(1)

(ii) Calculate the standard enthalpy change, ΔH^\ominus , for the reaction. Include a sign and units in your answer.

(2)

(iii) Calculate the percentage error in the temperature change caused by an uncertainty of 0.1 °C in each thermometer reading.

(2)



(iv) The experiment was repeated using 50.0 cm^3 of a 1.00 mol dm^{-3} solution of nitric acid instead of the hydrochloric acid. Explain why the temperature change was the same in both experiments.

(1)

(v) The experiment was repeated again using 25 cm^3 of 2.00 mol dm^{-3} hydrochloric acid. Predict the temperature change in this experiment.

(1)

(vi) Which of the experiments in (iv) and (v) gave the least error in the temperature change? Justify your answer.


(1)

(Total for Question 3 = 20 marks)



N 3 4 1 0 4 A 0 9 1 6

- 4 An experiment to prepare a sample of 2-chloro-2-methylpropane uses the reaction of 2-methylpropan-2-ol with concentrated hydrochloric acid.



The steps of the experimental procedure are as follows

1. Place 0.20 mol of 2-methylpropan-2-ol and 70 cm³ of concentrated hydrochloric acid in a large conical flask.
2. Stopper and shake the flask at intervals, releasing any pressure after each shaking.
3. Separate the 2-chloro-2-methylpropane from the aqueous solution using a separating funnel.
4. To the 2-chloro-2-methylpropane in the separating funnel, add 20 cm³ of sodium hydrogencarbonate solution. Shake the separating funnel, carefully releasing carbon dioxide frequently.
5. Separate the 2-chloro-2-methylpropane and repeat the washing with sodium hydrogencarbonate solution until this washing step is no longer necessary.
6. Transfer the 2-chloro-2-methylpropane to a small conical flask and add a suitable drying agent.
7. Filter off the drying agent, collecting the 2-chloro-2-methylpropane into a distillation flask. Heat the flask, collecting the fraction that distils off between 50 °C and 52 °C.

Data

Property	2-methylpropan-2-ol	2-chloro-2-methylpropane
Density / g cm ⁻³	0.789	0.842
Molar mass / g mol ⁻¹	74.1	92.6
Boiling temperature / °C	82.4	50.8



(a) Calculate the volume of 2-methylpropan-2-ol used in the preparation.

(1)

(b) Draw a diagram of a separating funnel that could be used in step 3.
Label the 2-chloro-2-methylpropane layer.

(2)

(c) (i) Suggest why the product is washed with sodium hydrogencarbonate solution (step 4).

(1)

(ii) How would you know that no further washing with sodium hydrogencarbonate was necessary?

(1)



(d) Suggest a suitable drying agent to dry the 2-chloro-2-methylpropane (step 6).

(1)

(e) Draw a labelled diagram of the apparatus you would use to carry out the final distillation (step 7).

(4)



(f) A reaction produced 18.7 cm^3 of 2-chloro-2-methylpropane starting from 0.20 mol 2-methylpropan-2-ol. Calculate the percentage yield of the product of this reaction. (2)

(g) A suitable chemical test for the chlorine in a chloroalkane, such as 2-chloro-2-methylpropane, is to add the chloroalkane to a mixture of aqueous silver nitrate solution and ethanol.

(i) Suggest why ethanol is added to the mixture. (1)

.....

.....

(ii) Give the expected result of this test. (1)

.....

(Total for Question 4 = 14 marks)

TOTAL FOR PAPER = 50 MARKS



BLANK PAGE



BLANK PAGE



The Periodic Table of Elements

1	2	3	4	5	6	7	0 (8)										
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)										
6.9 Li lithium 3	9.0 Be beryllium 4	11 Na sodium 11	12 Mg magnesium 12	13 B boron 5	14 C carbon 6	15 N nitrogen 7	16 O oxygen 8	17 F fluorine 9	18 Ne neon 10								
19 K potassium 19	20 Ca calcium 20	21 Sc scandium 21	22 Ti titanium 22	23 V vanadium 23	24 Cr chromium 24	25 Mn manganese 25	26 Fe iron 26	27 Co cobalt 27	28 Ni nickel 28	29 Cu copper 29	30 Zn zinc 30	31 Ga gallium 31	32 Ge germanium 32	33 As arsenic 33	34 Se selenium 34	35 Br bromine 35	36 Kr krypton 36
37 Rb rubidium 37	38 Sr strontium 38	39 Y yttrium 39	40 Zr zirconium 40	41 Nb niobium 41	42 Mo molybdenum 42	43 Tc technetium [98]	44 Ru ruthenium 44	45 Rh rhodium 45	46 Pd palladium 46	47 Ag silver 47	48 Cd cadmium 48	49 In indium 49	50 Sn tin 50	51 Sb antimony 51	52 Te tellurium 52	53 I iodine 53	54 Xe xenon 54
55 Cs caesium 55	56 Ba barium 56	57 La* lanthanum 57	72 Hf hafnium 72	73 Ta tantalum 73	74 W tungsten 74	75 Re rhenium 75	76 Os osmium 76	77 Ir iridium 77	78 Pt platinum 78	79 Au gold 79	80 Hg mercury 80	81 Tl thallium 81	82 Pb lead 82	83 Bi bismuth 83	84 Po polonium [209]	85 At astatine [210]	86 Rn radon [222]
87 Fr francium 87	88 Ra radium 88	89 Ac* actinium 89	104 Rf rutherfordium [261]	105 Db dubnium [262]	106 Sg seaborgium [266]	107 Bh bohrium [264]	108 Hs hassium [277]	109 Mt meitnerium [268]	110 Ds darmstadtium [271]	111 Rg roentgenium [272]	Elements with atomic numbers 112-116 have been reported but not fully authenticated						

140 Ce cerium 58	141 Pr praseodymium 59	144 Nd neodymium 60	147 Pm promethium 61	150 Sm samarium 62	152 Eu europium 63	157 Gd gadolinium 64	159 Tb terbium 65	163 Dy dysprosium 66	165 Ho holmium 67	167 Er erbium 68	169 Tm thulium 69	173 Yb ytterbium 70	175 Lu lutetium 71
232 Th thorium 90	231 Pa protactinium 91	238 U uranium 92	237 Np neptunium 93	242 Pu plutonium 94	243 Am americium 95	247 Cm curium 96	245 Bk berkelium 97	251 Cf californium 98	254 Es einsteinium 99	253 Fm fermium 100	256 Md mendelevium 101	254 No nobelium 102	257 Lr lawrencium 103

* Lanthanide series
* Actinide series

