


# **Mark Scheme 4733 June 2007**

4733

Mark Scheme

June 2007

1	(i)	$\hat{\mu} = 4830.0/100 = 48.3$ $249509.16/100 - (\text{their } \bar{x}^2)$ $\times 100/99$ $= 163.84$	B1 M1 M1 A1	4	48.3 seen Biased estimate: 162.2016: can get B1M1M0 Multiply by $n/(n - 1)$ Answer, 164 or 163.8 or 163.84
	(ii)	No, Central Limit theorem applies, so can assume distribution is normal	B2	2	“No” with statement showing CLT is understood (though CLT does not need to be mentioned) [SR: No with reason that is not wrong: B1]
2	$B(130, 1/40)$ $\approx \text{Po}(3.25)$ $e^{-\lambda} \frac{\lambda^3}{4!}$ $= 0.180$	B1 M1 A1 $\sqrt{}$ M1 A1	5	$B(130, 1/40)$ stated or implied Poisson, <i>or</i> correct N on their $B(n, p)$ Parameter their $np$ , <i>or</i> correct parameter(s) $\sqrt{}$ Correct formula, or interpolation Answer, 0.18 or a.r.t. 0.180 [SR: N(3.25, 3.17) or N(3.25, 3.25): B1M1A1]	
3	(i)	Binomial	B1	1	Binomial stated or implied
	(ii)	Each element equally likely Choices independent	B1 B1	2	All elements, or selections, equally likely stated Choices independent [not just “independent”] [can get B2 even if (i) is wrong]
4	(i)	Two of: Distribution symmetric No substantial truncation Unimodal/Increasingly unlikely further from $\mu$ , etc	B1 B1	2	One property Another definitely different property Don’t give both marks for just these two “Bell-shaped”: B1 only unless “no truncation”
	(ii)	Variance $8^2/20$ $z = \frac{47.0 - 50.0}{\sqrt{8^2/20}} = -1.677$ $\Phi(1.677) = 0.9532$	M1 A1 A1 A1	4	Standardise, allow cc, don’t need $n$ Denominator ( $8$ or $8^2$ or $\sqrt{8}$ ) $\div$ ( $20$ or $\sqrt{20}$ or $20^2$ ) $z$ -value, a.r.t. $-1.68$ or $+1.68$ Answer, a.r.t. 0.953
5	(i)	$H_1: \lambda > 2.5$ or 15	B1	1	$\lambda > 2.5$ or 15, allow $\mu$ , don’t need “ $H_1$ ”
	(ii)	Use parameter 15 $P(> 23)$  $1 - 0.9805 = 0.0195$ or 1.95%	M1 M1  A1	3	$\lambda = 15$ used [N(15, 15) gets this mark only] Find $P(> 23$ or $\geq 23)$ , final answer $< 0.5$ eg 0.0327 or 0.0122 Answer, 1.95% or 2% or 0.0195 or 0.02 [SR: 2-tailed, 3.9% gets 3/3 here]
	(iii)	$P(\leq 23   \lambda = 17) = 0.9367$ $P(\leq 23   \lambda = 18) = 0.8989$ Parameter = 17  $\lambda = 17/6$ or 2.83	M1  A1  M1	3	One of these, or their complement: .9367, .8989, 0.9047, 0.8551, .9317, .8933, .9907, .9805 Parameter 17 [17.1076], needs $P(\leq 23)$ , cwo [SR: if insufficient evidence can give B1 for 17] Their parameter $\div 6$ [2.85] [SR: Solve $(23.5 - \lambda)/\sqrt{\lambda} = 1.282$ M1; 18.05 A0]
6	(i)	$H_0: p = 0.19, H_1: p < 0.19$ where $p$ is population proportion $0.81^{20} + 20 \times 0.81^{19} \times 0.19$ $= 0.0841$ Compare 0.1	B2 M1 A1 A1 B1		Correct, B2. One error, B1, but $x$ or $\bar{x}$ or $r$ : B0 Binomial probabilities, allow 1 term only Correct expression [0.0148 + 0.0693] Probability, a.r.t. 0.084 Explicit comparison of “like with like”
	or	Add binomial probs until ans $> 0.1$ Critical region $\leq 1$	A1 B1		$[P(\leq 2) = 0.239]$
		Reject $H_0$ Significant evidence that proportion of $e$ 's in language is less than 0.19	M1 A1 $\sqrt{}$	8	Correct deduction and method [needs $P(\leq 1)$ ] Correct conclusion in context [SR: N(3.8, 3.078): B2M1A0B1M0]
	(ii)	Letters not independent	B1	1	Correct modelling assumption, stated in context Allow “random”, “depends on message”, etc

7	(i)		B1 B1 B1	3	Horizontal straight line Positive parabola, symmetric about 0 Completely correct, including correct relationship between two Don't need vertical lines or horizontal lines outside range, but don't give last B1 if horizontal line continues past "±1"	
	(ii)	$S$ is equally likely to take any value in range, $T$ is more likely at extremities	B2	2	Correct statement about distributions ( <i>not</i> graphs) [Partial statement, or correct description for one only: B1]	
	(iii)	$\int_{-1}^1 \frac{3}{2}x^2 dx = \left[ \frac{x^3}{2} \right]_{-1}^1$ $\frac{1}{2}(1 - (-1)^3) = 0.2$ or $\frac{1}{2}((-1)^3 + 1) = 0.8$ $t^3 = 0.6$ $t = 0.8434$	M1 B1 M1 M1 A1	5	Integrate $f(x)$ with limits $(-1, t)$ or $(t, 1)$ [recoverable if $t$ used later] Correct indefinite integral Equate to 0.2, or 0.8 if $[-1, t]$ used Solve cubic equation to find $t$ Answer, in range $[0.843, 0.844]$	
8	(i)	$\frac{64.2 - 63}{\sqrt{12.25/23}} = 1.644$ $P(z > 1.644) = 0.05$	M1dep A1 dep M1 A1	4	Standardise 64.2 with $\sqrt{12.25}$ $z = 1.644$ or $1.645$ , must be + Find $\Phi(z)$ , answer $< 0.5$ Answer, a.r.t. 0.05 or 5.0%	
	(ii)	(a)	$63 + 1.645 \times \frac{3.5}{\sqrt{50}}$ $\geq 63.81$	M1 B1 A1	3	$63 + 3.5 \times k / \sqrt{50}$ , $k$ from $\Phi^{-1}$ , <i>not</i> $-k = 1.645$ (allow 1.64, 1.65) Answer, a.r.t. 63.8, allow $>$ , $\geq$ , $=$ , c.w.o.
		(b)	$P(< 63.8 \mid \mu = 65)$ $\frac{63.8 - 65}{3.5/\sqrt{50}} = -2.3956$ 0.0083	M1 M1 A1 A1	4	Use of correct meaning of Type II Standardise their $c$ with $\sqrt{50}$ $z = (\pm) 2.40$ [or $-2.424$ or $-2.404$ etc] Answer, a.r.t. 0.008 [eg, 0.00767]
	(iii)	B better: Type II error smaller (and same Type I error)	B2√	2	This answer: B2. "B because sample bigger": B1. [SR: Partial answer: B1]	
9	(a)	$np > 5$ and $nq > 5$ $0.75n > 5$ is relevant $n > 20$	M2 A1	3	Use either $nq > 5$ or $npq > 5$ [SR: If M0, use $np > 5$ , or "n = 20" seen: M1] Final answer $n > 20$ or $n \geq 20$ only	
	(b)	(i)	$70.5 - \mu = 1.75\sigma$ $\mu - 46.5 = 2.25\sigma$  Solve simultaneously $\mu = 60$ $\sigma = 6$	M1 A1 B1 M1 A1√ A1√	6	Standardise once, and equate to $\Phi^{-1}$ , $\pm$ cc Standardise twice, signs correct, cc correct Both 1.75 and 2.25 Correct solution method to get one variable $\mu$ , a.r.t. 60.0 or $\pm 154.5$ $\sigma$ , a.r.t. 6.00 [Wrong cc (below): A1 both] [SR: $\sigma^2$ : M1A0B1M1A1A0]
		(ii)	$np = 60$ , $npq = 36$ $q = 36/60 = 0.6$ $p = 0.4$ $n = 150$	M1dep depM1 A1√ A1√	4	$np = 60$ and $npq = 6^2$ or 6 Solve to get $q$ or $p$ or $n$ $p = 0.4$ √ on wrong cc or $z$ $n = 150$ √ on wrong cc or $z$

$\sigma$	$\mu$	$q$	$p (\pm 0.01)$	$n$
6	60	0.6	0.4	150
	60.062			
6.25	5	0.6504	0.3496	171.8
	60.562			
6.25	5	0.6450	0.3550	170.6
	59.562			
6.25	5	0.6558	0.3442	173.0
6.5	60.125	0.7027	0.2973	202.2
6	59.5	0.6050	0.3950	150.6