






# **GCE MARKING SCHEME**

**CHEMISTRY  
AS/Advanced**

**JANUARY 2013**


**GCE CHEMISTRY - CH4**  
**JANUARY 2013 MARK SCHEME**

**Section A**



(ii) The isomers rotate the plane of polarised light in opposite directions [1]

(iii) Side effects from other optical isomer / lower dose needed / improved pharmacological activity / only one isomer has correct orientation to bind with biological molecule [1]



(1 mark for acid (accept aldehyde), 1 mark for ketone)

(c)(i) Ethylamine, ethanol, phenol, ethanoic acid [1]

(ii) Ethylamine is basic because it accepts a proton readily (1) due to the lone pair of electrons on the nitrogen. (1)  
Phenol is acidic because it loses a proton / the anion formed is stabilised (1) by delocalisation of the negative charge over the benzene ring. (1)  
(Accept description e.g. in phenoxide ion lone pairs of electrons on oxygen become delocalised with electrons in benzene ring.) [4]

Total [14]

2. (a)

	Butan-2-ol	Ethanal	Ethanol	Propanone
2,4-DNP	No reaction	Yellow-orange precipitate	No reaction	Yellow-orange precipitate
Tollens' reagent	No reaction	Silver mirror	No reaction	No reaction
I <sub>2</sub> /NaOH	Yellow precipitate	Yellow precipitate	Yellow precipitate	Yellow precipitate

(1 mark for each column) [4]


(b)(i) Electrophilic addition [1]

(ii) Carbonium ion / carbocation / electrophile [1]

(iii) Bromination / HBr addition / hydrogenation [1]

(iv) Secondary carbocation more stable than primary carbocation [1]

(c)(i) Nucleophilic addition [1]



1 mark electron movement  
1 mark charges

1 mark intermediate  
and electron movement


(Accept  $\text{CN}^{\delta-} - \text{H}^{\delta+}$  for  $\text{CN}^-$ )

Total [12]

3. (a) Intermolecular bond formed (1) when hydrogen attached to a highly electronegative atom (1) is bonded to an electronegative atom attached to hydrogen (in another molecule) (1) forming a very strong dipole – dipole attraction (1) [3]  
(maximum 3 marks)

*QWC* Legibility of text; accuracy of spelling, punctuation and grammar, clarity of meaning [1]

(b)




[1]

- (c) Behaves as / can react with an acid or a base (1)

-COOH is an acidic group / donates proton, -NH<sub>2</sub> is a basic group / accepts proton (1)  
[2]

(d)



[1]

- (e) Moles MSG =  $1/169.08 = 5.91 \times 10^{-3}$  (1)

Concentration =  $5.91 \times 10^{-3} / 0.1 = 5.91 \times 10^{-2}$  (1) [2]

- (f) (Neutral) FeCl<sub>3</sub> / Br<sub>2</sub> (1)

Purple colour / white precipitate (1) [2]

- (g) 2,4-Dinitrophenylhydrazine / acidified sodium dichromate (1)

Yellow-orange precipitate / orange to green colour change (1) [2]

Total [14]

**Total Section A [40]**


**Section B**

4. (a) For synthetic polymer:


Monomers: 1,6-Diaminohexane / ethane-1, 2-diol (1)

Hexanedioic acid / benzene-1,4-dioic acid (1)

Structure:




or



For natural polymer:

Monomers: aminoethanoic acid / 2-aminopropanoic acid (1)

Structure: e.g.



[5]

*QWC* Selection of a form and style of writing appropriate to purpose and to complexity of subject matter [1]




(ii) 1,4-dichlorobenzene [1]

(iii) Chlorine (in the absence of ultraviolet light) (1)  
 $\text{AlCl}_3$  /  $\text{FeCl}_3$  (as a halogen carrier) (1) [2]

(iv) Heat with  $\text{NaOH}$  (aq) (1)  
 add  $\text{HNO}_3$ (aq) followed by  $\text{AgNO}_3$ (aq) (1)  
**F** gives white precipitate, **G** does not (1)  
 In **F**, the C–Cl bond is polarised / contains  $\text{C}^{\delta+}$  or undergoes nucleophilic substitution (1)  
 In **G** due to delocalisation of the  $\pi$  electron cloud of the ring with the p-orbital electrons of the chlorine (1)  
 the C–Cl bond is too strong to break/ does not undergo nucleophilic substitution (1)  
 [6]

*QWC* The information is organised clearly and coherently, using specialist vocabulary where appropriate [1]

(c)(i) To prevent decomposition of benzenediazonium chloride /  $\text{HNO}_2$  [1]




(iii) A chromophore is the group of atoms responsible for the colour of the compound (by causing absorption in the visible region of the spectrum) [1]

Total [20]



5. (a)(i) Moles HCl =  $5.4 \times 10^{-3}$  (1)  
 $M_r \text{ B} = \frac{0.395}{0.0054} = 73.1$  (1) [2]

- (ii) **B** is basic therefore must be amine (1)  
**C** reacts with  $\text{Na}_2\text{CO}_3$  therefore must be an acid (1)  
**D** is oxidised to **C** therefore must be an alcohol (1)  
**A** hydrolyses to acid but does not contain oxygen therefore must be nitrile (1)  
**B** is  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{NH}_2$  (1)  
**C** is  $\text{CH}_3\text{CH}_2\text{CH}_2\text{COOH}$  (1)  
**D** is  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$  (1)  
**A** is  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CN}$  (1) [8]  
 (4 marks structures – if 3 carbons in chains penalise only once  
 4 marks reasons – accept alternative reasons)




Peak areas (1)

Chemical shifts (1)



Peak areas (1)

Chemical shifts (1)



- (c)(i) 2 steps instead of 3 /  $\text{CH}_3\text{COCH}_3$  can be sold / reagents are cheaper / gives a higher yield / easier to extract phenol / phenol formed more quickly / fewer reactants [2]  
 (Accept any 2)

- (ii) Lower temperature required / catalyst costs less / catalyst less likely to break up / catalyst less toxic or safer [1]

(d) Moles phenol =  $58.75/94.06 = 0.625$  (1)  
 Maximum mass aspirin =  $0.625 \times 180.08 = 112.55 \text{ g}$  (1)  
 65% yield, therefore mass aspirin = 73.16 g (1) [3]

Total [20]

**Total Section B [40]**